

## 2001 Paper 10 Question 10

### Continuous Mathematics

- (a) The MacLaurin series for a continuous, infinitely differentiable function,  $f(x)$ , is:

$$f(x) = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots + \frac{f^{(k)}(0)}{k!}x^k + \dots$$

Derive the MacLaurin series for each of  $\sin(x)$ ,  $\cos(x)$ , and  $e^x$ . [6 marks]

- (b) Hence, or otherwise, prove that:

$$e^{i\phi} = \cos \phi + i \sin \phi$$

where  $i = \sqrt{-1}$  [3 marks]

- (c) Prove that the box function,  $b(x)$ :

$$b(x) = \begin{cases} 1, & |x| \leq \frac{1}{2} \\ 0, & \text{otherwise} \end{cases}$$

has the Fourier transform,  $B(\nu)$ :

$$B(\nu) = \frac{\sin \pi\nu}{\pi\nu}$$

where  $\nu$  is frequency measured in Hertz (cycles per second). [7 marks]

- (d) The convolution of  $b(x)$  with itself is  $t(x)$ :

$$t(x) = b(x) * b(x) = \begin{cases} 1+x, & -1 \leq x \leq 0 \\ 1-x, & 0 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

Hence, or otherwise, find the Fourier transform,  $T(\nu)$ , of  $t(x)$ . [4 marks]