

MCMC methods

A simple technique is to introduce a random walk, so

$$\mathbf{w}_{i+1} = \mathbf{w}_i + \epsilon$$

where ϵ is zero mean spherical Gaussian and has small variance. Obviously the sequence \mathbf{w}_i does not have the required distribution. However, we can use the *Metropolis algorithm*, which does *not* accept all the steps in the random walk:

1. If $p(\mathbf{w}_{i+1}|\mathbf{y}) > p(\mathbf{w}_i|\mathbf{y})$ then accept the step.
2. Else accept the step with probability $\frac{p(\mathbf{w}_{i+1}|\mathbf{y})}{p(\mathbf{w}_i|\mathbf{y})}$.

In practice, the Metropolis algorithm has several shortcomings, and a great deal of research exists on improved methods, see:

R. Neal, "Probabilistic inference using Markov chain Monte Carlo methods," University of Toronto, Department of Computer Science Technical Report CRG-TR-93-1, 1993.

Approximate inference for Bayesian networks

MCMC methods also provide a method for performing *approximate inference* in *Bayesian networks*.

Say a system can be in a state s and moves from state to state in discrete time steps according to a probabilistic transition

$$\Pr(s \rightarrow s')$$

Let $\pi_t(s)$ be the probability distribution for the state after t steps, so

$$\pi_{t+1}(s') = \sum_s \Pr(s \rightarrow s') \pi_t(s)$$

If at some point we obtain $\pi_{t+1}(s) = \pi_t(s)$ for all s then we have reached a *stationary distribution* π . In this case

$$\forall s' \pi(s') = \sum_s \Pr(s \rightarrow s') \pi(s)$$

There is exactly one stationary distribution for a given $\Pr(s \rightarrow s')$ provided the latter obeys some simple conditions.

Approximate inference for Bayesian networks

The condition of *detailed balance*

$$\forall s, s' \pi(s) \Pr(s \rightarrow s') = \pi(s') \Pr(s' \rightarrow s)$$

is sufficient to provide a π that is a stationary distribution. To see this simply sum:

$$\begin{aligned} \sum_s \pi(s) \Pr(s \rightarrow s') &= \sum_s \pi(s') \Pr(s' \rightarrow s) \\ &= \pi(s') \underbrace{\sum_s \Pr(s' \rightarrow s)}_{=1} \\ &= \pi(s') \end{aligned}$$

If all this is looking a little familiar, it's because we now have an excellent application for the material in *Mathematical Methods for Computer Science*. That course used the alternative term *local balance*.

Approximate inference for Bayesian networks

Recalling once again the basic equation for performing probabilistic inference

$$\Pr(Q|e) = \frac{1}{Z} \Pr(Q \wedge e) = \frac{1}{Z} \sum_u \Pr(Q, u, e)$$

where

- Q is the query variable.
- e is the evidence.
- u are the unobserved variables.
- $1/Z$ normalises the distribution.

We are going to consider obtaining samples from the distribution $\Pr(Q, U|e)$.

Approximate inference for Bayesian networks

The evidence is fixed. Let the *state* of our system be a specific set of values for the *query variable and the unobserved variables*

$$\mathbf{s} = (q, u_1, u_2, \dots, u_n) = (s_1, s_2, \dots, s_{n+1})$$

and define \bar{s}_i to be the state vector *with s_i removed*

$$\bar{s}_i = (s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_{n+1})$$

To move from \mathbf{s} to \mathbf{s}' we replace one of its elements, say s_i , with a new value s'_i sampled according to

$$s'_i \sim \Pr(S_i | \bar{s}_i, e)$$

This has detailed balance, and has $\Pr(Q, U|e)$ as its stationary distribution.

5

Approximate inference for Bayesian networks

To see that $\Pr(Q, U|e)$ is the stationary distribution

$$\begin{aligned} \pi(\mathbf{s})\Pr(\mathbf{s} \rightarrow \mathbf{s}') &= \Pr(\mathbf{s}|e)\Pr(s'_i | \bar{s}_i, e) \\ &= \Pr(s_i, \bar{s}_i | e)\Pr(s'_i | \bar{s}_i, e) \\ &= \Pr(s_i | \bar{s}_i, e)\Pr(\bar{s}_i | e)\Pr(s'_i | \bar{s}_i, e) \\ &= \Pr(s_i | \bar{s}_i, e)\Pr(s'_i, \bar{s}_i | e) \\ &= \Pr(\mathbf{s}' \rightarrow \mathbf{s})\pi(\mathbf{s}') \end{aligned}$$

As a further simplification, sampling from $\Pr(S_i | \bar{s}_i, e)$ is equivalent to sampling S_i conditional on its parents, children and children's parents.

6

Approximate inference for Bayesian networks

So:

- We successively sample the query variable and the unobserved variables, conditional on their parents, children and children's parents.
- This gives us a sequence s_1, s_2, \dots which has been sampled according to $\Pr(Q, U|e)$.

Finally, note that as

$$\Pr(Q|e) = \sum_u \Pr(Q, u|e)$$

we can just ignore the values obtained for the unobserved variables. This gives us q_1, q_2, \dots with

$$q_i \sim \Pr(Q|e)$$

7

Approximate inference for Bayesian networks

To see that the final step works, consider what happens when we estimate the expected value of some function of Q .

$$\begin{aligned} \mathbb{E}[f(Q)] &= \sum_q f(q)\Pr(q|e) \\ &= \sum_q f(q) \sum_u \Pr(q, u|e) \\ &= \sum_q \sum_u f(q)\Pr(q, u|e) \end{aligned}$$

so sampling using $\Pr(q, u|e)$ and ignoring the values for u obtained works exactly as required.

8