

# *Mathematical Methods for Computer Science*



UNIVERSITY OF  
CAMBRIDGE

Computer Laboratory

---

Computer Science Tripos, Part IB

Michaelmas Term 2011/12

R.J. Gibbens

Problem sheet  
Part B: Probability methods

William Gates Building  
15 JJ Thomson Avenue  
Cambridge  
CB3 0FD

<http://www.cl.cam.ac.uk/>

### §1: Limits and inequalities

1. Suppose that  $X$  is a random variable with the  $U(-1, 1)$  distribution. Find the exact value of  $\mathbb{P}(|X| \geq a)$  for each  $a > 0$  and compare it to the upper bounds obtained from the Markov and Chebychev inequalities.
2. Let  $X$  be the random variable giving the number of heads obtained in a sequence of  $n$  independent fair coin flips. Compare the upper bounds on  $\mathbb{P}(X \geq 3n/4)$  obtained from the Markov and Chebychev inequalities.
3. Let  $A_i$  ( $i = 1, 2, \dots, n$ ) be a collection of random events and set  $N = \sum_{i=1}^n \mathbb{I}(A_i)$ . By considering Markov's inequality applied to  $\mathbb{P}(N \geq 1)$  show Boole's inequality, namely,

$$\mathbb{P}(\cup_{i=1}^n A_i) \leq \sum_{i=1}^n \mathbb{P}(A_i).$$

4. Let  $h : \mathbb{R} \rightarrow [0, \infty)$  be a non-negative function. Show that

$$\mathbb{P}(h(X) \geq a) \leq \frac{\mathbb{E}(h(X))}{a} \quad \text{for all } a > 0.$$

By making suitable choices of  $h(x)$ , show that we may obtain the Markov and Chebychev inequalities as special cases.

5. Show the following properties of the moment generating function.
  - (a) If  $X$  has mgf  $M_X(t)$  then  $Y = aX + b$  has mgf  $M_Y(t) = e^{bt}M_X(at)$ .
  - (b) If  $X$  and  $Y$  are independent then  $X + Y$  has mgf  $M_{X+Y}(t) = M_X(t)M_Y(t)$ .
  - (c)  $\mathbb{E}(X^n) = M_X^{(n)}(0)$  where  $M_X^{(n)}$  is the  $n^{\text{th}}$  derivative of  $M_X$ .
  - (d) If  $X$  is a discrete random variable taking values  $0, 1, 2, \dots$  with probability generating function  $G_X(z) = \mathbb{E}(z^X)$  then  $M_X(t) = G_X(e^t)$ .
6. Let  $X$  be a random variable with moment generating function  $M_X(t)$  which you may assume exists for any value of  $t$ . Show that for any  $a > 0$

$$\mathbb{P}(X \leq a) \leq e^{-ta}M_X(t) \quad \text{for all } t < 0.$$

7. Show that, if  $X_n \xrightarrow{D} X$ , where  $X$  is a degenerate random variable (that is,  $\mathbb{P}(X = \mu) = 1$  for some constant  $\mu$ ) then  $X_n \xrightarrow{P} X$ .
8. Suppose that you estimate your monthly phone bill by rounding all amounts to the nearest pound. If all rounding errors are independent and distributed as  $U(-0.5, 0.5)$ , estimate the probability that the total error exceeds one pound when your bill has 12 items. How does this procedure suggest an approximate method for constructing Normal random variables?
9. 2006 Paper 3 Question 10
10. 2011 Paper 6 Question 7

## §2: Markov chains

1. Suppose that  $(X_n)$  is a Markov chain with  $n$ -step transition matrix,  $P^{(n)}$ , and let  $\lambda_i^{(n)} = \mathbb{P}(X_n = i)$  be the elements of a row vector  $\lambda^{(n)}$  ( $n = 0, 1, 2, \dots$ ). Show that

(a)  $P^{(m+n)} = P^{(m)}P^{(n)}$  for  $m, n = 0, 1, 2, \dots$

(b)  $\lambda^{(n)} = \lambda^{(0)}P^{(n)}$  for  $n = 0, 1, 2, \dots$

2. Suppose that  $(X_n)$  is a Markov chain with transition matrix  $P$ . Define the relations “state  $j$  is accessible from state  $i$ ” and “states  $i$  and  $j$  communicate”. Show that the second relation is an equivalence relation and define the communicating classes as the equivalence classes under this relation. What is meant by the terms *closed class*, *absorbing class* and *irreducible*?

3. Show that

$$P_{ij}(z) = \delta_{ij} + F_{ij}(z)P_{jj}(z)$$

where

$$P_{ij}(z) = \sum_{n=0}^{\infty} p_{ij}^{(n)} z^n, \quad F_{ij}(z) = \sum_{n=0}^{\infty} f_{ij}^{(n)} z^n$$

and  $\delta_{ij} = 1$  if  $i = j$  and 0 otherwise. [You should assume that  $p_{ij}^{(n)}$  and  $f_{ij}^{(n)}$  are as defined in lectures with  $p_{ij}^{(0)} = \delta_{ij}$  and  $f_{ij}^{(0)} = 0$  for all states  $i, j$ .]

4. Suppose that  $(X_n)$  is a finite state Markov chain and that for some state  $i$  and for all states  $j$

$$\lim_{n \rightarrow \infty} p_{ij}^{(n)} = \pi_j$$

for some collection of numbers  $(\pi_j)$ . Show that  $\pi = (\pi_j)$  is a stationary distribution.

5. Consider the Markov chain with transition matrix

$$P = \begin{pmatrix} 0.128 & 0.872 \\ 0.663 & 0.337 \end{pmatrix}$$

for Markov’s example of a chain on the two states {vowel, consonant} for consecutive letters in a passage of text. Find the stationary distribution for this Markov chain. What are the mean recurrence times for the two states?

6. Define what is meant by saying that  $(X_n)$  is a reversible Markov chain and write down the local balance conditions. Show that if a vector  $\pi$  is a distribution over the states of the Markov chain that satisfies the local balance conditions then it is a stationary distribution.

7. Consider the Ehrenfest model for  $m$  balls moving between two containers with transition matrix

$$p_{i,i+1} = 1 - \frac{i}{m}, \quad p_{i,i-1} = \frac{i}{m}$$

where  $i$  ( $0 \leq i \leq m$ ) is the number of balls in a given container. Show that the Markov chain is irreducible and periodic with period 2. Derive the stationary distribution.

8. Consider a random walk,  $(X_n)$ , on the states  $i = 0, 1, 2, \dots$  with transition matrix

$$\begin{aligned}p_{i,i-1} &= p & i = 1, 2, \dots \\p_{i,i+1} &= 1 - p & i = 0, 1, \dots \\p_{0,0} &= p\end{aligned}$$

where  $0 < p < 1$ . Show that the Markov chain is irreducible and aperiodic. Find a condition on  $p$  to make the Markov chain positive recurrent and find the stationary distribution in this case.

9. Describe PageRank as a Markov chain model for the motion between nodes in a graph. Explain the main mathematical results that underpin PageRank's connection to a notion of web page "*importance*".
10. 2007 Paper 4 Question 5
11. 2010 Paper 6 Question 8
12. 2011 Paper 6 Question 8 NB The correct formula in (d) is:  $\sin(\theta) = (e^{i\theta} - e^{-i\theta})/2i$ .