

# *Mathematical Methods for Computer Science*



UNIVERSITY OF  
CAMBRIDGE

Computer Laboratory

---

Computer Science Tripos Part IB

Michaelmas Term 2012/13

Professor J. Daugman

Problem sheet  
Part A: Fourier and related methods

15 JJ Thomson Avenue  
Cambridge CB3 0FD

<http://www.cl.cam.ac.uk/>

## Part A: Fourier and related methods

1. Given a complex linear space,  $V$ , define the notion of an *inner product* and in the case of  $V = \mathbb{C}^n$  show that for any two vectors  $x, y \in \mathbb{C}^n$

$$\langle x, y \rangle = \sum_{i=1}^n x_i \bar{y}_i$$

where  $x = (x_1, x_2, \dots, x_n)$  and  $y = (y_1, y_2, \dots, y_n)$  defines an inner product.

2. Suppose that  $V$  is a complex inner product space. Show the *Cauchy-Schwarz inequality*, namely, that for all  $u, v \in V$

$$|\langle u, v \rangle|^2 \leq \langle u, u \rangle \langle v, v \rangle.$$

Define the notion of a *norm* for  $V$  and show that

$$\|v\| = \sqrt{\langle v, v \rangle}$$

is a norm.

3. Calculate the Fourier series of the function  $f(x)$  ( $x \in [-\pi, \pi]$ ) defined by

$$f(x) = \begin{cases} 1 & 0 \leq x < \pi \\ 0 & -\pi \leq x < 0. \end{cases}$$

Find also the complex Fourier series for  $f(x)$ .

4. Suppose that  $f(x)$  is a  $2\pi$ -periodic function with complex Fourier series

$$\sum_{n=-\infty}^{\infty} c_n e^{inx}.$$

Now consider the shifted version of  $f(x)$  given by

$$g(x) = f(x - x_0)$$

where  $x_0$  is a constant. Find the relationship between the complex Fourier coefficients of  $g(x)$  in terms of those of  $f(x)$ . How do the magnitudes of the corresponding coefficients compare?

5. Suppose that  $f(x)$  and  $g(x)$  are two functions defined for real  $x$  and that they have Fourier transforms  $F(\omega)$  and  $G(\omega)$ , respectively. Show that

$$\int_{-\infty}^{\infty} f(x)G(x)dx = \int_{-\infty}^{\infty} F(x)g(x)dx.$$

You may assume that the above integrals exist and that you may change the order of integration in your calculations.

6. Consider the functions  $f_b(x)$  and  $g(x)$  defined by

$$f_b(x) = \begin{cases} 0 & x > b \\ 1 & -b < x \leq b \\ 0 & x \leq -b \end{cases}$$

where  $b > 0$  is a constant and

$$g(x) = \begin{cases} 0 & x > 4 \\ 1 & 3 < x \leq 4 \\ 1.5 & 2 < x \leq 3 \\ 1 & 1 < x \leq 2 \\ 0 & x \leq 1. \end{cases}$$

Use the Fourier transform of  $f_b(x)$  (derived in lectures) together with properties of Fourier transforms (which you should state carefully) to construct the Fourier transform of  $g(x)$ .

7. Suppose that the  $N$ -point DFT of the sequence  $f[n]$  is given by  $F[k]$  where  $f(n)$  is itself a  $N$ -periodic sequence, that is  $f(n + N) = f(n)$  for  $n = 0, 1, \dots, N - 1$ . Show that the shifted sequence  $f[n - m]$  has DFT

$$e^{-2\pi imk/N} F[k]$$

where  $m$  is a constant integer. Show also that  $\overline{f[n]}$ , the complex conjugate of  $f[n]$ , has DFT  $\overline{F[-k]}$ . Suppose that  $f[-2] = -1$ ,  $f[-1] = -2$ ,  $f[0] = 0$ ,  $f[1] = 2$ ,  $f[2] = 1$ . Find the 5-point DFT of  $f[n]$ . Can you explain why it is purely imaginary?

8. Suppose that the sequences  $f[n]$  and  $g[n]$  have  $N$ -point DFTs given by  $F[k]$  and  $G[k]$ , respectively. By expanding  $F[k]G[k]$  show that the *cyclical convolution*

$$\sum_{m=0}^{N-1} f[m]g[n - m]$$

has DFT  $F[k]G[k]$ .

9. 2006 Paper 4 Question 6  
 10. 2007 Paper 3 Question 6  
 11. 2008 Paper 3 Question 5  
 12. 2012 Paper 6 Question 7