# CST 2016 Part IA Discrete Mathematics Formal Languages and Automata Exercise Sheet

### **1** Inductive definitions

**Exercise 1.1.** Let *L* be the subset of  $\{a, b\}^*$  inductively defined by the axiom  $-\frac{\varepsilon}{\varepsilon}$  and the

rules  $\frac{u}{aub}$  (one rule for each  $u \in \{a, b\}^*$ ).

- (a) Use rule induction to prove that every string in *L* is of the form  $a^n b^n$  for some  $n \in \mathbb{N}$ .
- (b) Use mathematical induction to prove  $\forall n \in \mathbb{N}$ .  $a^n b^n \in L$ .

Conclude that  $L = \{a^n b^n \mid n \in \mathbb{N}\}.$ 

**Exercise 1.2.** Suppose  $R \subseteq X \times X$  is a binary relation on a set *X*. Let  $R^{\dagger} \subseteq X \times X$  be inductively defined by the following axioms and rules:

$$\frac{(x,y)\in R^{\dagger}}{(x,x)\in R^{\dagger}} \ (x\in X), \qquad \frac{(x,y)\in R^{\dagger}}{(x,z)\in R^{\dagger}} \ (x\in X \text{ and } (y,z)\in R).$$

- (a) Show that  $R^{\dagger}$  is reflexive and that  $R \subseteq R^{\dagger}$ .
- (b) Use rule induction to show that  $R^{\dagger}$  is a subset of  $\{(y, z) \in X \times X \mid \forall x \in X. (x, y) \in R^{\dagger} \Rightarrow (x, z) \in R^{\dagger}\}$ . Deduce that  $R^{\dagger}$  is transitive.
- (c) Suppose  $S \subseteq X \times X$  is a reflexive and transitive binary relation and that  $R \subseteq S$ . Use rule induction to show that  $R^{\dagger} \subseteq S$ .
- (d) Deduce from (a)–(c) that  $R^+$  is equal to  $R^*$ , the reflexive-transitive closure of R.

**Exercise 1.3.** Let *L* be the subset of  $\{a, b\}^*$  inductively defined by the axiom  $-\frac{ab}{ab}$  and the

rules  $\frac{au}{au^2}$  and  $\frac{ab^3u}{au}$  (for all  $u \in \{a, b\}^*$ ).

- (a) Is  $ab^5$  in *L*? Give a derivation, or show there isn't one.
- (b) Use rule induction to show that every  $u \in L$  is of the form  $ab^n$  with  $n = 2^k 3m \ge 0$  for some  $k, m \in \mathbb{N}$ .
- (c) Is  $ab^3$  in L? Give a derivation, or show there isn't one.
- (d) Can you characterize exactly which strings are in *L*?

#### 2 Regular expressions

**Exercise 2.1.** Find regular expressions over  $\{0, 1\}$  that determine the following languages:

- (a)  $\{u \mid u \text{ contains an even number of } 1's\}$
- (b)  $\{u \mid u \text{ contains an odd number of } 0's\}$

**Exercise 2.2.** Show that the inductive definition of the matching relation between strings and regular expressions given on Slide 36 of the lecture notes has the properties listed on Slide 35.

**Exercise 2.3.** Show that  $b^*a(b^*a)^*$  and  $(a|b)^*a$  are equivalent regular expressions, that is, a string matches one iff it matches the other.

Tripos questions 2012.2.8(a-d) 2005.2.1(d) 1999.2.1(s) 1997.2.1(q) 1996.2.1(i)

# 3 Finite automata

**Exercise 3.1.** For each of the two languages mentioned in Exercise **??** find a DFA that accepts exactly that set of strings.

**Exercise 3.2.** Given an NFA<sup> $\varepsilon$ </sup>  $M = (Q, \Sigma, \Delta, s, F, T)$ , we write  $q \stackrel{u}{\Rightarrow} q'$  to mean that there is a path in M from state q to state q' whose non- $\varepsilon$  labels form the string  $u \in \Sigma^*$ . Show that  $\{(q, u, q') \mid q \stackrel{u}{\Rightarrow} q'\}$  is equal to the subset of  $Q \times \Sigma^* \times Q$  inductively defined by the axioms and rules

$$(q, \varepsilon, q)$$

$$\frac{(q, u, q')}{(q, u, q'')} \text{ if } q' \xrightarrow{\varepsilon} q'' \text{ in } M$$

$$\frac{(q, u, q')}{(q, ua, q'')} \text{ if } q' \xrightarrow{a} q'' \text{ in } M.$$

**Exercise 3.3.** The example of the subset construction given in the lecture notes constructs a DFA with eight states whose language of accepted strings happens to be  $L(a^*b^*)$ . Give a DFA with the same language of accepted strings, but fewer states. Give an NFA with even fewer states that does the same job.

**Tripos questions** 2010.2.9 2009.2.9 2004.2.1(d) 2001.2.1(d) 2000.2.1(b) 1998.2.1(s) 1995.2.19

# 4 Regular Languages

**Exercise 4.1.** Why can't the automaton Star(M) used in step (iv) of the proof of part (a) of Kleene's Theorem be constructed simply by taking *M*, making its start state the only accepting state and adding new  $\varepsilon$ -transitions back from each old accepting state to its start state?

**Exercise 4.2.** Construct an NFA<sup> $\varepsilon$ </sup> *M* satisfying  $L(M) = L((\varepsilon|b)^*aab^*)$ .

Exercise 4.3. Show that any finite set of strings is a regular language.

**Exercise 4.4.** Use the construction given in the proof of part (b) of Kleene's Theorem to find a regular expression for the DFA *M* whose state set is  $\{0, 1, 2\}$ , whose start state is 0, whose only accepting state is 2, whose alphabet of input symbols is  $\{a, b\}$ , and whose next-state function is given by the following table.

δ	а	b
0	1	2
1	2	1
2	2	1

**Exercise 4.5.** If  $M = (Q, \Sigma, \Delta, s, F)$  is an NFA, let Not(M) be the NFA  $(Q, \Sigma, \Delta, s, Q \setminus F)$  obtained from M by interchaning the role of accepting and non-accepting states. Give an example of an alphabet  $\Sigma$  and an NFA M with set of input symbols  $\Sigma$ , such that  $\{u \in \Sigma^* \mid u \notin L(M)\}$  is *not* the same set as L(Not(M)).

**Exercise 4.6.** Let  $r = (a|b)^* ab(a|b)^*$ . Find a complement for r over the alphabet  $\{a, b\}$ , i.e. a regular expressions  $\sim r$  over the alphabet  $\{a, b\}$  satisfying  $L(\sim r) = \{u \in \{a, b\}^* \mid u \notin L(r)\}$ .

**Exercise 4.7.** Given DFAs  $M_i = (Q_i, \Sigma, \delta_i, s_i, F_i)$  for i = 1, 2, let  $And(M_1, M_2)$  be the DFA  $(Q_1 \times Q_2, \Sigma, \delta, (s_1, s_2), F_1 \times F_2)$  where  $\delta : (Q_1 \times Q_2) \times \Sigma \to (Q_1 \times Q_2)$  is given by:

 $\delta((q_1, q_2), a) = (\delta_1(q_1, a), \delta_2(q_2, a))$ 

for all  $q_1 \in Q_1$ ,  $q_2 \in Q_2$  and  $a \in \Sigma$ . Show that  $L(And(M_1, M_2))$  is the intersection of  $L(M_1)$  and  $L(M_2)$ .

Tripos questions 2003.2.9 2000.2.7 1995.2.20

### 5 The Pumping Lemma

**Exercise 5.1.** Show that there is no DFA M for which L(M) is the language on Slide 107. [Hint: argue by contradiction. If there were such an M, consider the DFA M' with the same states as M, with alphabet of input symbols just consisting of a and b, with transitions all those of M which are labelled by a or b, with start state  $\delta_M(s_M, c)$  (where  $s_M$  is the start state of M), and with the same accepting states as M. Show that the language accepted by M' has to be  $\{a^n b^n \mid n \ge 0\}$  and deduce that no such M can exist.]

**Tripos questions** 2011.2.8 2006.2.8 2004.2.9 2002.2.9 2001.2.7 1999.2.7 1998.2.7 1996.2.1(j) 1996.2.8 1995.2.27