

# 1995 Paper 13 Question 14

## Numerical Analysis II

With reference to solution of the differential equation  $y' = f(x, y)$ , explain the conventional notation  $x_n, y(x_n), y_n, f_n$ . [4 marks]

Derive Euler's method

$$y_{n+1} = y_n + hf(x_n, y_n). \quad (1)$$

[3 marks]

Euler's method has *local error*

$$\frac{h^2}{2}y''(\xi).$$

Explain the terms *local error*, *global error*.

[2 marks]

The multistep formula

$$y_{n+1} = y_{n-3} + \frac{4h}{3}(2f_n - f_{n-1} + 2f_{n-2}) \quad (2)$$

has *local error*

$$\frac{14}{45}h^5y^{(5)}(\xi).$$

Outline the technique for deriving multistep formulae such as (2). (Omit algebraic details.) [2 marks]

Suppose Euler's formula is used as a starting procedure for formula (2). How many initial steps of formula (2) need to be evaluated using Euler? [2 marks]

Estimate very roughly the number  $N$  of Euler steps needed to approximate  $f_1$ . (Assume that  $|y^{(5)}(x)| \simeq 30$ , and Euler's method has *global error*  $h/N$ .) [5 marks]

What is the most important requirement for a starting procedure? Suggest a more suitable starting procedure than Euler. [2 marks]