

## 2005 Paper 8 Question 12

### Numerical Analysis II

The best  $L_\infty$  approximation to  $f(x) \in C[-1, 1]$  by a polynomial  $p_{n-1}(x)$  of degree  $n - 1$  has the property that

$$\max_{x \in [-1, 1]} |e(x)|$$

is attained at  $n + 1$  distinct points  $-1 \leq \xi_0 < \xi_1 < \dots < \xi_n \leq 1$  such that  $e(\xi_j) = -e(\xi_{j-1})$  for  $j = 1, 2, \dots, n$  where  $e(x) = f(x) - p_{n-1}(x)$ .

- (a) Let  $f(x) = x^2$ . Show, by means of a clearly labelled sketch graph, that the best polynomial approximation of degree 1 is a constant. [3 marks]
- (b) Now suppose  $f(x) = (x + 1)/(x + \frac{5}{3})$  is the function to be approximated over  $[-1, 1]$ . By sketching the graph, deduce properties of the best linear approximation  $p_1(x)$ . By differentiating  $e(x)$ , find  $p_1(x)$ . [9 marks]
- (c) Now consider  $f(x) = x/(9x^2 + 16)$ . Explain why the best approximation over  $[-1, 1]$  of degree 2 or less is of the form  $p_2(x) = ax$ , and sketch the graph to show the extreme values of  $e(x)$ . Verify that  $x = 4/9$  is one of the extreme values and find  $a$ . [8 marks]