

2000 Paper 13 Question 10

Numerical Analysis II

Explain the terms (a) *positive definite*, (b) *positive semi-definite* for a symmetric matrix \mathbf{A} . If a square matrix \mathbf{B} is non-singular, which of the properties (a) or (b) most accurately describes $\mathbf{B}^T \mathbf{B}$? What if \mathbf{B} is singular? [4 marks]

State *Schwarz's inequality* for the product $\mathbf{A}\mathbf{B}$. In what way is this modified for the product $\mathbf{A}\mathbf{x}$, where \mathbf{x} is a vector? What are the *singular values* of \mathbf{A} , and how are they related to the l_2 norm of \mathbf{A} ? In the *singular value decomposition* $\mathbf{A} = \mathbf{U}\mathbf{W}\mathbf{V}^T$, what is \mathbf{W} ? [5 marks]

Let $\hat{\mathbf{x}}$ be an approximate solution of $\mathbf{A}\mathbf{x} = \mathbf{b}$, and write $\mathbf{r} = \mathbf{b} - \mathbf{A}\hat{\mathbf{x}}$, $\mathbf{e} = \mathbf{x} - \hat{\mathbf{x}}$. Find an expression which is an upper bound for the relative error $\|\mathbf{e}\|/\|\mathbf{x}\|$ in terms of computable quantities. Explain how this result may be interpreted if the l_2 norm is used. [8 marks]

Suppose \mathbf{A} is a 5×5 matrix and $\mathbf{A}\mathbf{x} = \mathbf{b}$ is to be solved by singular value decomposition. If *machine epsilon* $\simeq 10^{-15}$ and the singular values of \mathbf{A} are $1, 10^{-6}, 10^{-10}, 10^{-17}, 0$ write down the generalised inverse \mathbf{W}^+ that you would use. [3 marks]