

4 Complexity Theory (mpf23)

- (a) For a complexity class \mathcal{C} , let $\text{co-}\mathcal{C} = \{\bar{L} \mid L \in \mathcal{C}\}$ and say that \mathcal{C} is closed under complementation whenever $\mathcal{C} = \text{co-}\mathcal{C}$.

Argue as to whether the following statements are true, false, or unknown.

- (i) All deterministic time complexity classes are closed under complementation. [3 marks]

- (ii) All non-deterministic time complexity classes are closed under complementation. [3 marks]

- (b) For a mapping $f : \Sigma \rightarrow \Sigma$ on an alphabet Σ and a language $L \subseteq \Sigma^*$, define $f[L] = \{f^n(w) \in \Sigma^* \mid w \in L\}$ where $f^n(a_1 \cdots a_n) = f(a_1) \cdots f(a_n)$.

Prove that $L \in \text{NP}$ implies $f[L] \in \text{NP}$. [4 marks]

- (c) Consider the following decision problem.

Q: Given natural numbers m and n in \mathbb{N} , and bits $a_{i,j}^{(k)}$ and b_k in $\{0, 1\}$ for $1 \leq k \leq m$ and $1 \leq i, j \leq n$, determine whether the system of equations $\sum_{1 \leq i, j \leq n} a_{i,j}^{(k)} x_i x_j = b_k$ ($1 \leq k \leq m$) with unknowns x_1, \dots, x_n has a solution in arithmetic modulo 2.

- (i) Prove that Q is in NP. [3 marks]

- (ii) By means of a polynomial-time reduction from the problem 3CNF, or otherwise, prove that Q is NP-hard. [Hint: Note, for instance, that $x = \neg y$ in the Boolean algebra $\{0, 1\}$ if, and only if, $xx + yy = 1$ in arithmetic modulo 2.] [7 marks]

You may use standard results provided that you state them clearly.