

8 Data Science (djw1005)

(a) In a COVID vaccine trial,  $n_0$  subjects were given a placebo and  $n_1$  were given the vaccine;  $x_0$  of the placebo subjects developed the disease and  $x_1$  of the vaccinated subjects. Considering the probability model  $X_k \sim \text{Binomial}(n_k, p_k)$ , the vaccine efficacy is defined to be  $e = 1 - p_1/p_0$ .

(i) State the maximum likelihood estimators for  $p_0$  and  $p_1$ . Give a formula for the maximum likelihood estimator for  $e$ . [2 marks]

(ii) Explain how to compute a 95% confidence interval for  $e$ . Also explain how to test whether  $e > 0.5$ . [7 marks]

(b) Further data about the trial has been made available, and we learn that subjects weren't all enrolled for the same length of time. We are given a full dataset consisting of three features, the predictor variable  $d_i$  and the response variables  $(t_i, c_i)$  for subject  $i$ . Here  $d_i = 1$  if the subject received the vaccine and  $d_i = 0$  otherwise;  $c_i = 1$  if the subject developed the disease and  $c_i = 0$  otherwise; and  $t_i$  is the day on which the subject developed the disease if  $c_i = 1$ , and the number of days enrolled in the trial otherwise.

Consider the following probability model. Among vaccinated subjects, the vaccine is effective with probability  $f$  and ineffective otherwise. Effectively vaccinated subjects never get the disease. For ineffectively vaccinated subjects, and for subjects on placebo, each day there is a probability  $q$  of developing the disease. The parameters  $f$  and  $q$  are unknown.

(i) For a subject  $i$  who received placebo, give an expression for the likelihood of the pair  $(t_i, c_i)$ . [4 marks]

(ii) For a subject  $i$  who received the vaccine, give an expression for the likelihood of the pair  $(t_i, c_i)$ . [4 marks]

(iii) Give an expression for the log likelihood of the entire dataset. [3 marks]

[Note: Your answers to this question should be symbolic. But you may like to know that in the low-dose part of the Oxford–AstraZeneca trial,  $n_0 = 1374$ ,  $n_1 = 1367$ ,  $x_0 = 30$ , and  $x_1 = 3$ .]