



---

On an Interpretation of Second Order Quantification in First Order Intuitionistic Propositional Logic

Author(s): Andrew M. Pitts

Source: *The Journal of Symbolic Logic*, Vol. 57, No. 1 (Mar., 1992), pp. 33-52

Published by: Association for Symbolic Logic

Stable URL: <https://www.jstor.org/stable/2275175>

Accessed: 20-12-2018 15:33 UTC

---

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



*Association for Symbolic Logic* is collaborating with JSTOR to digitize, preserve and extend access to *The Journal of Symbolic Logic*

JSTOR

## ON AN INTERPRETATION OF SECOND ORDER QUANTIFICATION IN FIRST ORDER INTUITIONISTIC PROPOSITIONAL LOGIC

ANDREW M. PITTS

**Abstract.** We prove the following surprising property of Heyting's intuitionistic propositional calculus, IpC. Consider the collection of formulas,  $\phi$ , built up from propositional variables ( $p, q, r, \dots$ ) and falsity ( $\perp$ ) using conjunction ( $\wedge$ ), disjunction ( $\vee$ ) and implication ( $\rightarrow$ ). Write  $\vdash \phi$  to indicate that such a formula is intuitionistically valid. We show that for each variable  $p$  and formula  $\phi$  there exists a formula  $A_p \phi$  (effectively computable from  $\phi$ ), containing only variables not equal to  $p$  which occur in  $\phi$ , and such that for all formulas  $\psi$  not involving  $p$ ,  $\vdash \psi \rightarrow A_p \phi$  if and only if  $\vdash \psi \rightarrow \phi$ . Consequently quantification over propositional variables can be modelled in IpC, and there is an interpretation of the second order propositional calculus, IpC<sup>2</sup>, in IpC which restricts to the identity on first order propositions.

An immediate corollary is the strengthening of the usual interpolation theorem for IpC to the statement that there are least and greatest interpolant formulas for any given pair of formulas. The result also has a number of interesting consequences for the algebraic counterpart of IpC, the theory of Heyting algebras. In particular we show that a model of IpC<sup>2</sup> can be constructed whose algebra of truth-values is equal to any given Heyting algebra.

**§1. Introduction.** This paper establishes a new and rather surprising property of Heyting's intuitionistic propositional calculus IpC. We show that quantification over propositional variables can be modelled in IpC, and hence that there is an interpretation of the second order propositional calculus IpC<sup>2</sup> in IpC which restricts to the identity on first order propositions. In order to state this result more precisely, we briefly recall the syntax and proof theory of first and second order intuitionistic propositional logic.

We will take the *first order propositions*,  $\phi$ , to be given by the following grammar:

$$\phi ::= p \mid \perp \mid \phi \wedge \phi' \mid \phi \vee \phi' \mid \phi \rightarrow \phi',$$

where  $p$  ranges over a set of *propositional variables*. Negation, truth and bi-implication can be defined in the usual way:

$$\neg \phi \stackrel{\text{def}}{=} \phi \rightarrow \perp, \quad \top \stackrel{\text{def}}{=} \neg \perp, \quad \phi \leftrightarrow \phi' \stackrel{\text{def}}{=} (\phi \rightarrow \phi') \wedge (\phi' \rightarrow \phi).$$

Table 1 gives a collection of natural deduction style rules for IpC. The premisses and conclusion of each rule are *sequents*,  $\Gamma \Rightarrow \phi$ , which we take to be specified by a

---

Received March 1, 1991.

This research was supported by the ESPRIT Basic Research Action No. 3003, "CLICS".

© 1992, Association for Symbolic Logic  
 0022-4812/92/5701-0004/\$03.00

finite *multiset* (unordered list)  $\Gamma$  of first order propositions and a single first order proposition  $\phi$ . The use of multisets rather than sets in sequents will be important when we consider the “size” of sequents in §2. For the moment, we note that since the order of formulas in  $\Gamma$  is immaterial, an explicit structural rule of exchange is not needed. As usual, we identify formulas with one-element multisets, and write  $\Gamma\Delta$  for the union of two multisets  $\Gamma$  and  $\Delta$ . We will write

$$\text{IpC} \vdash \Gamma \Rightarrow \phi$$

to indicate that the sequent is provable using the rules in Table 1. The new property of IpC which we establish here is:

**THEOREM 1.** *Given a propositional variable  $p$ , for each first order proposition  $\phi$  there is a first order proposition  $A_p\phi$ , containing only variables not equal to  $p$  which occur in  $\phi$ , and satisfying:*

- (i) *If  $\text{IpC} \vdash \Gamma \Rightarrow \phi$ , then  $\text{IpC} \vdash \Gamma \Rightarrow A_p\phi$ , provided  $p$  does not occur in  $\Gamma$ .*
- (ii) *If  $\text{IpC} \vdash \Gamma \Rightarrow A_p\phi$ , then for all  $\psi$ ,  $\text{IpC} \vdash \Gamma \Rightarrow \phi[\psi/p]$  (where  $\phi[\psi/p]$  denotes the result of substituting  $\psi$  for  $p$  throughout  $\phi$ ).*

This theorem will be proved in §2 using proof-theoretic methods. The key tool is the use of a Gentzen-style sequent calculus for IpC for which there is a well-founded relation on sequents making the hypotheses of each rule of the calculus less than its conclusion. (The first order proposition  $A_p\phi$  will be defined by recursion over this well-founded relation.) The particular sequent calculus we use is that given (independently) by Hudelmaier [6] and Dyckhoff [4]; its implicational part (the most important part) also occurs in the work of Lincoln, Scedrov and Shankar [8]. In fact, essentially similar refinements of the sequent calculus for IpC were developed by the Soviet school of proof theory some time ago—see Vorob’ev [13].

TABLE 1. Natural deduction rules for IpC.

---

$\frac{\Gamma \Rightarrow \psi}{\Gamma\phi \Rightarrow \psi}$ (Weaken)	$\frac{\Gamma\phi\phi \Rightarrow \psi}{\Gamma\phi \Rightarrow \psi}$ (Contract)
$\frac{}{\phi \Rightarrow \phi}$ (Id)	
$\frac{\Gamma \Rightarrow \perp}{\Gamma \Rightarrow \phi}$ ( $\perp$ Elim)	
$\frac{\Gamma \Rightarrow \phi \quad \Delta \Rightarrow \psi}{\Gamma\Delta \Rightarrow \phi \wedge \psi}$ ( $\wedge$ In <sub>tr</sub> )	$\frac{\Gamma \Rightarrow \phi \wedge \psi}{\Gamma \Rightarrow \phi}$ ( $\wedge$ Elim <sub>1</sub> )
	$\frac{\Gamma \Rightarrow \phi \wedge \psi}{\Gamma \Rightarrow \psi}$ ( $\wedge$ Elim <sub>2</sub> )
$\frac{\Gamma \Rightarrow \phi}{\Gamma \Rightarrow \phi \vee \psi}$ ( $\vee$ In <sub>tr</sub> <sub>1</sub> )	$\frac{\Gamma \Rightarrow \psi}{\Gamma \Rightarrow \phi \vee \psi}$ ( $\vee$ In <sub>tr</sub> <sub>2</sub> )
$\frac{\Gamma\phi \Rightarrow \theta \quad \Delta\psi \Rightarrow \theta \quad \Theta \Rightarrow \phi \vee \psi}{\Gamma\Delta\Theta \Rightarrow \theta}$ ( $\vee$ Elim)	
$\frac{\Gamma\phi \Rightarrow \psi}{\Gamma \Rightarrow \phi \rightarrow \psi}$ ( $\rightarrow$ In <sub>tr</sub> )	$\frac{\Gamma \Rightarrow \phi \rightarrow \psi \quad \Delta \Rightarrow \phi}{\Gamma\Delta \Rightarrow \psi}$ ( $\rightarrow$ Elim)

---

REMARK 2. It is perhaps worth pointing out that the analogue of Theorem 1 for classical logic is rather trivially true, since there we may take  $A_p\phi$  to be  $\phi[\top/p] \wedge \phi[\perp/p]$ .

The existence of first order propositions  $A_p\phi$  satisfying Theorem 1 enables one to interpret in IpC the *second order intuitionistic propositional logic* IpC<sup>2</sup>, a logic which extends IpC with quantification over propositional variables. As is well known, in this logic implication and universal quantification suffice to define the other connectives and existential quantification—see [11]. However, we will take the *second order propositions* to be given by the grammar

$$\phi ::= p \mid \perp \mid \phi \wedge \phi' \mid \phi \vee \phi' \mid \phi \rightarrow \phi' \mid \forall p\phi$$

in order that they be a superset of the first order propositions. The natural deduction rules for  $\forall p\phi$  are given in Table 2. Occurrences of  $p$  in  $\phi$  become bound in  $\forall p\phi$ ; all other types of occurrence of variables are free. We will write

$$\text{IpC}^2 \vdash \Gamma \Rightarrow \phi$$

to indicate that a sequent of second order propositions can be proved using the rules in Tables 1 and 2.

TABLE 2. Natural deduction rules for  $\forall$   
(\*provided  $p$  is not free in  $\Gamma$ ).

$\frac{\Gamma \Rightarrow \phi}{\Gamma \Rightarrow \forall p\phi} (\forall\text{Intr})^*$	$\frac{\Gamma \Rightarrow \forall p\phi}{\Gamma \Rightarrow \phi[\psi/p]} (\forall\text{Elim})$
---	---

In §3 we use Theorem 1 to define a translation of second order propositions,  $\phi$ , into first order ones,  $\phi^*$ . The translation has the following properties:

- (i) If  $\text{IpC}^2 \vdash \Gamma \Rightarrow \phi$ , then  $\text{IpC} \vdash \Gamma^* \Rightarrow \phi^*$  (where  $\Gamma^*$  indicates the translation applied elementwise to the multiset  $\Gamma$ ).
- (ii)  $p^* = p$ ,  $\perp^* = \perp$ ,  $(\phi \# \psi)^* = \phi^* \# \psi^*$  (for  $\# = \wedge, \vee, \rightarrow$ ), and hence in particular  $\phi^* = \phi$  for all first order propositions  $\phi$ .

See Proposition 9 below for more details. An immediate corollary of the existence of such an interpretation is a strengthening of the usual interpolation theorem for IpC. Recall that the latter says that, given first order propositions  $\phi$  and  $\psi$  for which  $\phi \Rightarrow \psi$  is provable in IpC, there is some first order proposition  $\theta$  containing only variables common to both  $\phi$  and  $\psi$ , and for which both  $\phi \Rightarrow \theta$  and  $\theta \Rightarrow \psi$  are provable. Here we establish (Proposition 11) that *the collection of such interpolant propositions is not merely nonempty but in fact contains least and greatest elements* (with respect to the provability ordering for IpC).<sup>1</sup>

Theorem 1 also has interesting consequences for the theory of Heyting algebras, a theory which bears the same relationship to IpC as does the theory of Boolean algebras to classical propositional logic. In §4 we establish a form of “second order completeness” for Heyting algebras: letting  $H[X]$  denote the Heyting algebra obtained by adjoining an indeterminate  $X$  to a given Heyting algebra  $H$ , we prove (Theorem 13) that *the inclusion  $i_H: H \hookrightarrow H[X]$  possesses both left and right adjoints,*

<sup>1</sup>The author is grateful to G. R. Renardel de Lavalette for pointing out that this solves his open problem 6.5 in [14].

i.e. there are monotone functions  $a_H, e_H: H[X] \rightarrow H$  satisfying  $(e_H \circ i_H) = \text{id}_H = (a_H \circ i_H)$  and  $(i_H \circ a_H) \leq \text{id}_{H[X]} \leq (i_H \circ e_H)$ . Moreover, these adjoints are natural in  $H$  in the category-theoretic sense. It follows from this that, given any Heyting algebra  $H$ , the generic model of the algebraic theory of “Heyting algebras equipped with a morphism from  $H$ ” is actually a model of  $\text{IpC}^2$ . Since the algebra of truth-values of this model is just  $H$ , we conclude (Proposition 18) that *any Heyting algebra can appear as the algebra of truth-values of a model of  $\text{IpC}^2$* . Equivalently, *any Heyting algebra can appear as the Lindenbaum algebra of a second order intuitionistic theory*.

The results presented in this paper have had a rather long gestation period. Some ten or so years ago I tried to prove the negation of Theorem 1 in connection with the higher order analogue of Proposition 18—the question of whether any Heyting algebra can appear as the algebra of truth-values of an elementary topos. I established that the free Heyting algebra on a countable infinity of generators does not so appear provided the property of  $\text{IpC}$  given in Theorem 1 does *not* hold. It seemed likely to me (and to others to whom I posed the question) that a first order proposition  $\phi$  could be found for which  $A_p\phi$  does not exist (although I could not find one!), thus settling the original question about toposes and Heyting algebras in the negative. That Theorem 1 is true is quite a surprise to me. Unfortunately, it appears that not all the results for second order logic reported here generalize to the setting of higher order logic. Whilst it is the case that Theorem 1 remains true if  $\text{IpC}$  is replaced by a quantifier-free fragment of intuitionistic higher order logic, the substitution property of Lemma 8 fails (so that one does not get an interpretation of full higher order logic in its quantifier-free fragment). It remains an open question whether every Heyting algebra can be the Lindenbaum algebra of a theory in intuitionistic higher order logic.

**Acknowledgement.** I would like to thank R. Dyckhoff for bringing to my attention the particular sequent calculus for  $\text{IpC}$  used in this paper. I would also like to thank him, A. Scedrov and G. Mints for elucidating its history.

**§2. Proof of Theorem 1.** Our proof will use the methods of proof theory. To be more precise, we will employ a certain refinement of the cut-free Gentzen sequent calculus for  $\text{IpC}$ , and we begin by explaining that.

Table 3 contains a fairly standard cut-free sequent calculus for  $\text{IpC}$ . This formulation of the sequent calculus has essential uses of the structural rules (Weaken) and (Contract) of Table 1 built in implicitly: weakening is built in via the (Atom) axiom, and an essential use of contraction is built in to the rule  $(\rightarrow \Rightarrow)$  by repeating the active proposition  $\phi \rightarrow \psi$  in the first premiss of the rule. We refer the reader to [3, Part 1, §3] for a proof of the fact that (Weaken), (Contract), (Id) and the cut rule

$$\frac{\Gamma \Rightarrow \phi \quad \phi \Delta \Rightarrow \psi}{\Gamma \Delta \Rightarrow \psi} \text{ (Cut)}$$

are all derivable from the rules in Table 3, and hence that these rules determine the same provable sequents as do those in Table 1.

TABLE 3. Sequent calculus for IpC  
 (\*where  $p$  is any propositional variable).

---


$$\frac{}{\Gamma p \Rightarrow p} \text{ (Atom)*}$$

$$\frac{}{\Gamma \perp \Rightarrow \phi} (\perp \Rightarrow)$$

$$\frac{\Gamma \Rightarrow \phi \quad \Gamma \Rightarrow \psi}{\Gamma \Rightarrow \phi \wedge \psi} (\Rightarrow \wedge) \quad \frac{\Gamma \phi \psi \Rightarrow \theta}{\Gamma(\phi \wedge \psi) \Rightarrow \theta} (\wedge \Rightarrow)$$

$$\frac{\Gamma \Rightarrow \phi}{\Gamma \Rightarrow \phi \vee \psi} (\Rightarrow \vee_1) \quad \frac{\Gamma \Rightarrow \psi}{\Gamma \Rightarrow \phi \vee \psi} (\Rightarrow \vee_2) \quad \frac{\Gamma \phi \Rightarrow \theta \quad \Gamma \psi \Rightarrow \theta}{\Gamma(\phi \vee \psi) \Rightarrow \theta} (\vee \Rightarrow)$$

$$\frac{\Gamma \phi \Rightarrow \psi}{\Gamma \Rightarrow \phi \rightarrow \psi} (\Rightarrow \rightarrow) \quad \frac{\Gamma(\phi \rightarrow \psi) \Rightarrow \phi \quad \Gamma \psi \Rightarrow \theta}{\Gamma(\phi \rightarrow \psi) \Rightarrow \theta} (\rightarrow \Rightarrow)$$


---

TABLE 4. Replacements for  $(\rightarrow \Rightarrow)$   
 (\*where  $p$  is any propositional variable).

---


$$\frac{\Gamma p \phi \Rightarrow \psi}{\Gamma p(p \rightarrow \phi) \Rightarrow \psi} (\rightarrow \Rightarrow_1)*$$

$$\frac{\Gamma(\phi_1 \rightarrow (\phi_2 \rightarrow \phi_3)) \Rightarrow \psi}{\Gamma((\phi_1 \wedge \phi_2) \rightarrow \phi_3) \Rightarrow \psi} (\rightarrow \Rightarrow_2)$$

$$\frac{\Gamma(\phi_1 \rightarrow \phi_3)(\phi_2 \rightarrow \phi_3) \Rightarrow \psi}{\Gamma((\phi_1 \vee \phi_2) \rightarrow \phi_3) \Rightarrow \psi} (\rightarrow \Rightarrow_3)$$

$$\frac{\Gamma(\phi_2 \rightarrow \phi_3) \Rightarrow \phi_1 \rightarrow \phi_2 \quad \Gamma \phi_3 \Rightarrow \psi}{\Gamma((\phi_1 \rightarrow \phi_2) \rightarrow \phi_3) \Rightarrow \psi} (\rightarrow \Rightarrow_4)$$


---

Note that  $(\rightarrow \Rightarrow)$  is the only rule in Table 3 which fails to have the property that its premisses are structurally simpler than its conclusion. Following Dyckhoff [4] and Hudelmaier [6], we can overcome this defect by replacing  $(\rightarrow \Rightarrow)$  by the four rules shown in Table 4. The new rules correspond to the possible forms of the antecedent of the introduced implication—except that a separate rule is not needed for the case when the antecedent is  $\perp$  (since the appropriate rule is an instance of weakening).

We will refer to the system of rules obtained from Table 3 by replacing  $(\rightarrow \Rightarrow)$  by the rules in Table 4 as LJ\*. Clearly any sequent provable in LJ\* is intuitionistically valid, since the rules in Table 4 are all derivable in IpC; the converse is also true, so that one has:

**THEOREM 3.**  $\text{IpC} \vdash \Gamma \Rightarrow \phi$  if and only if the sequent is provable in LJ\*.

We refer the reader to [4, Theorem 1] for a proof of this result.

**DEFINITION 4.** The *weight*,  $\text{wt}(\phi)$ , of a first order proposition  $\phi$  is a positive integer defined by induction on the structure of  $\phi$  as follows:

$$\begin{aligned}\text{wt}(p) &= \text{wt}(\perp) = 1, \\ \text{wt}(\phi \vee \psi) &= \text{wt}(\phi \rightarrow \psi) = \text{wt}(\phi) + \text{wt}(\psi) + 1, \\ \text{wt}(\phi \wedge \psi) &= \text{wt}(\phi) + \text{wt}(\psi) + 2.\end{aligned}$$

This weight function defines a well-ordering  $<$  on first order propositions via the definition

$$\phi < \psi \quad \text{if and only if} \quad \text{wt}(\phi) < \text{wt}(\psi).$$

Now extend  $<$  to a relation between finite multisets of propositions via the *multiset ordering* construction of Dershowitz and Manna. Thus  $\Gamma < \Delta$  holds if and only if  $\Delta = \Delta_1 \Delta_2$  and  $\Gamma = \Delta_1 \Gamma_2$ , for some  $\Delta_1, \Delta_2$  and  $\Gamma_2$  with  $\Delta_2$  nonempty and such that for all  $\gamma \in \Gamma_2$  there exists  $\delta \in \Delta_2$  with  $\gamma < \delta$ . As shown in [2], this relation between multisets is well-founded because the original relation on propositions is.

Finally, define a well-founded relation on sequents by declaring

$$(\Gamma \Rightarrow \phi) < (\Delta \Rightarrow \psi)$$

to hold just in case  $\Gamma\phi < \Delta\psi$ . Note that each rule in LJ\* has the property that a (premiss, conclusion)-pair lies in this relation between sequents.

We turn now to the proof of Theorem 1. Recall that in second order intuitionistic propositional logic, existentially quantified propositions  $\exists p\phi$  are definable in terms of  $\forall$  and  $\rightarrow$ :

$$\exists p\phi \stackrel{\text{def}}{=} \forall q(\forall p(\phi \rightarrow q) \rightarrow q)$$

(where  $q$  is not free in  $\phi$ ). It follows that the existence of the first order propositions  $A_p\phi$  for all  $\phi$ , with properties as in Theorem 1, entails the existence of first order propositions  $E_p\phi$  which model the existentially quantified proposition  $\exists p\phi$ . In fact to prove the theorem, we will need to define  $A_p\phi$  and  $E_p\phi$  simultaneously via mutual recursion. Moreover, we will need to give the definitions for multisets of formulas rather than for single formulas  $\phi$ , in order to utilize LJ\* to prove the required properties of the construction.

**PROPOSITION 5.** *Let  $\Delta$  be a finite multiset of first order propositions and  $\phi$  a single first order proposition. For each variable  $p$  there are first order propositions  $E_p(\Delta)$  and  $A_p(\Delta; \phi)$  satisfying the following conditions:*

- (i) (a)  $\text{Var}(E_p(\Delta)) \subseteq \text{Var}(\Delta) \setminus \{p\}$ , and
- (b)  $\text{Var}(A_p(\Delta; \phi)) \subseteq \text{Var}(\Delta\phi) \setminus \{p\}$ ,

where  $\text{Var}(\Delta)$  denotes the finite set of propositional variables in  $\Delta$ .

- (ii) (a)  $\text{IpC} \vdash \Delta \Rightarrow E_p(\Delta)$ , and
- (b)  $\text{IpC} \vdash \Delta A_p(\Delta; \phi) \Rightarrow \phi$ .

Moreover, for all finite multisets  $\Gamma$  of first order propositions not containing  $p$ , if  $\text{IpC} \vdash \Gamma\Delta \Rightarrow \phi$ , then:

- (iii) (a)  $\text{IpC} \vdash \Gamma E_p(\Delta) \Rightarrow \phi$ , provided  $p \notin \text{Var}(\phi)$ , and
- (b)  $\text{IpC} \vdash \Gamma E_p(\Delta) \Rightarrow A_p(\Delta; \phi)$ .

Theorem 1 follows immediately from Proposition 5 if we make the definition

$$(1) \quad A_p\phi \stackrel{\text{def}}{=} A_p(\emptyset; \phi).$$

For if  $\text{IpC} \vdash \Gamma \Rightarrow \phi$  with  $p \notin \text{Var}(\Gamma)$ , then by 5(iii)(b)

$$\text{IpC} \vdash \Gamma E_p(\emptyset) \Rightarrow A_p(\emptyset; \phi)$$

and by 5(ii)(a)

$$\text{IpC} \vdash \emptyset \Rightarrow E_p(\emptyset),$$

which together give  $\text{IpC} \vdash \Gamma \Rightarrow A_p\phi$ , as required for part (i) of the theorem. For part (ii) of the theorem, note that 5(ii)(b) gives  $\text{IpC} \vdash A_p\phi \Rightarrow \phi$ . Hence the sequent obtained from this one by substituting a proposition  $\psi$  for  $p$  throughout is also provable; but by 5(i)(b)  $p \notin \text{Var}(A_p\phi)$ , so the result of such a substitution is  $\text{IpC} \vdash A_p\phi \Rightarrow \phi[\psi/p]$ . So  $\text{IpC} \vdash \Gamma \Rightarrow \phi[\psi/p]$  holds whenever  $\text{IpC} \vdash \Gamma \Rightarrow A_p\phi$  does, as required.

REMARK 6. Properties (i)(a), (ii)(a) and (iii)(a) together imply that  $E_p(\Delta)$  acts like the existentially quantified formula  $\exists p(\bigwedge \Delta)$  (where  $\bigwedge \Delta$  denotes the conjunction of the formulas in  $\Delta$ ). However, properties (i)(b), (ii)(a), (ii)(b) and (iii)(b) imply that it is  $E_p(\Delta) \rightarrow A_p(\Delta; \phi)$ , rather than just  $A_p(\Delta; \phi)$ , which acts like the universally quantified formula  $\forall p(\bigwedge \Delta \rightarrow \phi)$ . This is because of the appearance of  $E_p(\Delta)$  in (iii)(b)—a complication which is needed to carry through the proof of (iii). This proof proceeds by induction on the structure of the proof of  $\Gamma \Delta \Rightarrow \phi$  in LJ\*, and it is the case where the proof ends with the active formula an implication contained in  $\Gamma$  which requires us to prove (iii)(b) rather than  $\text{IpC} \vdash \Gamma \Rightarrow A_p(\Delta; \phi)$ .

The rest of this section will be devoted to the proof of Proposition 5.

The formulas  $E_p(\Delta)$  and  $A_p(\Delta; \phi)$  are defined simultaneously by  $\prec$ -induction on  $\Delta\phi$  (where  $\prec$  is the well-founded relation of Definition 4). At each stage, we define  $E_p(\Delta)$  as the conjunction of a finite set of formulas  $\mathcal{E}_p(\Delta)$  and  $A_p(\Delta; \phi)$  as the disjunction of a finite set of formulas  $\mathcal{A}_p(\Delta; \phi)$ :

$$E_p(\Delta) \stackrel{\text{def}}{=} \bigwedge \mathcal{E}_p(\Delta), \quad A_p(\Delta; \phi) \stackrel{\text{def}}{=} \bigvee \mathcal{A}_p(\Delta; \phi).$$

The elements of the finite sets  $\mathcal{E}_p(\Delta)$  and  $\mathcal{A}_p(\Delta; \phi)$  are given by Table 5, with one element for each match of  $\Delta$  and  $\Delta; \phi$  to the patterns listed in the left-hand column of the table. It is quite possible that in a particular case there are no matches, so that  $\mathcal{E}_p(\Delta)$  or  $\mathcal{A}_p(\Delta; \phi)$  is empty—in which case  $E_p(\Delta) = \bigwedge \emptyset \stackrel{\text{def}}{=} \top$  and  $A_p(\Delta; \phi) = \bigvee \emptyset \stackrel{\text{def}}{=} \perp$ .

It follows easily by  $\prec$ -induction on  $\Delta\phi$  that  $E_p(\Delta)$  and  $A_p(\Delta; \phi)$  are built up from subformulas of  $\Delta\phi$  and that they do not contain  $p$ . So 5(i) holds.

The validity of the sequents in 5(ii) is proved simultaneously by  $\prec$ -induction on  $\Delta\phi$ . At each stage we have to show that  $\text{IpC} \vdash \Delta \Rightarrow \varepsilon$  and  $\text{IpC} \vdash \Delta\alpha \Rightarrow \phi$  hold for each  $\varepsilon \in \mathcal{E}_p(\Delta)$  and each  $\alpha \in \mathcal{A}_p(\Delta; \phi)$ . For each of the cases (E0)–(E8) and (A1)–(A13) of Table 5, this follows from the induction hypothesis by straightforward proofs in IpC.

We turn now to 5(iii); this is proved by induction on the structure of a proof of  $\Gamma \Delta \Rightarrow \phi$  in LJ\*, with one case for each proof rule:



TABLE 5. Definition of  $E_p(\Delta)$  and  $A_p(\Delta; \phi)$   
( $q$  denotes any propositional variable not equal to  $p$ ).

	$\Delta$ matches	$\mathcal{E}_p(\Delta)$ contains
(E0)	$\Delta' \perp$	$\perp$
(E1)	$\Delta' q$	$E_p(\Delta') \wedge q$
(E2)	$\Delta'(\delta_1 \wedge \delta_2)$	$E_p(\Delta' \delta_1 \delta_2)$
(E3)	$\Delta'(\delta_1 \vee \delta_2)$	$E_p(\Delta' \delta_1) \vee E_p(\Delta' \delta_2)$
(E4)	$\Delta'(q \rightarrow \delta)$	$q \rightarrow E_p(\Delta' \delta)$
(E5)	$\Delta' p(p \rightarrow \delta)$	$E_p(\Delta' p \delta)$
(E6)	$\Delta'((\delta_1 \wedge \delta_2) \rightarrow \delta_3)$	$E_p(\Delta'(\delta_1 \rightarrow (\delta_2 \rightarrow \delta_3)))$
(E7)	$\Delta'((\delta_1 \vee \delta_2) \rightarrow \delta_3)$	$E_p(\Delta'(\delta_1 \rightarrow \delta_3)(\delta_2 \rightarrow \delta_3))$
(E8)	$\Delta'((\delta_1 \rightarrow \delta_2) \rightarrow \delta_3)$	$[E_p(\Delta'(\delta_2 \rightarrow \delta_3)) \rightarrow A_p(\Delta'(\delta_2 \rightarrow \delta_3); \delta_1 \rightarrow \delta_2)] \rightarrow E_p(\Delta' \delta_3)$
	$\Delta; \phi$ matches	$\mathcal{A}_p(\Delta; \phi)$ contains
(A1)	$\Delta' q; \phi$	$A_p(\Delta'; \phi)$
(A2)	$\Delta'(\delta_1 \wedge \delta_2); \phi$	$A_p(\Delta' \delta_1 \delta_2; \phi)$
(A3)	$\Delta'(\delta_1 \vee \delta_2); \phi$	$[E_p(\Delta' \delta_1) \rightarrow A_p(\Delta' \delta_1; \phi)] \wedge [E_p(\Delta' \delta_2) \rightarrow A_p(\Delta' \delta_2; \phi)]$
(A4)	$\Delta'(q \rightarrow \delta); \phi$	$q \wedge A_p(\Delta' \delta; \phi)$
(A5)	$\Delta' p(p \rightarrow \delta); \phi$	$A_p(\Delta' p \delta; \phi)$
(A6)	$\Delta'((\delta_1 \wedge \delta_2) \rightarrow \delta_3); \phi$	$A_p(\Delta'(\delta_1 \rightarrow (\delta_2 \rightarrow \delta_3)); \phi)$
(A7)	$\Delta'((\delta_1 \vee \delta_2) \rightarrow \delta_3); \phi$	$A_p(\Delta'(\delta_1 \rightarrow \delta_3)(\delta_2 \rightarrow \delta_3); \phi)$
(A8)	$\Delta'((\delta_1 \rightarrow \delta_2) \rightarrow \delta_3); \phi$	$[E_p(\Delta'(\delta_2 \rightarrow \delta_3)) \rightarrow A_p(\Delta'(\delta_2 \rightarrow \delta_3); \delta_1 \rightarrow \delta_2)] \wedge A_p(\Delta' \delta_3; \phi)$
(A9)	$\Delta; q$	$q$
(A10)	$\Delta' p; p$	$\top$
(A11)	$\Delta; \phi_1 \wedge \phi_2$	$A_p(\Delta; \phi_1) \wedge A_p(\Delta; \phi_2)$
(A12)	$\Delta; \phi_1 \vee \phi_2$	$A_p(\Delta; \phi_1) \vee A_p(\Delta; \phi_2)$
(A13)	$\Delta; \phi_1 \rightarrow \phi_2$	$E_p(\Delta \phi_1) \rightarrow A_p(\Delta \phi_1; \phi_2)$

*Case (Atom).* So  $\phi$  is a propositional variable and is an element of  $\Gamma \Delta$ , i.e.  $\phi \in \Gamma$  or  $\phi \in \Delta$ . We will split the argument into two subcases according to whether  $\phi$  is the variable  $p$  or not.

*Subcase  $\phi = p$ .* In this case we just have to check that (b) holds for  $\Gamma \Delta \Rightarrow \phi$ . Since  $p \notin \Gamma$ , we must have  $\Delta = \Delta' p$ . Then case (A10) of Table 5 gives  $\text{IpC} \vdash \top \Rightarrow A_p(\Delta' p; p)$ , from which (b) follows.

*Subcase  $\phi \neq p$ .* We know that either  $\phi \in \Gamma$  or  $\phi \in \Delta$ . In the first case (a) holds for  $\Gamma \Delta \Rightarrow \phi$  by (Atom), and (b) follows from (a) because, by case (A9) of Table 5,  $\text{IpC} \vdash \phi \Rightarrow A_p(\Delta; \phi)$ . On the other hand, if  $\phi \in \Delta$ , say  $\Delta = \Delta' \phi$ , case (E1) of Table 5 gives  $\text{IpC} \vdash E_p(\Delta' \phi) \Rightarrow E_p(\Delta') \wedge \phi$ , from which (a) for  $\Gamma \Delta \Rightarrow \phi$  follows; and as before (b) follows from (a) by case (A9).

*Case ( $\perp \Rightarrow$ ).* So  $\perp \in \Gamma \Delta$ . If  $\perp \in \Gamma$  then (a) and (b) hold by ( $\perp \Rightarrow$ ). If  $\perp \in \Delta$ , then, by case (E0) of Table 5,  $\vdash E_p(\Delta) \Rightarrow \perp$ , from which (a) and (b) follow.

*Case* ( $\Rightarrow \wedge$ ). So  $\phi = \phi_1 \wedge \phi_2$ ,  $\vdash \Gamma \Delta \Rightarrow \phi_i$  ( $i = 1, 2$ ), and (a) and (b) hold for these sequents by the induction hypothesis.

(a) If  $p \notin \text{Var}(\phi)$  then  $p \notin \text{Var}(\phi_i)$ , so by (a) for  $\Gamma \Delta \Rightarrow \phi_i$  we have

$$\text{IpC} \vdash \Gamma E_p(\Delta) \Rightarrow \phi_i$$

for  $i = 1, 2$ , from which (a) for  $\Gamma \Delta \Rightarrow \phi$  follows.

(b) By (b) for  $\Gamma \Delta \Rightarrow \phi_i$  we have  $\vdash \Gamma E_p(\Delta) \Rightarrow A_p(\Delta; \phi_i)$  for  $i = 1, 2$ , and hence

$$\text{IpC} \vdash \Gamma E_p(\Delta) \Rightarrow A_p(\Delta; \phi_1) \wedge A_p(\Delta; \phi_2).$$

But by case (A11) of Table 5

$$\text{IpC} \vdash A_p(\Delta; \phi_1) \wedge A_p(\Delta; \phi_2) \Rightarrow A_p(\Delta; \phi).$$

So (b) for  $\Gamma \Delta \Rightarrow \phi$  follows.

*Case* ( $\wedge \Rightarrow$ ). *Subcase 1.*  $\Gamma = \Gamma'(\gamma_1 \wedge \gamma_2)$ ,  $\text{IpC} \vdash \Gamma' \gamma_1 \gamma_2 \Delta \Rightarrow \phi$ , and (a) and (b) hold for this sequent by the induction hypothesis. Hence (a) and (b) hold for  $\Gamma \Delta \Rightarrow \phi$  by an application of ( $\wedge \Rightarrow$ ).

*Subcase 2.*  $\Delta = \Delta'(\delta_1 \wedge \delta_2)$ ,  $\text{IpC} \vdash \Gamma \Delta' \delta_1 \delta_2 \Rightarrow \phi$ , and (a) and (b) hold for this sequent by the induction hypothesis. From this it follows that (a) and (b) hold for  $\Gamma \Delta \Rightarrow \phi$ , since by cases (E2) and (A2) of Table 5 we have

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow E_p(\Delta' \delta_1 \delta_2), \quad \text{IpC} \vdash A_p(\Delta' \delta_1 \delta_2; \phi) \Rightarrow A_p(\Delta; \phi).$$

*Case* ( $\Rightarrow \vee_i$ ). This case is analogous to that for ( $\Rightarrow \wedge$ ), using case (A12) of Table 5.

*Case* ( $\vee \Rightarrow$ ). *Subcase 1.*  $\Gamma = \Gamma'(\gamma_1 \vee \gamma_2)$ ,  $\text{IpC} \vdash \Gamma' \gamma_i \Delta \Rightarrow \phi$  ( $i = 1, 2$ ), and (a) and (b) hold for these sequents by the induction hypothesis. Hence (a) and (b) hold for  $\Gamma \Delta \Rightarrow \phi$  by an application of ( $\vee \Rightarrow$ ).

*Subcase 2.*  $\Delta = \Delta'(\delta_1 \vee \delta_2)$ ,  $\text{IpC} \vdash \Gamma \Delta' \delta_i \Rightarrow \phi$  ( $i = 1, 2$ ), and (a) and (b) hold for these sequents by the induction hypothesis.

(a) If  $p \notin \text{Var}(\phi)$ , then by (a) for  $\Gamma \Delta' \delta_i \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma E_p(\Delta' \delta_i) \Rightarrow \phi$$

for  $i = 1, 2$ , and thus

$$\text{IpC} \vdash \Gamma (E_p(\Delta' \delta_1) \vee E_p(\Delta' \delta_2)) \Rightarrow \phi.$$

But by case (E3) of Table 5

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow E_p(\Delta' \delta_1) \vee E_p(\Delta' \delta_2),$$

and hence (a) holds for  $\Gamma \Delta \Rightarrow \phi$ .

(b) By (b) for  $\Gamma \Delta' \delta_i \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma E_p(\Delta' \delta_i) \Rightarrow A_p(\Delta' \delta_i; \phi)$$

for  $i = 1, 2$ , and thus

$$\text{IpC} \vdash \Gamma \Rightarrow (E_p(\Delta' \delta_1) \rightarrow A_p(\Delta' \delta_1; \phi)) \wedge (E_p(\Delta' \delta_2) \rightarrow A_p(\Delta' \delta_2; \phi)).$$

But by case (A3) of Table 5

$$\text{IpC} \vdash (E_p(\Delta' \delta_1) \rightarrow A_p(\Delta' \delta_1; \phi)) \wedge (E_p(\Delta' \delta_2) \rightarrow A_p(\Delta' \delta_2; \phi)) \Rightarrow A_p(\Delta; \phi),$$

so that  $\text{IpC} \vdash \Gamma \Rightarrow A_p(\Delta; \phi)$ , and hence in particular (b) holds for  $\Gamma \Delta \Rightarrow \phi$ .

*Case* ( $\Rightarrow \rightarrow$ ). So  $\phi = \phi_1 \rightarrow \phi_2$ ,  $\text{IpC} \vdash \Gamma \Delta \phi_1 \Rightarrow \phi_2$ , and (a) and (b) hold for this sequent by the induction hypothesis.

(a) If  $p \notin \text{Var}(\phi)$ , then  $p \notin \text{Var}(\phi_i)$  for  $i = 1, 2$ . So by (a) for  $\Gamma \Delta \phi_1 \Rightarrow \phi_2$  we have

$$\text{IpC} \vdash \Gamma \phi_1 E_p(\Delta) \Rightarrow \phi_2,$$

and hence, by an application of ( $\Rightarrow \rightarrow$ ), (a) holds for  $\Gamma \Delta \Rightarrow \phi$ .

(b) By (b) for  $\Gamma \Delta \phi_1 \Rightarrow \phi_2$  we have

$$\text{IpC} \vdash \Gamma E_p(\Delta \phi_1) \Rightarrow A_p(\Delta \phi_1; \phi_2).$$

But since

$$\text{IpC} \vdash (E_p(\Delta \phi_1) \rightarrow A_p(\Delta \phi_1; \phi_2)) \Rightarrow A_p(\Delta; \phi)$$

holds by case (A13) of Table 5, we get  $\text{IpC} \vdash \Gamma \Rightarrow A_p(\Delta; \phi)$ , and so in particular (b) holds for  $\Gamma \Delta \Rightarrow \phi$ .

*Case* ( $\rightarrow \Rightarrow_1$ ). For this rule there are four subcases to consider according to how the active formula and its atomic antecedent occur in  $\Gamma \Delta$ .

*Subcase 1.*  $\Gamma = \Gamma' q (q \rightarrow \gamma)$  with  $q \neq p$  (because  $p \notin \text{Var}(\Gamma)$ ),  $\text{IpC} \vdash \Gamma' q \gamma \Delta \Rightarrow \phi$ , and (a) and (b) hold for this sequent by the induction hypothesis. Consequently (a) and (b) hold for  $\Gamma \Delta \Rightarrow \phi$  as well, by an application of ( $\rightarrow \Rightarrow_1$ ).

*Subcase 2.*  $\Gamma = \Gamma' q$ ,  $\Delta = \Delta' (q \rightarrow \delta)$ ,  $\text{IpC} \vdash \Gamma' q \Delta' \delta \Rightarrow \phi$ , and (a) and (b) hold for this sequent by the induction hypothesis.

(a) If  $p \notin \text{Var}(\phi)$ , then by (a) for  $\Gamma' q \Delta' \delta \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma' q E_p(\Delta' \delta) \Rightarrow \phi,$$

and hence

$$\text{IpC} \vdash \Gamma (q \rightarrow E_p(\Delta' \delta)) \Rightarrow \phi$$

by an application of ( $\rightarrow \Rightarrow_1$ ). But by case (E4) of Table 5 we also have

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow q \rightarrow E_p(\Delta' \delta),$$

and hence (a) holds for  $\Gamma \Delta \Rightarrow \phi$ .

(b) By (b) for  $\Gamma' q \Delta' \delta \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma' q E_p(\Delta' \delta) \Rightarrow A_p(\Delta' \delta; \phi),$$

and hence

$$\text{IpC} \vdash \Gamma (q \rightarrow E_p(\Delta' \delta)) \Rightarrow q \wedge A_p(\Delta' \delta; \phi).$$

But by cases (E4) and (A4) of Table 5, we also have

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow q \rightarrow E_p(\Delta' \delta),$$

$$\text{IpC} \vdash q \wedge A_p(\Delta' \delta; \phi) \Rightarrow A_p(\Delta; \phi),$$

and hence (b) holds for  $\Gamma \Delta \Rightarrow \phi$ .

*Subcase 3.*  $\Gamma = \Gamma' (q \rightarrow \gamma)$ ,  $\Delta = \Delta' q$ ,  $\text{IpC} \vdash \Gamma' q \gamma \Delta' \Rightarrow \phi$ , and (a) and (b) hold for this sequent by the induction hypothesis.

(a) If  $p \notin \text{Var}(\phi)$  then by (a) for  $\Gamma' q \gamma \Delta' \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma' q \gamma E_p(\Delta') \Rightarrow \phi,$$

and hence

$$\text{IpC} \vdash \Gamma(q \wedge E_p(\Delta')) \Rightarrow \phi.$$

But by case (E1) of Table 5 we have

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow q \wedge E_p(\Delta'),$$

and hence (a) holds for  $\Gamma\Delta \Rightarrow \phi$ .

(b) By (b) for  $\Gamma'q\gamma\Delta' \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma'q\gamma E_p(\Delta') \Rightarrow A_p(\Delta'; \phi),$$

and hence

$$\vdash \Gamma(q \wedge E_p(\Delta')) \Rightarrow A_p(\Delta'; \phi).$$

But by cases (E1) and (A1) of Table 5, we have

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow q \wedge E_p(\Delta'),$$

$$\text{IpC} \vdash A_p(\Delta'; \phi) \Rightarrow A_p(\Delta; \phi),$$

and hence (b) holds for  $\Gamma\Delta \Rightarrow \phi$ .

*Subcase 4.*  $\Delta = \Delta'd(d \rightarrow \delta)$  with  $d$  a propositional variable,  $\text{IpC} \vdash \Gamma\Delta'd\delta \Rightarrow \phi$ , and (a) and (b) hold for this sequent by the induction hypothesis.

*Subsubcase*  $d = p$ . (a) If  $p \notin \text{Var}(\phi)$ , then by (a) for  $\Gamma\Delta'p\delta \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma E_p(\Delta'p\delta) \Rightarrow \phi.$$

But by case (E5) of Table 5 we also have

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow E_p(\Delta'p\delta),$$

and hence (a) holds for  $\Gamma\Delta \Rightarrow \phi$ .

(b) By (b) for  $\Gamma\Delta'p\delta \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma E_p(\Delta'p\delta) \Rightarrow A_p(\Delta'p\delta; \phi).$$

Hence, using cases (E5) and (A5) of Table 5, we get (a) for  $\Gamma\Delta \Rightarrow \phi$ .

*Subsubcase*  $d \neq p$ . (a) If  $p \notin \text{Var}(\phi)$ , then by (a) for  $\Gamma\Delta'd\delta \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma d E_p(\Delta'\delta) \Rightarrow \phi.$$

But by cases (E1) and (E4) of Table 5 we have

$$\text{IpC} \vdash E_p(\Delta) \Rightarrow E_p(\Delta'(d \rightarrow \delta)) \wedge d \Rightarrow (d \rightarrow E_p(\Delta'\delta)) \wedge d \Rightarrow E_p(\Delta'\delta) \wedge d,$$

and hence (a) holds for  $\Gamma\Delta \Rightarrow \phi$ .

(b) By (b) for  $\Gamma\Delta'd\delta \Rightarrow \phi$  we have

$$\text{IpC} \vdash \Gamma d E_p(\Delta'\delta) \Rightarrow A_p(\Delta'\delta; \phi),$$

and hence

$$\text{IpC} \vdash \Gamma E_p(\Delta'\delta) \wedge d \Rightarrow d \wedge A_p(\Delta'\delta; \phi).$$

But as above, by cases (E1) and (E4) of Table 5 we have  $\text{IpC} \vdash E_p(\Delta) \Rightarrow E_p(\Delta'\delta) \wedge d$ ;

and similarly, by cases (A4) and (A1) we have

$$\text{IpC} \vdash d \wedge A_p(\Delta' \delta; \phi) \Rightarrow A_p(\Delta'(d \rightarrow \delta); \phi) \Rightarrow A_p(\Delta' d(d \rightarrow \delta); \phi).$$

Hence (b) holds for  $\Gamma \Delta \Rightarrow \phi$ .

*Case* ( $\rightarrow \Rightarrow_2$ ). This case is analogous to that for ( $\wedge \Rightarrow$ ), using cases (E6) and (A6) of Table 5.

*Case* ( $\rightarrow \Rightarrow_3$ ). This case is analogous to that for ( $\wedge \Rightarrow$ ), using cases (E7) and (A7) of Table 5.

*Case* ( $\rightarrow \Rightarrow_4$ ). *Subcase 1.*  $\Gamma = \Gamma'((\gamma_1 \rightarrow \gamma_2) \rightarrow \gamma_3)$ ,  $\text{IpC} \vdash \Gamma'(\gamma_2 \rightarrow \gamma_3) \Delta \Rightarrow \gamma_1 \rightarrow \gamma_2$ ,  $\text{IpC} \vdash \Gamma' \gamma_3 \Delta \Rightarrow \phi$ , and (a) and (b) hold for these sequents by the induction hypothesis, i.e.

$$\begin{aligned} \text{IpC} \vdash \Gamma'(\gamma_2 \rightarrow \gamma_3) E_p(\Delta) &\Rightarrow \gamma_1 \rightarrow \gamma_2, \\ \text{IpC} \vdash \Gamma' \gamma_3 E_p(\Delta) &\Rightarrow A_p(\Delta; \phi), \end{aligned}$$

and

$$\text{IpC} \vdash \Gamma' \gamma_3 E_p(\Delta) \Rightarrow \phi,$$

when  $p \notin \text{Var}(\phi)$ . Then (a) and (b) for  $\Gamma \Delta \Rightarrow \phi$  follow from these by an application of ( $\rightarrow \Rightarrow_4$ ).

*Subcase 2.*  $\Delta = \Delta'((\delta_1 \rightarrow \delta_2) \rightarrow \delta_3)$ ,  $\text{IpC} \vdash \Gamma \Delta'(\delta_2 \rightarrow \delta_3) \Rightarrow \delta_1 \rightarrow \delta_2$ ,  $\text{IpC} \vdash \Gamma \Delta' \delta_3 \Rightarrow \phi$ , and (a) and (b) hold for these sequents by the induction hypothesis, i.e.

$$(2) \quad \text{IpC} \vdash \Gamma E_p(\Delta'(\delta_2 \rightarrow \delta_3)) \Rightarrow A_p(\Delta'(\delta_2 \rightarrow \delta_3); \delta_1 \rightarrow \delta_2),$$

$$(3) \quad \text{IpC} \vdash \Gamma E_p(\Delta' \delta_3) \Rightarrow A_p(\Delta' \delta_3; \phi),$$

and

$$(4) \quad \text{IpC} \vdash \Gamma E_p(\Delta' \delta_3) \Rightarrow \phi$$

when  $p \notin \text{Var}(\phi)$ —in which case, combining (2), (4) and case (E8) of Table 5, we have that (a) holds for  $\Gamma \Delta \Rightarrow \phi$ . For (b), from (2) we get

$$(5) \quad \text{IpC} \vdash \Gamma \Rightarrow E_p(\Delta'(\delta_2 \rightarrow \delta_3)) \rightarrow A_p(\Delta'(\delta_2 \rightarrow \delta_3); \delta_1 \rightarrow \delta_2),$$

and this together with (3) and case (E8) of Table 5 yields

$$(6) \quad \text{IpC} \vdash \Gamma E_p(\Delta) \Rightarrow A_p(\Delta' \delta_3; \phi).$$

Then (5), (6) and case (A8) of Table 5 together give that (b) holds for  $\Gamma \Delta \Rightarrow \phi$ .

This completes the proof of Proposition 5.  $\square$

**§3. Interpreting  $\text{IpC}^2$  in  $\text{IpC}$ .** Using the propositions  $A_p \phi$  defined in the previous section, we can translate second order propositions into first order ones.

**DEFINITION 7.** For each second order proposition  $\phi$ , define a first order proposition  $\phi^*$  by induction on the structure of  $\phi$  as follows:

$$\begin{aligned} p^* &\stackrel{\text{def}}{=} p, \\ \perp^* &\stackrel{\text{def}}{=} \perp, \\ (\phi \# \psi)^* &\stackrel{\text{def}}{=} \phi^* \# \psi^* \quad (\# = \wedge, \vee, \rightarrow), \\ (\forall p \phi)^* &\stackrel{\text{def}}{=} A_p \phi^*. \end{aligned}$$

In order to see that this translation sends  $\text{IpC}^2$ -provable sequents to  $\text{IpC}$ -provable ones, we need to establish a crucial property of the mapping  $\phi \mapsto A_p\phi$ , namely that it commutes with substitution. It is a peculiarity of second order logic (compared with third or higher order logic) that this follows automatically from the properties of  $A_p\phi$  established in Theorem 1.

LEMMA 8. *Given distinct propositional variables  $p$  and  $q$ , and first order propositions  $\phi$  and  $\psi$  with  $p, q \notin \text{Var}(\psi)$ , one has*

$$\text{IpC} \vdash \emptyset \Rightarrow A_p(\phi[\psi/q]) \leftrightarrow (A_p\phi)[\psi/q].$$

PROOF. By part (ii) of Theorem 1 we have  $\text{IpC} \vdash A_p\phi \Rightarrow \phi$ , and hence  $\text{IpC} \vdash (A_p\phi)[\psi/q] \Rightarrow \phi[\psi/q]$ ; but since  $p$  does not occur in  $(A_p\phi)[\psi/q]$ , we can apply part (i) of the theorem to conclude that  $\text{IpC} \vdash (A_p\phi)[\psi/q] \Rightarrow A_p(\phi[\psi/q])$ .

To prove the converse, we use the following congruence property of  $\leftrightarrow$  in  $\text{IpC}$ :

$$(7) \quad \text{IpC} \vdash (\psi \leftrightarrow \psi') \Rightarrow \theta[\psi/q] \leftrightarrow \theta[\psi'/q].$$

From part (ii) of Theorem 1 we have

$$\text{IpC} \vdash A_p(\phi[\psi/q]) \Rightarrow \phi[\psi/q],$$

and so, using (7), we also have

$$\text{IpC} \vdash (\psi \leftrightarrow q)A_p(\phi[\psi/q]) \Rightarrow \phi.$$

Since  $p$  does not occur in the left-hand side of this sequent, part (i) of the theorem implies that we also have

$$\text{IpC} \vdash (\psi \leftrightarrow q)A_p(\phi[\psi/q]) \Rightarrow A_p\phi.$$

On substituting  $\psi$  for  $q$  throughout this sequent, we obtain

$$\text{IpC} \vdash A_p(\phi[\psi/q]) \Rightarrow (A_p\phi)[\psi/q],$$

as required. □

PROPOSITION 9. *The translation  $\phi \mapsto \phi^*$  has the following properties:*

(i) *For all sequents of second order propositions, if  $\text{IpC}^2 \vdash \Gamma \Rightarrow \phi$ , then  $\text{IpC} \vdash \Gamma^* \Rightarrow \phi^*$  (where  $\Gamma^*$  indicates the translation applied elementwise to the multiset  $\Gamma$ ).*

(ii) *If  $\phi$  is a first order proposition, then  $\phi^* = \phi$ .*

*Thus  $(-)^*$  gives an interpretation of  $\text{IpC}^2$  into  $\text{IpC}$  which restricts to the identity on first order propositions.*

PROOF. Part (i) is proved by induction on the structure of the proof of  $\Gamma \Rightarrow \phi$  from the rules in Tables 1 and 2. The induction step for rule  $(\forall\text{Intr})$  uses part (i) of Theorem 1. The induction step for rule  $(\forall\text{Elim})$  uses part (ii) of the theorem in conjunction with Lemma 8.

Part (ii) is immediate from the definition of  $(-)^*$ . □

Since  $\text{IpC}$ -provability is decidable whereas  $\text{IpC}^2$ -provability is not (see [5], for example), the interpretation  $(-)^*$  cannot be conservative. Here is an interesting example of an unprovable second order proposition whose interpretation is provable.

EXAMPLE 10. For all second order propositions  $\phi$  and  $\psi$ ,

$$\text{IpC} \vdash \emptyset \Rightarrow [\forall p(\phi \vee \psi) \leftrightarrow (\forall p\phi \vee \forall p\psi)]^*.$$

For combining Definition 7 with the definition of  $A_p$  in (1), we have

$$(\forall p(\phi \vee \psi))^* = A_p(\emptyset; \phi^* \vee \psi^*).$$

The construction of  $A_p$  according to Table 5 then gives

$$A_p(\emptyset; \phi^* \vee \psi^*) = A_p(\emptyset; \phi^*) \vee A_p(\emptyset; \psi^*) = (\forall p\phi)^* \vee (\forall p\psi)^*.$$

Second order intuitionistic propositional logic has an interpolation property which is a trivial consequence of quantification over propositions. Using the interpretation  $(-)^*$ , we can transfer this to a nontrivial interpolation property for IpC which strengthens the usual interpolation theorem for this logic.

**PROPOSITION 11.** *Let  $\phi$  and  $\psi$  be first order propositions for which  $\text{IpC} \vdash \phi \Rightarrow \psi$  holds. Call a first order proposition  $\theta$  an **interpolant** for  $(\phi, \psi)$  if*

- $\text{Var}(\theta) \subseteq \text{Var}(\phi) \cap \text{Var}(\psi)$ , and
- $\text{IpC} \vdash \phi \Rightarrow \theta$  and  $\text{IpC} \vdash \theta \Rightarrow \psi$ .

*There exist first order propositions  $\varepsilon$  and  $\alpha$  which are respectively least and greatest interpolants for  $(\phi, \psi)$ , in the sense that any interpolant  $\theta$  satisfies  $\text{IpC} \vdash \varepsilon \Rightarrow \theta$  and  $\text{IpC} \vdash \theta \Rightarrow \alpha$ .*

**PROOF.** Suppose that

$$\text{Var}(\phi) = \{p_1, \dots, p_u, q_1, \dots, q_v\}, \quad \text{Var}(\psi) = \{q_1, \dots, q_v, r_1, \dots, r_w\},$$

with the variables  $p_i, q_j, r_k$  pairwise distinct. Since  $\text{IpC} \vdash \phi \Rightarrow \psi$ , it is also the case that  $\text{IpC}^2 \vdash \phi \Rightarrow \psi$ , and hence

$$\text{IpC}^2 \vdash \exists p_1 \dots \exists p_u \phi \Rightarrow \psi, \quad \text{IpC}^2 \vdash \phi \Rightarrow \forall r_1 \dots \forall r_w \psi,$$

since  $p_1, \dots, p_u \notin \text{Var}(\psi)$  and  $r_1, \dots, r_w \notin \text{Var}(\phi)$ . Furthermore, one always has

$$\text{IpC}^2 \vdash \phi \Rightarrow \exists p_1 \dots \exists p_u \phi, \quad \text{IpC}^2 \vdash \forall r_1 \dots \forall r_w \psi \Rightarrow \psi.$$

So, if we set  $\varepsilon \stackrel{\text{def}}{=} (\exists p_1 \dots \exists p_u \phi)^*$  and  $\alpha \stackrel{\text{def}}{=} (\forall r_1 \dots \forall r_w \psi)^*$ , properties (i) and (ii) in Proposition 9 applied to the above sequents imply that  $\varepsilon$  and  $\alpha$  are indeed interpolants for  $(\phi, \psi)$  in IpC. Moreover, if  $\theta$  is any other first order interpolant, from  $\text{IpC} \vdash \theta \Rightarrow \psi$  we get  $\text{IpC}^2 \vdash \theta \Rightarrow \psi$  and hence  $\text{IpC}^2 \vdash \theta \Rightarrow \forall r_1 \dots \forall r_w \psi$  (since  $r_1, \dots, r_w \notin \text{Var}(\theta)$ ); and then on applying  $(-)^*$  we get  $\text{IpC} \vdash \theta \Rightarrow \alpha$ . Similarly,  $\text{IpC} \vdash \phi \Rightarrow \theta$  implies  $\text{IpC} \vdash \varepsilon \Rightarrow \theta$ .  $\square$

**§4. Heyting algebra applications.** Recall that a *Heyting algebra*  $H$  is a (distributive) lattice in which every pair of elements  $h, h' \in H$  possesses a relative pseudocomplement,  $h \rightarrow h'$  (the greatest element whose meet with  $h$  lies underneath  $h'$  in the ordering on  $H$ ). A *morphism of Heyting algebras* is a function preserving all finite meets, finite joins and relative pseudocomplements. We will denote by **Heyt** the category of Heyting algebras and morphisms. As their name suggests, Heyting algebras are the models of an algebraic theory: see for example Balbes and Dwinger [1, Chapter IX] for an equational presentation and further information on the theory of Heyting algebras.

The relationship between Heyting algebras and intuitionistic logic is exactly analogous to that between Boolean algebras and classical logic. In particular, there is a correspondence between Heyting algebras and first order intuitionistic propositional theories, induced by the process of forming the Lindenbaum algebra of a theory. Thus for each set  $G$ , let  $F\langle G \rangle$  denote the set of first order propositions built up from the elements of  $G$  regarded as propositional variables. Given a subset  $R \subseteq F\langle G \rangle$ , let

$$\begin{aligned} F\langle G \rangle &\rightarrow F\langle G; R \rangle, \\ \phi &\mapsto [\phi], \end{aligned}$$

denote the quotient of  $F\langle G \rangle$  by the equivalence relation identifying two first order propositions  $\phi$  and  $\psi$  if and only if  $\text{IpC} \vdash \Gamma \Rightarrow \phi \leftrightarrow \psi$  holds for some finite  $\Gamma \subseteq R$ . Endowing  $F\langle G; R \rangle$  with the partial order

$$[\phi] \leq [\psi] \text{ if and only if for some finite } \Gamma \subseteq R, \text{IpC} \vdash \Gamma \Rightarrow \phi \rightarrow \psi,$$

one obtains a Heyting algebra—the Lindenbaum algebra of the IpC-theory (determined by)  $R$  over the language  $G$ . Moreover, every Heyting algebra  $H$  can be presented in this way: for example, take  $G$  to be the set  $H$  itself and  $R$  to be those propositions mapped to  $\top$  by the obvious evaluation function  $F\langle G \rangle \rightarrow H$  induced by the identity function  $G \rightarrow H$ .

Theorem 1 can be rephrased as a statement about the relationship between a Heyting algebra  $H$  and the algebra obtained from it by freely adjoining an indeterminate  $X$ .

**DEFINITION 12.** Given a Heyting algebra  $H$ , a *Heyting polynomial algebra over  $H$*  is a Heyting algebra  $H[X]$  equipped with a distinguished element  $X$  and a morphism  $i_H: H \rightarrow H[X]$  with the following universal property:

For each morphism  $f: H \rightarrow K$  in **Heyt** and each element  $k \in K$ , there is a unique morphism  $g: H[X] \rightarrow K$  such that  $g \circ i_H = f$  and  $g(X) = k$ .

As usual, the universal property in the definition determines  $H[X]$  uniquely up to isomorphism (over  $H$ ). Given a presentation of  $H$  as  $F\langle G; R \rangle$ , one can present  $H[X]$  as  $F\langle G \cup \{p\}; R \rangle$ , where  $p$  is any element not contained in  $G$ ; the morphism  $i_H$  is induced by the inclusion  $F\langle G \rangle \subseteq F\langle G \cup \{p\} \rangle$ , and the distinguished element  $X$  is the equivalence class  $[p]$ .

When  $R = \emptyset$ ,  $H = F\langle G; R \rangle$  is the *free* Heyting algebra on the set of generators  $G$  and  $H[X]$  is free on  $G \cup \{p\}$ . Except for the case of a single generator, the structure of free Heyting algebras is not well understood. The following theorem sheds some new light on this structure.

**THEOREM 13.** For any Heyting algebra  $H$ , the morphism  $i_H: H \rightarrow H[X]$  possesses both left and right adjoints. In other words, there exist functions  $e_H: H[X] \rightarrow H$  and  $a_H: H[X] \rightarrow H$  satisfying


$$(8) \quad e_H(P) \leq h \quad \text{if and only if} \quad P \leq i_H(h),$$

$$(9) \quad h \leq a_H(P) \quad \text{if and only if} \quad i_H(h) \leq P,$$

for all  $P \in H[X]$  and  $h \in H$ .



These adjoints are natural in  $H$ . In other words, for each morphism  $f: H \rightarrow H'$  in **Heyt** the following squares commute



(where  $f[X]$  is the unique morphism  $g$  satisfying  $(g \circ i_H) = (i_{H'} \circ f)$  and  $g(X) = X$  whose existence is guaranteed by the universal property of  $H[X]$  in Definition 12).

PROOF. As remarked above, we can assume  $H = F\langle G; R \rangle$  for some  $G$  and  $R$ , and then take  $H[X] = F\langle G \cup \{p\}; R \rangle$ . Thus each element  $h \in H$  is of the form  $h = [\psi]$ , where  $\psi$  is some first order proposition with variables in  $G$ . Similarly, each  $P \in H[X]$  is of the form  $P = [\phi]$ , where  $\phi$  may involve  $p$  as well as variables in  $G$ . Moreover,  $i_H(h) \leq P$  in  $H[X]$  if and only if  $\text{IpC} \vdash \Gamma \Rightarrow \psi \rightarrow \phi$  holds for some finite  $\Gamma \subseteq R$ . But by Theorem 1 we have

$$\begin{aligned}
 \text{IpC} \vdash \Gamma \Rightarrow \psi \rightarrow \phi & \text{ iff } \text{IpC} \vdash \Gamma \psi \Rightarrow \phi \\
 & \text{ iff } \text{IpC} \vdash \Gamma \psi \Rightarrow A_p \phi \\
 & \text{ iff } \text{IpC} \vdash \Gamma \Rightarrow \psi \rightarrow A_p \phi.
 \end{aligned}$$

So on defining  $a_H(P) = [A_p \phi]$  we get (9). (Clearly the definition of  $a_H(P)$  is independent of the choice of representative for  $P$ .) The naturality of  $a_H$  follows from Lemma 8, because any morphism  $F\langle G; R \rangle \rightarrow F\langle G'; R' \rangle$  is induced by a function  $F\langle G \rangle \rightarrow F\langle G' \rangle$  of the form

$$\psi \mapsto \psi[\gamma'_g/g \mid g \in G]$$

for some  $G$ -indexed family  $(\gamma'_g \in F\langle G' \rangle \mid g \in G)$ . Alternatively, the naturality of  $a_H$  can be deduced from an interpolation property of pushout squares in **Heyt**—see [9, Theorem B].

The existence of the left adjoints  $e_H$  (and their naturality) follows from the existence of the natural right adjoints by an algebraic version of the proof that  $\exists$  is definable from  $\forall$  and  $\rightarrow$  in  $\text{IpC}^2$ . Alternatively, we can use Proposition 5 to give a direct definition:

$$e_H([\phi]) = [E_p(\phi)]. \qquad \square$$

REMARKS 14. (i) The morphism  $i_H: H \rightarrow H[X]$  is always a monomorphism (since the universal property of  $H[X]$  implies that  $i_H$  has a left inverse sending  $X$  to, for example, the top element of  $H$ ). Consequently the adjoints to  $i_H$  satisfy

$$e_H \circ i_H = \text{id}_H = a_H \circ i_H.$$

(Of course they also satisfy  $i_H \circ a_H \leq \text{id}_{H[X]} \leq i_H \circ e_H$ .)

(ii) (cf. Remark 2). The analogue of Theorem 13 for Boolean algebras is rather trivial. This is because a Boolean polynomial in a single indeterminate  $X$  with coefficients in a given Boolean algebra  $B$  can always be put in the normal form

$$(b \wedge X) \vee (b' \wedge \neg X) \quad (b, b' \in B).$$

Then the inclusion of  $B$  into its algebra of Boolean polynomials is given by

$$b \mapsto (b \wedge X) \vee (b \wedge \neg X),$$

and it is easy to see that this has left and right adjoints, given respectively by  $(b \wedge X) \vee (b' \wedge \neg X) \mapsto (b \vee b')$  and  $(b \wedge X) \vee (b' \wedge \neg X) \mapsto (b \wedge b')$ .

Consider the morphism

$$k: F\langle G; R \rangle \rightarrow F\langle G \cup \{p\}; R \cup \{\rho\} \rangle$$

induced by the inclusion  $F\langle G \rangle \subseteq F\langle G \cup \{p\} \rangle$ , where as before  $p$  is some new element not in  $G$  and now we have also extended  $R$  by adding in some  $\rho \in F\langle G \cup \{p\} \rangle$ . The proof of Theorem 13 extends to show that this morphism  $k$  also has left and right adjoints,  $e_k$  and  $a_k$ , given by:

$$e_k([\phi]) = [E_p(\rho \wedge \phi)], \quad a_k([\phi]) = [A_p(\rho \rightarrow \phi)].$$

Iterating, we get adjoints for the morphism induced by extending  $G$  and  $R$  by finitely many elements. Such morphisms are precisely the finitely presented objects in the locally finitely presentable category  $H/\mathbf{Heyt}$  of Heyting algebras equipped with a morphism from  $H (= F\langle G; R \rangle)$  (and whose morphisms are commutative triangles). So we get the following corollary (the second paragraph of which follows from the first and the interpolation property [9, Theorem B] of pushout squares in  $\mathbf{Heyt}$ ):

**COROLLARY 15.** *Any morphism  $k: H \rightarrow K$  in  $\mathbf{Heyt}$  which makes  $K$  finitely presented over  $H$  possesses both left and right adjoints  $e_k, a_k: K \rightarrow H$ .*

*Furthermore, given any morphism  $f: H \rightarrow H'$ , forming the pushout square*

$$\begin{array}{ccc}
 K & \xrightarrow{f'} & K' \\
 \uparrow k & & \uparrow k' \\
 H & \xrightarrow{f} & H'
 \end{array}$$

*in  $\mathbf{Heyt}$ , ( $k'$  necessarily makes  $K'$  finitely presented over  $H'$  and) the adjoints satisfy  $f \circ e_k = e_{k'} \circ f'$  and  $f \circ a_k = a_{k'} \circ f'$ .  $\square$*

These results enable us to construct, for each Heyting algebra  $H$ , a model of the second order calculus  $\text{IpC}^2$ . To explain further, we must describe what is needed to specify such a model. The particular notion of model we will use is the specialization from categories to partial orders of the notion of model of the second order lambda calculus described in [10] and [12].

First note that since the notion of Heyting algebra is algebraic, it makes sense to speak of a *Heyting algebra object*,  $U$ , in any category  $\mathcal{C}$  with finite products: such an object comes equipped with morphisms

$$\top, \perp : 1 \rightarrow U, \quad \wedge, \vee, \rightarrow : U \times U \rightarrow U,$$

making various diagrams (derived from the defining equations of the theory of Heyting algebras) commute in  $\mathcal{C}$ . As usual, this structure induces an ordinary Heyting algebra structure on the hom-sets  $\mathcal{C}(I, U)$ ; and precomposition with  $f: I \rightarrow I'$  gives a morphism of Heyting algebras,  $f^*: \mathcal{C}(I', U) \rightarrow \mathcal{C}(I, U)$ .

DEFINITION 16. Say that a Heyting algebra object  $U$  in  $\mathcal{C}$  possesses *internal  $U$ -indexed meets* if for each object  $I \in \mathcal{C}$  there is a right adjoint

$$\bigwedge_I : \mathcal{C}(I \times U, U) \rightarrow \mathcal{C}(I, U)$$

to the morphism  $\pi_1^*$  induced by composition with the first projection morphism  $\pi_1: I \times U \rightarrow I$ , and moreover these right adjoints are natural in  $I$ .

REMARK 17. The import of this condition on  $U$  becomes more apparent from the point of view of the internal higher order logic of the topos  $[\mathcal{C}^{\text{op}}, \mathcal{S}et]$  of presheaves on  $\mathcal{C}$ . Identifying the objects  $I$  of  $\mathcal{C}$  with their corresponding presheaves  $\mathcal{C}(-, I)$ , we see that the functions  $\bigwedge_I$  constitute a morphism  $U^U \rightarrow U$  giving the meet of an internal  $U$ -indexed family of elements of  $U$ .

A Heyting algebra object  $U$  possessing internal  $U$ -indexed meets in a category  $\mathcal{C}$  with finite products determines a model of  $\text{IpC}^2$ . For each second order proposition  $\phi$  with free variables in the list  $\vec{p} = p_1, \dots, p_n$  of distinct variables, we get a morphism

$$\llbracket \phi(\vec{p}) \rrbracket : U^n \rightarrow U$$

defined by induction on the structure of  $\phi$  as follows:

$$\begin{aligned} \llbracket p_i(\vec{p}) \rrbracket &\stackrel{\text{def}}{=} (U^n \xrightarrow{\pi_i} U), \\ \llbracket \perp(\vec{p}) \rrbracket &\stackrel{\text{def}}{=} (U^n \rightarrow 1 \xrightarrow{\perp} U), \\ \llbracket (\phi \# \psi)(\vec{p}) \rrbracket &\stackrel{\text{def}}{=} (U^n \xrightarrow{\langle \llbracket \phi(\vec{p}) \rrbracket, \llbracket \psi(\vec{p}) \rrbracket \rangle} U \times U \xrightarrow{\#} U), \\ \llbracket (\forall p\phi)(\vec{p}) \rrbracket &\stackrel{\text{def}}{=} \bigwedge_{U^n} (U^n \times U \xrightarrow{\llbracket \phi(\vec{p}p) \rrbracket} U) \end{aligned}$$

(where  $\# = \wedge, \vee, \text{ or } \rightarrow$ ). In particular, when  $\phi$  contains no free variables we can take  $\vec{p}$  to be empty and obtain a global element  $\llbracket \phi \rrbracket \in \mathcal{C}(1, U)$  of  $U$  for each closed second order proposition  $\phi$ . We will call the Heyting algebra  $\mathcal{C}(1, U)$  the *algebra of truth-values* of the model  $(\mathcal{C}, U)$ .

We say that  $(\mathcal{C}, U)$  *satisfies* such a closed proposition  $\phi$ , and write  $(\mathcal{C}, U) \models \phi$ , if  $\llbracket \phi \rrbracket$  is the top element of  $\mathcal{C}(1, U)$ . This notion of satisfaction is sound for provability in  $\text{IpC}^2$ : if  $\text{IpC}^2 \vdash \emptyset \Rightarrow \phi$  then  $(\mathcal{C}, U) \models \phi$ . Conversely, it is not hard to prove (by a term model construction) that it is also complete:  $\text{IpC}^2 \vdash \emptyset \Rightarrow \phi$  holds if  $\phi$  is satisfied by all  $(\mathcal{C}, U)$ .

Returning now to Theorem 13, given  $H \in \mathbf{Heyt}$ , let  $\mathcal{C}$  be the opposite of the full subcategory of  $H/\mathbf{Heyt}$  consisting of the finitely generated free objects. More concretely, we can take the objects of  $\mathcal{C}$  to be finite ordinals,  $[n]$ , and the morphisms  $[n] \rightarrow [m]$  to be  $m$ -tuples of elements of the polynomial Heyting algebra in  $n$  indeterminates,  $H[X_1, \dots, X_n]$ . Composition is given by substitution, and the

identity on  $[n]$  is  $(X_1, \dots, X_n)$ . Lawvere’s categorical treatment of algebraic theories (see [7], for example) tells us that  $\mathcal{C}$  (has finite products and) contains the generic model of the algebraic theory of “Heyting algebras equipped with a morphism from  $H$ ”. In particular,  $\mathcal{C}$  does contain a Heyting algebra object, namely  $U = [1]$ : its top and bottom elements are  $(\top), (\perp): [0] \rightarrow [1]$  and its meet, join and pseudocomplementation operations are  $(X_1 \# X_2): [1] \times [1] = [2] \rightarrow [1]$  (for  $\# = \wedge, \vee, \rightarrow$ ).

For this Heyting algebra object we have for each object  $[n] \in \mathcal{C}$  that

$$\begin{aligned} \mathcal{C}([n], U) &= \mathcal{C}([n], [1]) = H[X_1, \dots, X_n], \\ \mathcal{C}([n] \times U, U) &= \mathcal{C}([n + 1], [1]) = H[X_1, \dots, X_n, X_{n+1}] \\ &\cong H[X_1, \dots, X_n][X], \end{aligned}$$

and  $\pi_1^*$  is  $i_{H[X_1, \dots, X_n]}$ . Consequently, Theorem 13 implies that  $U$  has internal  $U$ -indexed meets and hence determines a model of  $\text{IpC}^2$ . Note that the algebra of truth-values of this model is  $\mathcal{C}(1, U) = \mathcal{C}([0], [1]) = H$ , the given Heyting algebra. We have thus proved:

**PROPOSITION 18.** *Given a Heyting algebra  $H$ , the algebraic theory of “Heyting algebras equipped with a morphism from  $H$ ” has the property that its generic model  $U$  has internal  $U$ -indexed meets, and hence provides a model of second order intuitionistic propositional logic. Since the algebra of truth-values of this model is just  $H$ , we conclude that every Heyting algebra appears as the algebra of truth-values of some model of  $\text{IpC}^2$ . □*

There is a correspondence between instances of the notion of model of  $\text{IpC}^2$  as we have defined it, and  $\text{IpC}^2$ -theories. (By such a theory we mean a suitable language together with a collection of axioms—second order propositions over the language.) Under this correspondence, the algebra of truth-values of a model is identified with the Lindenbaum algebra of the theory (i.e. the collection of closed second order propositions over the given language, quotiented by provability in  $\text{IpC}^2$  augmented with the given axioms). Consequently, Proposition 18 implies:

*Every Heyting algebra is the Lindenbaum algebra of some  $\text{IpC}^2$ -theory.*

In fact one can see this without recourse to the correspondence between models and theories, using the interpretation of  $\text{IpC}^2$  into  $\text{IpC}$  developed in §3. For given a Heyting algebra  $H$ , choose a presentation for it as  $H = F\langle G; R \rangle$ . Then the set of second order propositions over  $G$

$$\{\phi \mid \text{IpC} \vdash \Gamma \Rightarrow \phi^*, \text{ for some finite } \Gamma \subseteq R\}$$

determines an  $\text{IpC}^2$ -theory over  $G$  whose Lindenbaum algebra is isomorphic to  $F\langle G; R \rangle$  (the isomorphism being induced by  $(-)^*$ ).

REFERENCES

[1] R. BALBES and P. DWINGER, *Distributive lattices*, University of Missouri Press, Columbia, Missouri, 1974.  
 [2] N. DERSHOWITZ and Z. MANNA, *Proving termination with multiset orderings*, *Communications of the Association for Computing Machinery*, vol. 22 (1979), pp. 465–476.  
 [3] A. G. DRAGALIN, *Mathematical intuitionism*, American Mathematical Society, Providence, Rhode Island, 1988.

- [4] R. DYCKHOFF, *Contraction-free sequent calculi for intuitionistic logic*, preprint, St. Andrews University, 1990.
- [5] D. M. GABBAY, *Semantical investigations in Heyting's intuitionistic logic*, Synthese Library, vol. 148, Reidel, Dordrecht, 1981.
- [6] J. HUDELMAIER, *Bounds for cut elimination in intuitionistic propositional logic*, Ph.D. Thesis, University of Tübingen, Tübingen, 1989.
- [7] A. KOCK and G. E. REYES, *Doctrines in categorical logic*, **Handbook of Mathematical Logic** (J. Barwise, editor), North-Holland, Amsterdam, 1977, Chapter A.8, pp. 283–313.
- [8] P. LINCOLN, A. SCEDROV and N. SHANKAR, *Linearizing intuitionistic implication*, **Proceedings of the sixth annual symposium on logic in computer science**, IEEE Computer Society Press, Washington, D.C., 1991 (to appear).
- [9] A. M. PITTS, *Amalgamation and interpolation in the category of Heyting algebras*, **Journal of Pure and Applied Algebra**, vol. 29 (1983), pp. 155–165.
- [10] ———, *Polymorphism is set theoretic, constructively*, **Category theory and computer science, proceedings, Edinburgh, 1987** (D. Pitt et al., editors), Lecture Notes in Computer Science, vol. 283, Springer-Verlag, Berlin, 1987, pp. 12–39.
- [11] D. PRAWITZ, *Natural deduction*, Almqvist & Wiksell, Stockholm, 1965.
- [12] R. A. G. SEELY, *Categorical semantics for higher order polymorphic lambda calculus*, this JOURNAL, vol. 52 (1987) pp. 969–989.
- [13] N. N. VOROB'EV, *A new algorithm for derivability in the constructive propositional calculus*, **American Mathematical Society Translations**, ser. 2, vol. 94 (1970), pp. 37–71.
- [14] G. R. RENARDEL DE LAVALETTE, *Interpolation in fragments of intuitionistic propositional logic*, this JOURNAL, vol. 54 (1989), pp. 1419–1430.

COMPUTER LABORATORY  
CAMBRIDGE UNIVERSITY  
CAMBRIDGE CB2 3QG, ENGLAND

*E-mail:* Andrew.Pitts@computer-lab.cam.ac.uk