HOL Light Very Quick Reference

compiled by John Harrison, mangled by Freek Wiedijk

Theorems (type `thm`)

```
ADD1                    |- SUC m = m + 1
ADD_AC                  |- m + n = n + m /\ (m + n) + p = m + n + p /\ m + n + p = n + m + p
ADD_ASSOC               |- m + n + p = (m + n) + p
ADD_CLAUSES             |- (!n. 0 + n = n) /\ (!m. m + 0 = m) /\ (!m n. SUC m + n = SUC (m + n)) /\ (!m n. m + SUC n = SUC (m + n))
ADD_SUB                 |- (m + n) - n = m
ADD_SYM                 |- m + n = n + m
ALL                     |- (ALL P [] <=> T) /\ (ALL P (CONS h t) <=> P h /\ ALL P t)
ALL2                    |- (ALL2 P [] [] <=> T) /\ ... /\ (ALL2 P (CONS h1 t1) (CONS h2 t2) <=> P h1 h2 /\ ALL2 P t1 t2)
APPEND                  |- (!l. APPEND [] l = l) /\ (!h t l. APPEND (CONS h t) l = CONS h (APPEND t l))
ARITH                   |- (NUMERAL 0 = 0 /\ BIT0 _0 = _0) /\ ((!n. SUC (NUMERAL n) = NUMERAL (SUC n)) /\ ...
ARITH_EQ                |- (!m n. NUMERAL m = NUMERAL n <=> m = n) /\ (_0 = _0 <=> T) /\ ...
CARD_CLAUSES            |- CARD {} = 0 /\ (!x s. FINITE s ==> CARD (x INSERT s) = (if x IN s then CARD s else SUC (CARD s)))
CART_EQ                 |- x = y <=> (!i. 1 <= i /\ i <= dimindex UNIV ==> x $ i = y $ i)
CONJ_ASSOC              |- t1 /\ t2 /\ t3 <=> (t1 /\ t2) /\ t3
DE_MORGAN_THM           |- (~(t1 /\ t2) <=> ~t1 \/ ~t2) /\ (~(t1 \/ t2) <=> ~t1 /\ ~t2)
DIVISION                |- ~(n = 0) ==> m = m DIV n * n + m MOD n /\ m MOD n < n
ETA_AX                  |- (\x. t x) = t
EVEN                    |- (EVEN 0 <=> T) /\ (!n. EVEN (SUC n) <=> ~EVEN n)
EXISTS_REFL             |- ?x. x = a
EXP                     |- (!m. m EXP 0 = 1) /\ (!m n. m EXP SUC n = m * m EXP n)
EXTENSION               |- s = t <=> (!x. x IN s <=> x IN t)
FACT                    |- FACT 0 = 1 /\ (!n. FACT (SUC n) = SUC n * FACT n)
FINITE_INDUCT_STRONG    |- P {} /\ (!x s. P s /\ ~(x IN s) /\ FINITE s ==> P (x INSERT s)) ==> (!s. FINITE s ==> P s)
FINITE_NUMSEG           |- FINITE (m .. n)
FINITE_RULES            |- FINITE {} /\ (!x s. FINITE s ==> FINITE (x INSERT s))
FINITE_SUBSET           |- FINITE t /\ s SUBSET t ==> FINITE s
FORALL_PAIR_THM         |- (!p. P p) <=> (!p1 p2. P (p1,p2))
FUN_EQ_THM              |- f = g <=> (!x. f x = g x)
GE                      |- m >= n <=> n <= m
HAS_SIZE                |- s HAS_SIZE n <=> FINITE s /\ CARD s = n
HD                      |- HD (CONS h t) = h
IMP_IMP                 |- p ==> q ==> r <=> p /\ q ==> r
IN                      |- x IN P <=> P x
IN_DELETE               |- x IN s DELETE y <=> x IN s /\ ~(x = y)
IN_ELIM_THM             |- (!P x. x IN GSPEC (\v. P (SETSPEC v)) <=> P (\p t. p /\ x = t)) /\ ...
IN_IMAGE                |- y IN IMAGE f s <=> (?x. y = f x /\ x IN s)
IN_INSERT               |- x IN y INSERT s <=> x = y \/ x IN s
IN_INTER                |- x IN s INTER t <=> x IN s /\ x IN t
IN_NUMSEG               |- p IN m .. n <=> m <= p /\ p <= n
IN_SING                 |- x IN {y} <=> x = y
IN_UNION                |- x IN s UNION t <=> x IN s \/ x IN t
IN_UNIV                 |- x IN UNIV
LAMBDA_BETA             |- 1 <= i /\ i <= dimindex UNIV ==> (lambda) g $ i = g i
LAST                    |- LAST (CONS h t) = (if t = [] then h else LAST t)
LE                      |- (!m. m <= 0 <=> m = 0) /\ (!m n. m <= SUC n <=> m = SUC n \/ m <= n)
LEFT_ADD_DISTRIB        |- m * (n + p) = m * n + m * p
LEFT_IMP_EXISTS_THM     |- (?x. P x) ==> Q <=> (!x. P x ==> Q)
LENGTH                  |- LENGTH [] = 0 /\ (!h t. LENGTH (CONS h t) = SUC (LENGTH t))
LENGTH_APPEND           |- LENGTH (APPEND l m) = LENGTH l + LENGTH m
LE_0                    |- 0 <= n
LE_ADD                  |- m <= m + n
LE_EXISTS               |- m <= n <=> (?d. n = m + d)
LE_MULT_LCANCEL         |- m * n <= m * p <=> m = 0 \/ n <= p
LE_REFL                 |- n <= n
LE_TRANS                |- m <= n /\ n <= p ==> m <= p
LT                      |- (!m. m < 0 <=> F) /\ (!m n. m < SUC n <=> m = n \/ m < n)
LT_0                    |- 0 < SUC n
LT_REFL                 |- ~(n < n)
MEM                     |- (MEM x [] <=> F) /\ (MEM x (CONS h t) <=> x = h \/ MEM x t)
MEMBER_NOT_EMPTY        |- (?x. x IN s) <=> ~(s = {})
MONO_EXISTS             |- (!x. P x ==> Q x) ==> (?x. P x) ==> (?x. Q x)
MONO_FORALL             |- (!x. P x ==> Q x) ==> (!x. P x) ==> (!x. Q x)
MULT_AC                 |- m * n = n * m /\ (m * n) * p = m * n * p /\ m * n * p = n * m * p
MULT_ASSOC              |- m * n * p = (m * n) * p
MULT_CLAUSES            |- (!n. 0 * n = 0) /\ ... /\ (!m n. m * SUC n = m + m * n)
MULT_SYM                |- m * n = n * m
NOT_CONS_NIL            |- ~(CONS h t = [])
NOT_EXISTS_THM          |- ~(?x. P x) <=> (!x. ~P x)
NOT_FORALL_THM          |- ~(!x. P x) <=> (?x. ~P x)
NOT_IMP                 |- ~(t1 ==> t2) <=> t1 /\ ~t2
NOT_IN_EMPTY            |- ~(x IN {})
NOT_LE                  |- ~(m <= n) <=> n < m
NOT_LT                  |- ~(m < n) <=> n <= m
NOT_SUC                 |- ~(SUC n = 0)
PAIR_EQ                 |- x,y = a,b <=> x = a /\ y = b
PRE                     |- PRE 0 = 0 /\ (!n. PRE (SUC n) = n)
```

```
REAL_ABS_MUL                    |- abs (x * y) = abs x * abs y
REAL_ABS_NEG                    |- abs (--x) = abs x
REAL_ABS_NUM                    |- abs (&n) = &n
REAL_ABS_POS                    |- &0 <= abs x
REAL_ABS_POW                    |- abs (x pow n) = abs x pow n
REAL_ADD_ASSOC                  |- x + y + z = (x + y) + z
REAL_ADD_LID                    |- &0 + x = x
REAL_ADD_LINV                   |- --x + x = &0
REAL_ADD_RID                    |- x + &0 = x
REAL_ADD_SYM                    |- x + y = y + x
REAL_ENTIRE                     |- x * y = &0 <=> x = &0 \/ y = &0
REAL_EQ_IMP_LE                  |- x = y ==> x <= y
REAL_INV_MUL                    |- inv (x * y) = inv x * inv y
REAL_LET_TRANS                  |- x <= y /\ y < z ==> x < z
REAL_LE_LMUL                    |- &0 <= x /\ y <= z ==> x * y <= x * z
REAL_LE_LT                      |- x <= y <=> x < y \/ x = y
REAL_LE_REFL                    |- x <= x
REAL_LE_SQUARE                  |- &0 <= x * x
REAL_LE_TOTAL                   |- x <= y \/ y <= x
REAL_LTE_TRANS                  |- x < y /\ y <= z ==> x < z
REAL_LT_01                      |- &0 < &1
REAL_LT_DIV                     |- &0 < x /\ &0 < y ==> &0 < x / y
REAL_LT_IMP_LE                  |- x < y ==> x <= y
REAL_LT_IMP_NZ                  |- &0 < x ==> ~(x = &0)
REAL_LT_LE                      |- x < y <=> x <= y /\ ~(x = y)
REAL_LT_MUL                     |- &0 < x /\ &0 < y ==> &0 < x * y
REAL_LT_REFL                    |- ~(x < x)
REAL_LT_TRANS                   |- x < y /\ y < z ==> x < z
REAL_MUL_AC                     |- m * n = n * m /\ (m * n) * p = m * n * p /\ m * n * p = n * m * p
REAL_MUL_ASSOC                  |- x * y * z = (x * y) * z
REAL_MUL_LID                    |- &1 * x = x
REAL_MUL_LINV                   |- ~(x = &0) ==> inv x * x = &1
REAL_MUL_LZERO                  |- &0 * x = &0
REAL_MUL_RID                    |- x * &1 = x
REAL_MUL_RINV                   |- ~(x = &0) ==> x * inv x = &1
REAL_MUL_RZERO                  |- x * &0 = &0
REAL_MUL_SYM                    |- x * y = y * x
REAL_NEGNEG                     |- -- --x = x
REAL_NEG_NEG                    |- -- --x = x
REAL_NOT_LE                     |- ~(x <= y) <=> y < x
REAL_NOT_LT                     |- ~(x < y) <=> y <= x
REAL_OF_NUM_ADD                 |- &m + &n = &(m + n)
REAL_OF_NUM_EQ                  |- &m = &n <=> m = n
REAL_OF_NUM_LE                  |- &m <= &n <=> m <= n
REAL_OF_NUM_LT                  |- &m < &n <=> m < n
REAL_OF_NUM_MUL                 |- &m * &n = &(m * n)
REAL_OF_NUM_POW                 |- &x pow n = &(x EXP n)
REAL_POS                        |- &0 <= &n
REAL_POW_2                      |- x pow 2 = x * x
REAL_POW_ADD                    |- x pow (m + n) = x pow m * x pow n
REAL_SUB_0                      |- x - y = &0 <=> x = y
REAL_SUB_LDISTRIB               |- x * (y - z) = x * y - x * z
REAL_SUB_LE                     |- &0 <= x - y <=> y <= x
REAL_SUB_LT                     |- &0 < x - y <=> y < x
REAL_SUB_REFL                   |- x - x = &0
REAL_SUB_RZERO                  |- x - &0 = x
RIGHT_ADD_DISTRIB               |- (m + n) * p = m * p + n * p
RIGHT_FORALL_IMP_THM            |- (!x. P ==> Q x) <=> P ==> (!x. Q x)
SKOLEM_THM                      |- (!x. ?y. P x y) <=> (?y. !x. P x (y x))
SUBSET                          |- s SUBSET t <=> (!x. x IN s ==> x IN t)
SUC_INJ                         |- SUC m = SUC n <=> m = n
TL                              |- TL (CONS h t) = t
TRUTH                           |- T
```

Inference rules (return type `thm`)

```
AC th tm                        Prove equivalence by associativity and commutativity
AP_TERM tm th                   From |- s = t to |- f s = f t
AP_THM th tm                    From |- f = g to |- f x = g x
ARITH_RULE tm                   Linear arithmetic prover over N
ASSUME tm                       Generate trivial theorem p |- p
BETA_RULE th                    Reduce all beta-redexes in theorem
CONJ th th                      From |- p and |- q to |- p /\ q
CONJUNCT1 th                    From |- p /\ q to |- p
CONJUNCT2 th                    From |- p /\ q to |- q
CONV_RULE conv th               Apply conversion to conclusion of theorem
DISCH tm th                     From p |- q to |- p ==> q
DISCH_ALL th                    From p1, ..., pn |- q to |- p1 ==> ... ==> pn ==> q
EQT_ELIM th                     From |- p <=> T to |- p
EQT_INTRO th                    From |- p to |- p <=> T
EQ_MP th th                     From |- p <=> q and |- p to |- q
GEN tm th                       From |- p[x] to |- !x. p[x]
GENL[tm] th                     From |- p[x1,...,xn] to |- !x1 .. xn. p[x1,...,xn]
GEN_ALL th                      From |- p[x1,...,xn] to |- !x1 .. xn. p[x1,...,xn], all variables
GEN_REWRITE_RULE cnvn [th] th   Rewrite conclusion of theorem using precise depth conversion
GSYM th                         Switch topmost equations, e.g. from |- !x. s[x] = t[x] to !x. t[x] = s[x]
INST[tm,tm] th                  Instantiate |- p[x1,...xn] to |- p[t1,...,tn]
INT_ARITH tm                    Linear arithmetic prover over Z
INT_OF_REAL_THM th              Map universal theorem from R to analog over Z
ISPEC tm th                     From |- !x. p[x] to |- p[t] with type instantiation
ISPECL[tm] th                   From |- !x1 .. xn. p[x1,...,xn] to |- p[t1,...,tn] with type instantiation
MATCH_MP th th                  From |- p ==> q and |- p' to |- q', instantiating first theorem to match
MK_COMB(th,th)                  From |- f = g and |- x = y to |- f(x) = g(y)
MP th th                        From |- p ==> q and |- p to |- q, no matching
ONCE_REWRITE_RULE[th] th        Rewrite conclusion of theorem once at topmost subterms
PART_MATCH tmfn th tm           Instantiate theorem by matching part of it to a term
PROVE_HYP th th                 From |- p and p |- q to |- q
REAL_ARITH tm                   Linear arithmetic prover over R
REFL tm                         Produce trivial theorem |- t = t
REWRITE_RULE[th] th             Rewrite conclusion of theorem with equational theorems
SPEC tm th                      From |- !x. p[x] to |- p[t]
SPECL[tm] th                    From |- !x1 .. xn. p[x1,...,xn] to |- p[t1,...,tn]
SPEC_ALL th                     From |- !x1 .. xn. p[x1,...,xn] to |- p[x1,...,xn]
SYM th                          From |- s = t to |- t = s
TAUT tm                         Prove propositional tautology like `p /\ q ==> p`
TRANS th th                     From |- s = t and |- t = u and |- s = u
UNDISCH th                      From |- p ==> q to p |- q
```

Inference rule with return type `thm list`

```
CONJUNCTS th                    From |- p1 /\ ... /\ pn to [|- p1; ...; |- pn]
```

Conversions (type `conv` = `term -> thm`)

```
BETA_CONV tm                    Reduce toplevel beta-redex |- (\x. s[x]) t = s[t]
CONTRAPOS_CONV                  From `p ==> q` give |- (p ==> q) <=> (~q ==> ~p)
GEN_BETA_CONV                   Reduce general beta-redex like |- (\(x,y). p[x,y]) (a,b) = p[a,b]
GEN_REWRITE_CONV cnvn [th]      Rewriting conversion using precise depth conversion
NUM_REDUCE_CONV                 Evaluate numerical expressions over N like `2 + 7 DIV (FACT 3)`
conv ORELSEC conv               Try to apply one conversion and if it fails, apply the other
REAL_RAT_REDUCE_CONV            Evaluate numerical expressions over R like `&22 / &7 - &3 * &1`
REWRITE_CONV[th]                Conversion to rewrite a term t to t' giving |- t = t'
REWR_CONV th                    Conversion to rewrite a term t once at top level giving |- t = t'
SYM_CONV                        Conversion to switch equations once |- P[s = t] <=> P[t = s]
conv THENC conv                 Apply one conversion then the other
TOP_DEPTH_CONV conv             Apply conversion once to top-level terms
```

Conversionals (type `conv -> conv`)

```
BINDER_CONV                     Apply conversion to body of quantifier etc.
LAND_CONV                       Apply conversion to LHS of binary operator, e.g. `s` in `s + t`
ONCE_DEPTH_CONV                 Apply conversion to first possible subterms top-down
RAND_CONV                       Apply conversion to rand of combination, e.g. x in f(x)
RATOR_CONV                      Apply conversion to rator of combination, e.g. f in f(x)
```

Tactics (return type `tactic`)

```
ABBREV_TAC tm                 Introduce abbreviation for t, from ?- p[t] to t = x ?- p[x]
ABS_TAC                       From ?- (\x. s[x]) = (\x. t[x]) to ?- s[x] = t[x]
ALL_TAC                       Tactic with no effect
ANTS_TAC                      From ?- (p ==> q) ==> r to ?- p and ?- q ==> r
AP_TERM_TAC                   From ?- f s = f t to ?- s = t
AP_THM_TAC                    From ?- f x = g x to ?- f = g
ARITH_TAC                     Tactic to solve linear arithmetic over N
ASM_CASES_TAC tm              Split ?- q into p ?- q and ~p ?- q
ASM_MESON_TAC[th]             Tactic for first-order logic including assumptions
ASM_REWRITE_TAC[th]           Rewrite goal by theorems including assumptions
ASM_SIMP_TAC[th]              Simplify goal by theorems including assumptions
BETA_TAC                      Reduce all beta-redexes in conclusion of goal
COND_CASES_TAC                From ?- P[if p then x else y] to p ?- p[x] and ~p ?- p[y]
CONJ_TAC                      Split ?- p /\ q into ?- p and ?- q
CONV_TAC conv                 Apply conversion to conclusion of goal
DISCH_TAC                     From ?- p ==> q to p ?- q
DISCH_THEN ttac               From ?- p ==> q to ?- q after using |- p
DISJ1_TAC                     From ?- p \/ q to ?- p
DISJ2_TAC                     From ?- p \/ q to ?- q
EQ_TAC                        Split ?- p <=> q into ?- p ==> q and ?- q ==> p
EVERY_ASSUM ttac              Apply function to each assumption of goal
EXISTS_TAC tm                 From ?- ?x. p[x] to ?- p[t]
EXPAND_TAC s                  Expand an abbreviation in a goal
FIRST_ASSUM ttac              Apply function to first possible assumption of goal
FIRST_X_ASSUM ttac            Apply function to and remove first possible assumption of goal
GEN_REWRITE_TAC cnvn [th]     Rewrite conclusion of goal using precise depth conversion
GEN_TAC                       From ?- !x. p[x] to ?- p[x]
INDUCT_TAC                    Apply ordinary mathematical induction to goal
LIST_INDUCT_TAC               Apply list induction to goal
MAP_EVERY atac [a]            Map tactic-producing function over a list of arguments, apply in sequence
MESON_TAC[th]                 Solve goal using first-order automation, ignoring assumptions
ONCE_REWRITE_TAC[th]          Rewrite conclusion of goal once at topmost subterms
tac ORELSE tac                Try to apply one tactic and if it fails, apply the other
POP_ASSUM ttac                Remove first assumption of goal and apply function to it
POP_ASSUM_LIST tltac          Remove assumptions of goal and apply function to it
REAL_ARITH_TAC                Tactic to solve linear arithmetic over R
REFL_TAC                      Solve trivial goal ?- t = t
REPEAT tac                    Apply a tactic repeatedly until it fails
REWRITE_TAC[th]               Rewrite conclusion of goal with equational theorems
RULE_ASSUM_TAC thfn           Apply inference rule to all hypotheses of goal
SET_TAC[th]                   Solve trivial set-theoretic goal like ‘x IN (x INSERT s)‘
SIMP_TAC[th]                  Simplify goal by theorems ignoring assumptions
SPEC_TAC(tm,tm)               From ?- p[t] to ?- !x. p[x]
STRIP_TAC                     Break down goal, ?- p /\ q to ?- p and ?- q etc. etc.
SUBGOAL_THEN tm ttac          Split off a separate subgoal
TRY tac                       Try a tactic but do nothing if it fails
tac THEN tac                  Apply one tactic then the other to all resulting subgoals
tac THENL [tac]               Apply one tactic then second list to corresponding subgoals
UNDISCH_TAC tm                From p ?- q to ?- p ==> q
USE_THEN s ttac               Apply function to assumption with particular label
X_GEN_TAC tm                  From ?- !x. p[x] to ?- p[y] with specified ‘y‘
```

Theorem-tactics (type `thm_tactic` = `thm -> tactic`)

```
ACCEPT_TAC                    Solve goal ?- p by theorem |- p
ANTE_RES_THEN ttac            Using |- p ==> q in goal p ?- r apply theorem-tactic to |- q
ASSUME_TAC                    Given |- p, from ?- q to p ?- q, no label on new assumption
CHOOSE_THEN ttac              Using |- ?x. p[x] apply theorem-tactic to |- p[x]
CONJUNCTS_THEN ttac           Using |- p /\ q apply theorem-tactic to |- p and |- q
CONJUNCTS_THEN2 ttac ttac     Using |- p /\ q apply respective theorem-tactics to |- p and |- q
DISJ_CASES_TAC                Use |- p \/ q, from ?- r to p ?- r and q ?- r
DISJ_CASES_THEN ttac          Use |- p \/ q, apply theorem-tactic to |- p and |- q separately
LABEL_TAC s                   Given |- p, from ?- q to p ?- q, labelling new assumption "s"
MATCH_ACCEPT_TAC              From |- p[x1,...,xn] solve goal ?- p[t1,...,tn] that's an instance
MATCH_MP_TAC                  Use |- p ==> q to go from ?- q' to ?- p', instantiation theorem to match
MP_TAC                        Use |- p to go from ?- q to ?- p ==> q
REPEAT_TCL ttacfn ttac        Apply theorem-tactical repeatedly until it fails
STRIP_ASSUME_TAC              Break theorem down into pieces and add them as assumptions
SUBST1_TAC                    Substitute equation in conclusion of goal, no matching
SUBST_ALL_TAC                 Substitute equation in hypotheses and conclusion of goal, no matching
X_CHOOSE_TAC tm               From |- ?x. p[x] and ?- q to p[y] ?- q, specified y
X_CHOOSE_THEN tm ttac         From |- ?x. p[x] apply theorem-tactic to |- p[y], specified y
```

tm	: term
[tm]	: term list
(tm,tm)	: term * term
[tm,tm]	: (term * term) list
tmfn	: term -> term
th	: thm
[th]	: thm list
(th,th)	: thm * thm
thfn	: thm -> thm
conv	: conv
cnvn	: conv -> conv
tac	: tactic
[tac]	: tactic list
ttac	: thm_tactic = thm -> tactic
tltac	: thm list -> tactic
ttacfn	: thm_tactical = thm_tactic -> thm_tactic
atac	: 'a -> tactic
[a]	: 'a list
s	: string