

Why bodies? Twelve reasons for including bodily expressions in affective neuroscience.

Journal:	<i>Philosophical Transactions B</i>
Manuscript ID:	RSTB-2009-0190
Article Type:	Review
Date Submitted by the Author:	24-Jul-2009
Complete List of Authors:	de Gelder, Beatrice; Tilburg University, Psychology
Issue Code: Click http://rstb.royalsocietypublishing.org/site/misc/issue-codes.xhtml target=_new>here to find the code for your issue.:	EMOTIONS
Subject:	Behaviour < BIOLOGY, Cognition < BIOLOGY, Neuroscience < BIOLOGY
Keywords:	bodily expression, affective neuroscience, emotions

1
2
3
4
5 **In: *Philosophical transactions of the Royal Society. B.***

6
7 *(in press)*

8
9
10 **Why bodies? Twelve reasons for including bodily expressions in affective**
11 **neuroscience.**
12
13

14
15
16
17
18 **Beatrice de Gelder**

19
20
21 *Laboratory of Cognitive and Affective Neuroscience, Tilburg University, The Netherlands and*
22 *Martinos Center for Biomedical Imaging, Massachusetts General Hospital, Charlestown,*
23 *Massachusetts and Harvard Medical School, Cambridge, USA.*
24
25

26
27
28 Why bodies? It is rather puzzling that given the massive interest in affective neuroscience in
29 the last decade it still seems to make sense to raise the question “Why bodies?” and to try to
30 provide an answer to it, as it the goal of this chapter. There are now hundreds of articles on
31 human emotion perception ranging from behavioural studies to brainimaging experiments.
32 Furthermore, decades of reports on affective disorders in neurological patient and clinical
33 studies of psychiatric populations have . The most cursory glance at the literature on emotion
34 in humans, now referred to with the umbrella term of social and affective neuroscience, shows
35 that over 95 % of them have used faces as stimuli. Of the remaining 5%, a few have used
36 scenes or auditory information including human voices, music or environmental sounds. But
37 by far the smallest number has looked into whole body expressions. As a rough estimate, a
38 search on PubMed today, May 1th 2009, yields 3521 hits for emotion X faces, 1003 hits for
39 emotion x music and 339 hits for emotion x bodies. When looking in more detail, the body x
40 emotion category in fact yields a majority of papers on wellbeing, nursing, sexual violence or
41 organ donation. But the number of cognitive and affective neuroscience studies of emotional
42 body perception as of today is lower than 20.
43
44
45
46
47
48
49
50
51
52
53
54

55
56
57 Why then have whole bodies and bodily expressions not attracted the attention of
58 researchers so far? The goal of this chapter is to contribute some elements for an answer to
59 this question. We believe that there is something to learn from the historical neglect of bodies
60

1
2
3 and bodily expressions. We next will address some historical misconceptions about whole
4 body perception and in the process we intend to provide not just an impetus for this kind of
5 work, but also to contribute in the process to a better understanding of the significance of the
6 affective dimension of behaviour, mind and brain as seen from the vantage point of bodily
7 communication. Subsequent sections discuss available evidence for the neurofunctional basis
8 of facial and bodily expressions as well as neuropsychological and clinical studies of bodily
9 expressions.
10
11
12
13
14
15
16

17 ***1. Bodily expressions are recognized as reliably as facial expressions.***

18
19 Prima facie, there is no historical explanation why faces have not captured much
20 attention. In fact two of the most illustrious theoreticians of emotion, Darwin and James,
21 discussed whole body expressions at great length. Darwin famously included postural
22 descriptions in his *The Expression of the Emotions in Man and Animals* (Darwin, 1872/1965),
23 and James (1932) investigated recognition of emotion with photographs of whole-body
24 posture. More recently, theoreticians of emotion like Tomkins (1995) and later Frijda (1988),
25 stressed the intimate link between emotion and action and were thus naturally led to
26 emphasize the importance of the body. If bodily expressions did never occupy centre stage in
27 emotion research, that may have been related to scepticism concerning consensus among
28 observers that has its roots not so much in theory than in empirical results dating from the first
29 generation of investigations of whole body stimuli. For example, Ekman (1965) performed
30 two studies on recognition of emotion from bodily expressions. But the results seem to have
31 led him to sharing this skepticism. He concentrated on facial expressions at the detriment of
32 bodily ones as chances to find evidence for emotion expression universals loomed larger in
33 the domain of facial expressions.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

But in recent decades researchers have taken up the issue of bodily expression
recognition and results from a number of behavioral experiments using independent stimulus
sets now allow us to conclude that recognition performance for bodily expressions is very
similar for face and body stimuli and this counts for both studies with static or dynamic
whole-body stimuli. Available studies whether focusing on recognition per se or preparing a
set of validated body stimuli, have indeed found a high degree of agreement among observers
(Atkinson et al., 2004; de Meijer, 1989; Dittrich et al., 1996; Hadjikhani and de Gelder, 2003;
Wallbott, 1998). A similar high consensus is found for videoclips depicting emotions or
emotion expressing instrumental actions (Grèzes et al., 2007; Pichon et al., 2008; 2009). As is

1
2
3 to be expected performance is lower when point light stimuli are used instead of full images.
4 Recognition rates are around 10% higher for dynamic images than for their still counterparts
5 and they increases when the face is not blanked out. These are all factors which may mainly
6 be related to the amount of information in the stimulus. In conclusion, when tested with
7 comparable stimuli and under comparable viewing conditions, there is as much consensus for
8 recognition of bodily expressions as there is for facial expressions.
9
10
11
12
13
14

15
16 Two other observations were made already in the earlier and more anecdotic reports
17 on bodily expressions and they have since taken on special importance. These relate to
18 casually observed phenomena that are nowadays referred to as respectively motor and
19 emotional contagion. Indeed, on the occasion of his own experiments W. James reported that
20 participants sometimes tended to imitate the posture they were looking at. Such emotional
21 motor reaction has since been studied systematically by Dimberg and collaborators. When
22 shown facial expressions and other affective pictures observers make emotion specific facial
23 movements. This is seldom visible to the naked eye and as James commented, observers do
24 not actually take the posture they are observing, -at least not in a way this is obvious to the
25 naked eye-, but rather experience some kind of kinesthesia in body parts that are normally
26 involved in the production of the observed posture. Furthermore, James also reported some
27 evidence for what has since been called emotion contagion. He noted that sometimes
28 observers seemed to experience the feelings and emotions typically associated with the
29 posture they saw and indicated that these feelings may be follow enacting the posture
30 observed (p. 419). Note that James reports these two observations, motor reaction and
31 emotional feeling reactions, as two separate phenomena. Of lately there has been a tendency
32 to conflate emotional motor reaction and motor contagion in a single phenomenon, emotional
33 contagion based on mirror neuron activation. But this obscures the fact that facial reactions do
34 not need to mimic the stimulus itself, but reflect its affective content whether it is a face or a
35 scene, or, an emotional voice (Magnee et al., 2006) or a bodily expression (Tamietto et al.,
36 2008; 2009). It also obscures the fact that there may be significant dissociations between the
37 emotion and the motor perception structures in the brain (Grezes & de Gelder, 2008).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 So far researchers have almost exclusively focused on measuring motor and emotional
57 contagion in the facial musculature. This type of work is consistent with the theoretical basis
58 of the FACS (Facial Action Coding System) developed by Ekman and collaborators. As to
59 spontaneous imitation of bodily postures by observers, we have been developing a BACS
60

1
2
3 (Bodily Action Coding system) based on whole body muscle movements (de Gelder ,van
4 Boxtel, in preparation) so far observed this indirectly.
5
6
7

8
9 To conclude this section, the notion that there is more consensus for recognition of
10 facial expressions than for bodily expressions has now turned out to be largely unfounded.
11 More importantly, some physiological and autonomic signatures of recognition do obtain as
12 well whether we view bodily or facial expressions. In the following sections we will see that
13 not only are bodily expressions well recognized, but they also trigger recognition under
14 conditions of limited attention and awareness in the same way as facial expressions do. On the
15 other hand, there are also important differences between facial and bodily expressions. We
16 will review differences between the two stimulus types and discuss why they are important.
17
18
19
20
21
22
23
24
25

26 *2. Understanding the sociocultural background of face centrism will enrich human* 27 *emotion research.* 28

29
30 Out of habit or out of principle, we tend to fill in automatically ‘facial’ when talking about
31 emotional expressions. This bias is very clearly reflected in the fact, already mentioned, that
32 studies of emotion recognition have hardly ever ventured away from the face. Yet it is a
33 truism that emotions are conveyed by a whole range of other cues besides the facial
34 expression. The continuity of facial expressions with postural, gestural and auditory signals
35 has tended to remain in the background. Before addressing specific issues it is worth asking
36 what may possibly be the roots of the face bias in our cultural and ideological heritage.
37
38
39
40
41
42
43

44 The emphasis on the face is present in western culture and art but not for example in
45 Islamic art. Well before neuropsychologists observed relative selective effects of lesions and
46 neurophysiologists recording from single cells provided evidence for specialised face
47 processes artists have focussed on the face. Traditionally the face is seen as the window of the
48 soul, it is our privileged access route to the thoughts and feelings of the people around us. In
49 their face we read what others think and feel and we see ourselves reflected, accepted or
50 rejected in their eyes. Faces presumably provide this information rapidly and automatically.
51 But there is also an added moral dimension to the facial communication channel. Indeed, we
52 do not merely learn about what a person thinks from watching his face, but at the same time
53 we evaluate his trustworthiness. And we seem to be able to do so at a glance. This theme has
54 been taken up recently in experiments investigating whether intuitive judgments of the
55
56
57
58
59
60

1
2
3 trustworthiness of human faces are accurate. There are now quite a few studies providing
4 evidence that are first impressions are strong and remain over time (Bar, et al., 2006). We are
5 quite good at telling trustworthiness at glance (Engell et al., 2007). But these two dimensions
6 of facial information reading, the emotional expression per se and the trustworthiness, are
7 traditionally linked. This cultural background is clear in the discussions of rapid facial
8 judgements of trustworthiness just mentioned but also in the notion that we can tell lies from
9 the face. Obviously, we also can tell higher order properties from body language as shown for
10 example for like deceit (Grèzes et al., 2004) or moral violation (Sinke et al., submitted).
11
12
13
14
15
16
17
18

19 The philosophical emotion literature from Aristotle to Spinoza, to name just these,
20 discusses emotions in the context of passion, reason and ethics. More fundamentally, this
21 special status of the face is linked to dualism, in recent times typically associated with
22 Descartes and his statement of the mind-body problem. If dualism is the notion that there is a
23 seemingly unbridgeable gap between our mental life and our material existence, then the face
24 falls on the side of the mind and the mental life while the body is relegated to the realm of the
25 machine. The face expresses the mind, but the body, as is typical of machines, does not have a
26 mind of its own and thus does not express anything. Just as the movements of the car do not
27 express the car's feelings or intentions, the movements of the body are equally mechanistic
28 and devoid of meaning.
29
30
31
32
33
34
35
36
37
38

39 Besides these, there are many other more directly cultural reasons why the body may
40 seem a less reliable source of affective information. Tradition, culture, religion and fashion
41 have a considerable impact on the public presentation of the body. Examples of extreme
42 influence are Greek aesthetics of ideal body proportions, the imperatives of the Victorian
43 dress code, or at the other extreme, the burqa completely hiding from sight all hints of facial
44 and bodily expression. All these factors together conspire to influence our attitude to bodies
45 by underscoring how its visual appearance can be manipulated and they end up casting doubt
46 on the body as a natural means of emotional communication.
47
48
49
50
51
52
53
54

55 Finally, consistent with the dualist framework just referred to and with the lingering
56 doubts about whether bodily expressions are easily recognized one understands that for an
57 universalist emotion theory the focus on facial expressions appears as the more promising.
58 And indeed over the last three decades the case for universal emotional expressions and
59
60

1
2
3 associated universal emotions has been argued with the help of analysis of facial expressions
4 most forcefully by Ekman and collaborators (e.g. this volume).
5
6
7
8
9

10 ***3. Investigations of bodies will extend the scope of face based research and provide***
11 ***evidence that human emotion theories based on studies of facial expression may generalize***
12 ***to other affective signals.***
13
14

15 It may very well turn out that current models of human emotion perception originally
16 developed for understanding facial expressions perception are just as valid for investigating
17 and understanding bodily expressions. There is very little evidence available at present to
18 support or refute this view.
19
20
21
22
23

24 On the positive side, one may argue that we already know from animal research that a
25 central structure in affective processes, the amygdala receives input from visual, auditory and
26 postural cues and is therefore likely to play a role in processing faces, bodily postures and
27 vocalizations as well. But AMG is one among other important structures involved in alerting
28 the organism to the presence of affective signals and preparing an adaptive response. For
29 example, the first brainimaging studies led to the impression that the role of the AMG in
30 emotion perception reflects a specialization for fearful facial expressions. Subsequent findings
31 have challenged this picture and pleaded in favor of sensitivity to salience or even to stimulus
32 ambiguity. Furthermore, AMG is widely connected to a number of other cortical and
33 subcortical brain structures (Amaral & Price, 1984.) And its embedding in these multiple
34 networks determines to a large extent its specific functional role in relation to the stimulus and
35 the behavioral context. It also determines the subjective emotional experience and the
36 behavioral consequences. For example, AMG is sensitive the presence of threat stimuli in the
37 environment whether or not the observer is not aware of them (Tamietto et al., 2009).
38
39
40
41
42
43
44
45
46
47
48
49
50

51 On the negative side, new investigations of other affective channels than the face may
52 challenge current face –based models. These issues are now beginning to be addressed and
53 some intriguing similarities and differences between the neural basis of facial and bodily
54 expressions emerge already. As direct comparisons become available, significant differences
55 between the neurofunctional basis of facial and bodily expressions are beginning to emerge.
56
57
58
59
60 An extensive overview of the currently available studies that have used behavioural,

1
2
3 electrophysiological and brainimaging methods is provide elsewhere (de Gelder et al., 2009).
4
5 Some main tendencies are briefly summarised in what follows here.
6
7

8
9 At the behavioural level there is clear evidence that both faces and bodies are
10 processed configurally rather than as an assemblage of features. This is assessed by measuring
11 the perceptual processes triggered when the stimuli are presented upside-down. The resulting
12 difference in performance is called the inversion effect, which refers to the loss of
13 performance when faces have to be recognized from upside down compared to upright
14 presented stimuli. Contrary to what is often assumed, this is not specific for faces. A similar
15 loss of performance is also observed for other stimuli like for example landscapes. Recent
16 findings show that recognition of faces and bodies presented upside-down is relatively more
17 impaired than recognition of inverted objects, like houses when each category is compared
18 with its own inverted counterpart (Reed et al., 2003).
19
20
21
22
23
24
25
26
27

28 Some electrophysiological studies have already been reported and others are under
29 way. The well known inversion effect measured in the time window of 150-200 ms and
30 labelled the N170 obtains similarly for faces and bodies. We established that in an ERP study
31 that used faces, bodies and shoes, each compared to their inverted counterpart (Stekelenburg
32 and de Gelder, 2004). This was confirmed and extended in a study using MRI constrained
33 MEG which allows a very good temporal resolution combined with a good spatial one. This
34 study showed very early inversion effects for faces and bodies between 70-100 ms
35 poststimulus with category-specific cortical distributions (Meeren et al., 2008). There is also
36 evidence that young infants are already sensitive to the orientation of body stimuli as
37 measured by EEG (Gliga and Dehaene-Lambertz, 2005).
38
39
40
41
42
43
44
45
46
47

48 Investigations of the neurofunctional basis of seeing bodily expressions have begun to
49 show that this activates the same brain areas that were are hitherto associated with the
50 perception of faces (for reviews, see de Gelder, 2006; Peelen and Downing, 2007). For
51 example, in the first report of the neural basis of perceiving bodily expressions we compared
52 neutral and fear expressions and found increased activity for fearful bodily expressions in
53 AMG and FG. The area that showed body responsiveness in FG was the same one as
54 identified in an separate study using a face localizer. Of course, a more fine grained analysis
55 of the fMRI signal may in turn show a partial separation as well as an area of overlap with
56 face and body sensitivity as has indeed been suggested in later studies (Kanwisher et al.,
57
58
59
60

1
2
3 1997). There is almost no evidence in the literature to answer this question. For this reason we
4
5 designed an fMRI study with the aim to investigate whether the brain shows distinctive
6
7 activation patterns for perception of faces and bodies.
8
9

10 We presented pictures of faces and bodies with blurred faces that either showed a
11
12 neutral, fearful or happy expression and instructed participants to categorize the stimuli. To
13
14 untangle brain activation related to faces and bodies, we compared how the brain responds to
15
16 both categories (irrespective of emotional expression). As expected given the part-whole
17
18 relation between bodies and faces, the results showed that the middle part of the fusiform
19
20 gyrus (FG) that is typically associated with the perception of facial identity, is more activated
21
22 for bodies than for faces (van de Riet et al., 2009). Previous studies have shown that there was
23
24 partial overlap between the face-selective and body selective region within the FG
25
26 (Hadjikhani and de Gelder, 2003; Peelen and Downing, 2005). In fact, viewing whole body
27
28 expressions elicited a wider network of brain areas compared to faces, including other areas
29
30 previously associated with perception of facial expressions, like STS. Other brain regions are
31
32 more active for bodies than for faces, the middle temporal/middle occipital gyrus (the so
33
34 called extra-striate body area, EBA (Downing et al., 2001), the superior occipital gyrus and
35
36 the parieto-occipital sulcus.
37

38 When the affective information conveyed by the bodies and faces, overall there is
39
40 comparably more activation for bodily than for facial expressions (Kret et al., submitted).
41
42 Interestingly, emotional body expressions activate cortical and subcortical motor areas like
43
44 caudate nucleus, putamen and inferior frontal gyrus (IFG), possibly reflecting the adaptive
45
46 action component implied in the body expression, which is less pronounced in facial
47
48 expressions (de Gelder et al., 2004). In a follow up study we presented video clips of dynamic
49
50 facial and bodily expressions that conveyed a neutral, fearful or angry expression instead of
51
52 static picture stimuli. The results were consistent with the previous study while broadening
53
54 the perspective: bodies compared to faces activated more areas than vice versa, including the
55
56 FG. Again, motor related areas were more activated by emotional body expressions (Kret et
57
58 al. submitted).
59

60
4. *Stimulating a more direct understanding of the significance of emotions as adaptive actions.*

1
2
3 Another benefit to be gained from using bodily expression stimuli is the broader emotion
4 perspective obtained by using affective signals that are in effect operational over longer
5 distances than faces. This also shifts the attention, away from personal identity shown by the
6 face and which may not always matter for rapid decoding of the expression, to action much
7 better conveyed by bodily expressions seen at considerable distance. A major difference
8 between facial and bodily expressions is that the latter can already be recognized from far
9 away while the former require the viewer to be nearby. This is potentially an important
10 difference between how facial and bodily expressions play their communicative role and it
11 should have consequences for the specific information conveyed.
12
13
14
15
16
17
18
19

20
21 Focussing on facial expressions seems tends to make us refer to a person's mental states.
22 But focussing on bodily expressions directs attention to a person or a group's actions. When
23 we talk about emotions, ascribe emotions we have in sight of in mind we implicitly seem
24 naturally to refer to mental states of the persons whose face we have in sight (or in mind). But
25 when we refer to emotions we see expressed in the body it is more frequent to have in mind
26 an action. Therefore, when due to distance of other peripheral visual conditions or
27 impairments we are unable to tell the emotional state from reading the face, we can still
28 perfectly clearly read the action from the sight of the body.
29
30
31
32
33
34
35
36

37 ***5. To contribute to some long standing debates on why facial expressions in isolation are***
38 ***often recognized less than perfectly.***
39

40
41 Common sense tends to hold that we read facial expressions like we read words on a
42 page, meaning that we directly and unambiguously access the meaning. But as is often the
43 case, the expressly held common sense beliefs and what people routinely do when they
44 behave commonsensical are two different things. In fact, in daily life we only seem to hold to
45 the belief that a facial expression is unambiguous in a few extreme circumstances, like for
46 example in case of a really menacing fury or of a panic-stricken expressions. Most of the time
47 the anger and fear faces we see do leave some room for interpretation as is increasingly
48 evidenced by semantic effects and context effects on face recognition (Feldman-Barrett, 2009;
49 for a review, see de Gelder & vanden Stock, 2009).
50
51
52
53
54
55
56
57

58 Yet as theorists, whether reasoning from common sense principles or from available
59 scientific data, we hold on to a preferred beliefs in basic emotional expressions universally
60 represented by some facial expressions. The notion that these universal or 'basic emotions'

1
2
3 expressions are the rock bottom of our mental life thus mirrors our belief that certain emotion
4 category labels correspond to universal mental states with an unique associated biological
5 basis and with evolutionary defined triggers. But since researchers began to use the stimulus
6 set provided by Ekman et al (1976), less than perfect recognition rates were reported. For
7 example, recognition rates are rarely above 80% and they often tend to be comparatively
8 lower for fear, which is otherwise seen as the best candidate for a hardwired mental state
9 expression cum biological substrate. This brings us to the next issue, that some of these basic
10 mental states are most clearly expressed by the face while others are least ambiguous when
11 expressed by the whole body.
12
13
14
15
16
17
18

19
20
21 ***6. To address situations where facial and bodily expressions do not provide the same***
22 ***meaning as when facial and bodily cues combine, interact, conflict. To provide the missing***
23 ***context to theories of face perception***
24
25

26 For a while "Headless Body in Topless Bar" counted as one of the funniest lines to have
27 appeared in US newspapers. But headless bodies and bodiless heads do figure only in crime
28 catalogues and police reports and are not part of our daily experience, at the very least not part
29 of the daily experience that constitutes the normal learning environment in which we acquire
30 our face and body expertise. Yet, except for a few isolated studies (de Gelder et al., 2006;
31 Righthart & de Gelder, 2006; Mobbs et al., 2006; for an extended review and discussion, see
32 de Gelder & van den Stock, in press) the literature on face recognition has not yet addressed
33 the issue of context effects in face perception. By 'context' we mean here the whole
34 naturalistic environment that is almost always present when we encounter a face).
35
36
37
38
39
40
41
42

43
44 Perception of facial expression is influenced by whatever expression the body shows.
45 A popular notion is that our body language gives away our real feelings, for example in
46 situations where we manage to control our facial expression. Typical examples are when one
47 is trying to keep a poker face in situations of social control and dominance and stress. We
48 don't show anger or nervousness, we smile all the way through the conversation or the
49 interview, however annoying or unenlightening the questions may be. Even children are
50 Research on the simultaneous perception of faces and bodies is still sparse. Two behavioural
51 studies directly investigated how our recognition of facial expressions is influenced by
52 accompanying whole body expressions (Meeren et al., 2005; Van den Stock et al., 2007).
53 Meeren et al. (2005) combined angry and fearful facial expressions with angry and fearful
54 whole body expressions to create both congruent (fearful face on fearful body and angry face
55
56
57
58
59
60

1
2
3 on angry body) and incongruent (fearful face on angry body and angry face on fearful body)
4 realistically looking compound stimuli. These were briefly (200ms) presented one by one
5 while the participants were instructed to categorize the emotion expressed by the face and
6 ignore the body. The results showed that recognition of the facial expression was biased
7 towards the emotion expressed by the body language, as reflected by both the accuracy and
8 reaction time data. In a follow-up study, facial expressions that were morphed on a continuum
9 between happy and fearful were once combined with a happy and once with a fearful whole
10 body expression (Van den Stock et al., 2007). The resulting compound stimuli were presented
11 one by one for 150ms, while the participants were instructed to categorize the emotion
12 expressed by the face in a 2 alternative forced choice paradigm (fear or happiness). Again, the
13 ratings of the facial expressions were influenced towards the emotion expressed by the body
14 and this influence was highest for facial expressions that were most ambiguous (expressions
15 that occupied an intermediate position on the morph continuum). Evidence from EEG-
16 recordings that were collected during the experiment shows that the brain responds to the
17 emotional face-body incongruence as early as 115ms post stimulus onset (Meeren et al.,
18 2005).

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34 ***7. To understand emotion specificity of affective signals. The relative importance of***
35 ***face vs. body may be a function of the specific emotional significance conveyed.***

36 To repeat a truism, emotions are complex, rich and multilevel phenomena. And in spite of the
37 all the research effort devoted to it, many researchers have the impression that progress is
38 slow. This has led some authors for example LeDoux (1996) to argue for a research strategy
39 of concentrating on one emotion, in this case, fear, rather than trying to make progress on all
40 fronts at the same time. But this has not stopped researchers from periodically advancing
41 general theories. For example, a powerful impetus of emotion research in the last years tries
42 to capitalize on the potential of mirror neuron activation in the brain. The specific finding that
43 prompted this generalisation was a study of disgust and the study used video clips of facial
44 expressions of disgust (Wicker et al., 2003). Disgust is clearly an emotion that centres around
45 activity in the mouth region and thus privileges the face as a bearer. The adaptive action
46 component of disgust is unlikely to involve much movement of the lower limbs. But other
47 emotions are more powerfully expressed in the arms and the lower limbs of the whole body
48 that in the face. Aggression is a case in point. When viewing aggressive body pictures
49 observers spend most of the time looking at the hands (Fridin et al., 2009). Therefore, a
50 comparison of emotional expressions that is sensitive to the specific emotion is likely to
51
52
53
54
55
56
57
58
59
60

1
2
3 reveal the relative prominence of the face vs. the body depending on the emotion considered
4 and differential contribution from body parts to the specific emotion.
5
6
7

8
9 **8. To understand gender specificity of the importance of body vs. face signals.**

10 Notwithstanding widespread stereotypes about gender specificity in emotions and body
11 related issues there are so far only very few studies available on these issues. We designed
12 study with the goal to compare the neural bases dedicated to processing facial and bodily
13 expressions using video clips of faces and bodies expressing threatening emotions (fear and
14 anger) and to assess the influence of gender from both the observer and the actor (Kret et al.,
15 2009). Male and female participants recognized all expressions very well (mean percentage
16 correct: 90% for fear, 95% for anger). There were no significant differences between accuracy
17 rates for male or female participants, and there was also no difference in the recognition of
18 male or female actors. In contrast to these behavioural results, we did find some striking
19 gender effects in the fMRI results. Activation of classical subcortical emotion areas (AMG,
20 hippocampus, putamen, thalamus and basal ganglia) only showed up when the observers,
21 both male and female, perceived threat from male actors. In male participants, the dorsal
22 stream was primarily involved in perceiving threat, -especially from angry male- body
23 language. Strikingly, the superior temporal sulcus, an area that has often found to be
24 implicated in emotional processing was absolutely not influenced by the type of emotion in
25 the female participants.
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 When females perceived *male threatening body language*, they activate the dorsal
41 pathway and a network that involves action preparation and observation. Apart from enhanced
42 activity in the visual areas, there are activations in precuneus, inferior and superior parietal
43 lobe (action observation), precentral gyrus, SMA and motor cingulate cortex (posterior and
44 anterior), as well as in the caudate nucleus and in the putamen (action preparation). The
45 activity found in the motor cingulate cortex corresponds to the area known to be involved in
46 arm movements (Pickard and Strick, 1996). This may imply that female participants felt the
47 urge to protect themselves (for example covering their faces, with their arms when watching
48 male body expressions of threat).
49
50
51
52
53
54
55
56
57

58 **9. Bodily expressions are perceived in a multisensory environment, combine with**
59 **audio signals, a phenomenon hitherto only reported for facial expressions.**
60

1
2
3 An interesting argument in favour of the primacy of the face may be that facial
4 expressions form a more natural pair with the voice. Research has shown that recognition of
5 the emotion in the target modality (the facial expression) is typically influenced towards the
6 emotion expressed in the task irrelevant modality (the affective prosody of the voice). (e.g., de
7 Gelder and Vroomen, 2000). In this type of study two modalities are typically combined to
8 create emotionally congruent and incongruent face–voice pairs in order to provide a window
9 into the integration process (de Gelder & Bertelson, 2003).

10
11
12 It is worth noting that the argument concerning the naturalness and ecological validity
13 of considering visual stimuli in a multisensory context applies just as well to the whole body
14 than to the face only. A trainee singer trying to sing only with the upper body will quickly
15 experience the limitations of that approach. Similarly, in trying to shout with the arms folded
16 over your chest one quickly becomes aware that vocalisations are produced by the body!

17
18
19 In two recent studies we have taken this issue beyond facial expressions and
20 investigated affective crossmodal influences in whole body expressions (Van den Stock et al.,
21 2007). We investigated naturalistic actions that are part of everyday life and focused on
22 instrumental actions, like grasping and drinking. Our data showed that affective crossmodal
23 effects occur with body–voice pairs and are thus very similar to previous findings about
24 combined perception of face-voice pairs (Van den Stock, 2008).

25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41 ***10. Perception or recognition of bodily expressions does not require full attention nor does***
42 ***it require that the visual stimulus be consciously seen.***

43
44 Over the last decades a number of research reports have concluded that emotional
45 information can be processed without observers being aware of it (Barrett et al., 2007; Kunst-
46 Wilson and Zajonc, 1980). But nonconscious affect perception has almost exclusively been
47 investigated with the use of facial expressions, either on their own or in combination with
48 other visual stimuli. Many studies now provide direct and indirect evidence for visual
49 discriminations of facial expressions in the absence of visual awareness of the stimulus (e.g.
50 de Gelder et al., 1999; Dimberg et al., 2000; Esteves et al., 1994a; Jolij and Lamme, 2005).
51 Theoretical models have been advanced arguing that separate pathways may sustain conscious
52 and nonconscious emotional perception (LeDoux, 1996; Morris et al., 1998). The notion of
53 separate pathways has to some extent been accepted by the community at large although some
54 core findings are still a matter of debate (Duncan and Barrett, 2007; Pessoa, 2005).

1
2
3
4
5 An issue that has so far not received much attention is whether there also exists
6 nonconscious emotional perception for bodily expressions. Data from patients with
7 hemianopia indicate that they may reliably discriminate between bodily expressions which
8 they are unable to see due to striate cortex lesion (de Gelder and Hadjikhani, 2006) or due to
9 an attentional disorder following parietal lesion (Tamietto and de Gelder, 2008) and, more
10 radically still, in patients with cortical blindness (Tamietto et al., 2009).
11
12
13
14
15
16

17 Other indicators besides behavioural measures also provide evidence for automatic
18 processing. A striking example is represented by the spontaneous tendency to synchronize our
19 facial expressions with those of another person during face-to-face situations. This
20 phenomenon of emotional contagion (Hatfield et al., 1994) is now widely observed but it is
21 still poorly understood. Recent proposals link emotional contagion directly to motor
22 resonance (i.e., stimulus/response motor matching) (Carr et al., 2003; Dimberg et al., 2000).
23
24
25
26
27
28

29 A directly related issue concerns the degree of automaticity of emotional contagion
30 and the role of visual awareness of the eliciting stimulus in the unfolding of affective
31 reactions at different levels of emotional experience. Available evidence shows that
32 nonconscious perceptual mechanisms are sufficient for processing emotional signals, most
33 notably so far facial expressions. The clearest evidence for processing without visual stimulus
34 awareness is obtained in patients with lesions to the primary visual cortex (V1). These
35 patients reliably discriminate the affective valence of facial expressions projected to their
36 clinically blind visual field by guessing (*affective blindsight*), despite having no conscious
37 perception of the stimuli they are responding to (Morris et al., 2001; Pegna et al., 2005).
38 Nevertheless, it is unknown at present whether nonconscious perception of emotions in
39 cortically blind patients may lead to spontaneous facial reactions or to other physiological
40 changes typically associated to emotional responses.
41
42
43
44
45
46
47
48
49
50

51
52 ***11. To reveal the neurofunctional correlates of category specific cq. body specific***
53 ***processes and deficits.***
54

55 A strong impetus for category specificity of neural substrates comes from
56 neuropsychological reports of patients with brain damage acquired in adulthood. There is a
57 well known neuropsychological deficit related to impaired face recognition, labelled
58 prosopagnosia. These patients are impaired in recognizing faces, and very often have no
59
60

1
2
3 recognition at all of an individual by the face only. Brain damage occurring in the normally
4 developed brain that affects face perception is often localized in occipitotemporal cortex and
5 temporal cortex (midfusiform gyrus and inferior occipital gyrus) unilaterally or bilaterally.
6
7 The developmental counterpart of acquired prosopagnosia is now also often reported. There is
8 substantial similarity between acquired and developmental prosopagnosia at the behaviour
9 level but there are many other differences (see de Gelder and Rouw, 2000) for a comparison).

10
11
12
13
14
15
16 It is important to specify the nature of the disorder though and this is still a matter of
17 debate. The short definition of prosopagnosia characterizes it as a deficit in face recognition.
18 But this is too broad and also too a-specific. We are in fact dealing with a deficit that affects
19 recognition of personal identity from the sight of the face. Other dimensions of face
20 information are processed mostly normally, like emotional expression, visual speech or
21 gender. In fact, a good means of defining the typical face deficit of prosopagnosics is by
22 establishing that there exists a dissociation between the different dimensions of face
23 perception some of which are impaired while others are intact. On the other hand, there are
24 too date only very few cases of pure prosopagnosia, where the perception and recognition
25 deficit is restricted to the face and does not affect in any way perception and recognition of
26 other object categories. Thus on this axis also, a dissociation must be established requiring
27 that the perception and recognition impairment is impairment is not present for non face
28 stimuli. To establish the presence of developmental prosopagnosia the same dimensions of
29 dissociation need to be assessed.

30
31
32
33
34
35
36
37
38
39
40
41
42 Using body stimuli offers a chance to advance the debate on category specificity. As a
43 matter of fact there are very few objects other than faces for which strong claims about
44 category specific representation have been made. An interesting object category not used so
45 far concerns human bodies. Recently, it has been shown in normal subjects that perceiving
46 human bodies or body parts activates an area in extrastriate cortex, labeled extrastriate body
47 area (EBA) (Downing et al., 2001) and more recently a second body specific area in the FG
48 (Hadjikhani and de Gelder, 2003; Peelen and Downing, 2005) overlapping partially with the
49 face-sensitive one and it has been termed the fusiform body area (FBA). These behavioral and
50 neuro-functional similarities between perceiving faces and bodies in neurologically normal
51 observers raise the issue how bodies are processed in DP. Our second main finding concerns
52 the categorical specificity of face vs. body perception in DPs. We compared the activation of
53 body conditions in the face selective regions and of the face conditions in the body selective
54
55
56
57
58
59
60

1
2
3 regions between both groups. On the one hand, our findings indicate that perceiving neutral
4 faces results in a higher activation of EBA in the DP group, compared to the control group.
5 Combined with the lower activation in FFA, this increased activation in EBA might indicate
6 an anomalous processing route in the brains of DPs. It may be the case that (neutral) faces are
7 processed in the areas more dominantly dedicated to body perception. Combined with the
8 lower activation in FFA, this increased activation in EBA might indicate an anomalous
9 processing route in the brains of DPs. It may be the case that (neutral) faces are processed in
10 the areas more dominantly dedicated to body perception. On the other hand, we find a higher
11 activation for perceiving bodies in IOG. These combined findings indicate that the neural
12 correlates of perceiving faces and bodies, as manifested in IOG and EBA show a lower degree
13 of specificity in DP.
14
15
16
17
18
19
20
21
22
23

24
25 ***12. Investigations of bodily expressions will enrich basic clinical research and lead to the***
26 ***development of new observational and diagnostic tools.***
27

28 Many studies of emotional communication disorders have reported deficits in face
29 recognition in clinical populations as well as in psychiatric disorders. These include autism
30 spectrum subjects, schizophrenics, mood disorders like depression and high anxiety
31 individuals. Autism spectrum disorder (ASD) is usually behaviorally defined as characterized
32 by mild to severe impairments in communication and reciprocal social interaction as well as
33 repetitive and stereotyped behaviors. Daily observations document difficulties ASD subjects
34 have in recognizing and appropriately reacting to other people's emotions, whether they are
35 communicated by facial expressions, vocal tone, gestures or bodily postures. Some of these
36 characteristics have been documented experimentally but much debate remains. Earlier
37 behavioral studies did not consistently find emotion perception deficits (Braverman et al.,
38 1989; Capps et al., 1992; Davies et al., 1994; Hobson et al., 1988; Macdonald et al., 1989;
39 Tantam et al., 1989), but see (Baron-Cohen et al., 1997; Gepner et al., 2001; Grossman et al.,
40 2000; Ozonoff et al., 1990) recent studies taking a more fine-grained approach have
41 documented emotion recognition impairments mainly in the perception of negative emotions,
42 especially fear (Ashwin et al., 2006; Baron-Cohen et al., 2000; Corden et al., 2006; Dawson et
43 al., 2004; Gaigg and Bowler, 2007; Humphreys et al., 2007; Welchew et al., 2005).
44
45
46
47
48
49
50
51
52
53
54
55
56

57
58 The above illustrates that to date, research on emotion and social communications
59 disorders has focused primarily on impairments in the neurofunctional processes associated
60 with viewing facial expressions. In view of reports that ASD may avoid attending to the face,

1
2
3 investigations of other channels of communication look particularly promising. We recently
4 performed two studies on bodily expression processing, one using still images (Hadjikhani,
5 2009) and another videoclips (Grèzes et al., 2009). The main finding of the first study using
6 still bodies is that brain activation patterns in individuals with ASD do not show evidence of
7 differentiation between bodily expressions of fear and bodies engaged in neutral actions. This
8 finding suggests an abnormality in the network of brain areas that are normally engaged in the
9 perception of bodily expressed emotions in NT individuals, and is consistent with recent
10 behavioral findings of Hubert et al. (2007) who reported normal perception of point-light
11 displays of neutral actions in ASD, but abnormal perception of emotions.
12
13
14
15
16
17
18
19

20
21 In the study using video clips of neutral and fear expressing whole body actions
22 normal perception of dynamic actions in ASD was also observed. Yet there were clear
23 anomalies linked to a failure to grasp the emotional dimension of the actions. We measured
24 brain activity using fMRI during perception of fearful or neutral actions and showed that
25 whereas similar activation of brain regions known to play a role in action perception was
26 revealed in both autistics and controls, autistics failed to activate amygdala, inferior frontal
27 gyrus and premotor cortex when viewing gestures expressing fear. Our results support the
28 notion that dysfunctions in this network may contribute significantly to the characteristic
29 communicative impairments present in autism. We observed that ASD subjects fail to engage
30 cerebral regions involved in grasping the emotional meaning of the actions they view. We
31 suggest that this deficit may reflect an important failure of the mechanisms that control
32 normal behavioral responses to emotional signals in the behavior of others. The ensuing
33 deficiency in the appraisal of emotional cues may lead to the inappropriate behavioral
34 responses and the social difficulties that are characteristic of this population. This suggestion
35 takes us well beyond conclusions reached in studies about communication deficits using facial
36 expressions because it allows specific hypothesis about social interactive impairments that are
37 so clearly present in many ASD individuals which are obviously more far reaching than the
38 unwillingness to attend to the face. This traditional stumble block for measuring emotion
39 recognition in ASD can be overcome by using bodily expressions instead.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 ***Conclusion.***

57
58 We have reviewed a series of arguments in favor of substantially extending and
59 enriching current human emotion theories adding investigations of bodily expressions. We
60 have also highlighted the importance of new research on bodily expressions for theories that

1
2
3 consider emotions to be closely linked to adaptive action. Finally, we have discussed some
4 recent studies illustrating the potential of bodily expression research for neuropsychological
5 investigations as well as for clinical research.
6
7
8
9

10
11
12 *Acknowledgements.* The study was partly supported by Human Frontier Science Program
13 HFSP - RGP0054/2004-C and by EC-contract number FP6-NEST-2005-Path-IMP-043403.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

REFERENCES

- Ashwin, C., Chapman, E., Colle, L., and Baron-Cohen, S. (2006). Impaired recognition of negative basic emotions in autism: a test of the amygdala theory. *Soc Neurosci* 1, 349-363.
- Atkinson, A. P., Dittrich, W. H., Gemmell, A. J., and Young, A. W. (2004). Emotion perception from dynamic and static body expressions in point-light and full-light displays. *Perception* 33, 717-746.
- Bar, M., Neta M, Linz, H. (2006). Very first impressions. *Emotion* 6 (2), 269-278.
- Baron-Cohen, S., Ring, H. A., Bullmore, E. T., Wheelwright, S., Ashwin, C., and Williams, S. C. (2000). The amygdala theory of autism. *Neurosci Biobehav Rev* 24, 355-364.
- Baron-Cohen, S., Wheelwright, S., and Jolliffe, T. (1997). Is there a "language of the eyes"? Evidence from normal adults and adults with autism or Asperger syndrome. *Visual Cognition* 4, 311-331.
- Barrett, L. F., Ochsner, K. N., and Gros, J. J. (2007). On the automaticity of emotion. In *Social psychology and the unconscious: the automaticity of higher mental processes.*, J. Bargh, ed. (New York, Psychology Press).
- Braverman, M., Fein, D., Lucci, D., and Waterhouse, L. (1989). Affect comprehension in children with pervasive developmental disorders. *J Autism Dev Disord* 19, 301-316.
- Capps, L., Yirmiya, N., and Sigman, M. (1992). Understanding of simple and complex emotions in non-retarded children with autism. *J Child Psychol Psychiatry* 33, 1169-1182.
- Carr, L., Iacoboni, M., Dubeau, M. C., Mazziotta, J. C., and Lenzi, G. L. (2003). Neural mechanisms of empathy in humans: a relay from neural systems for imitation to limbic areas. *Proc Natl Acad Sci U S A* 100, 5497-5502.
- Corden, B., Critchley, H. D., Skuse, D., and Dolan, R. J. (2006). Fear recognition ability predicts differences in social cognitive and neural functioning in men. *J Cogn Neurosci* 18, 889-897.
- Darwin, C. (1872/1965). *The expressions of emotions in Man and Animals*. John Marry, London.
- Davies, S., Bishop, D., Manstead, A. S., and Tantam, D. (1994). Face perception in children with autism and Asperger's syndrome. *J Child Psychol Psychiatry* 35, 1033-1057.
- Dawson, G., Webb, S. J., Carver, L., Panagiotides, H., and McPartland, J. (2004). Young children with autism show atypical brain responses to fearful versus neutral facial expressions of emotion. *Dev Sci* 7, 340-359.
- de Gelder, B. (2006). Towards the neurobiology of emotional body language. *Nature Reviews Neuroscience* 7, 242-249.
- de Gelder, B., and Bertelson, P. (2003). Multisensory integration, perception and ecological validity. *Trends Cogn Sci* 7, 460-467.
- de Gelder, B., and Hadjikhani, N. (2006). Non-conscious recognition of emotional body language. *Neuroreport* 17, 583-586.
- de Gelder, B., and Rouw, R. (2000). Configural face processes in acquired and developmental prosopagnosia: evidence for two separate face systems? *Neuroreport* 11, 3145-3150.
- de Gelder, B., Snyder, J., Greve, D., Gerard, G., and Hadjikhani, N. (2004). Fear fosters flight: A mechanism for fear contagion when perceiving emotion expressed by a whole body. *The Proceedings of the National Academy of Sciences Online (US)* 101, 16701-16706.
- de Gelder, B. & vanden Stock, J. (2009). Real faces, real emotions: perceiving facial expressions in naturalistic contexts of voices, bodies and scenes. In Calder, A., Haxby, J., Johnson, M., & Rhodes, G. (Eds.). *The Handbook of Face Perception*. Oxford University Press.
- de Gelder, B., and Vroomen, J. (2000). The perception of emotions by ear and by eye. *Cognition and Emotion* 14, 289-311.

- 1
2
3 de Gelder, B., Vroomen, J., Pourtois, G., and Weiskrantz, L. (1999). Non-conscious
4 recognition of affect in the absence of striate cortex. *Neuroreport* 10, 3759-3763.
5
6 de Meijer, M. (1989). The contribution of general features of body movement to the
7 attribution of emotions. *J Nonverbal Behavior* 13, 247-268.
8
9 Dimberg, U., Thunberg, M., and Elmehed, K. (2000). Unconscious facial reactions to
10 emotional facial expressions. *Psychological Science* 11, 86-89.
11
12 Dittrich, W. H., Troscianko, T., Lea, S. E., and Morgan, D. (1996). Perception of emotion
13 from dynamic point-light displays represented in dance. *Perception* 25, 727-738.
14
15 Downing, P. E., Jiang, Y., Shuman, M., and Kanwisher, N. (2001). A cortical area selective
16 for visual processing of the human body. *Science* 293, 2470-2473.
17
18 Duncan, S., and Barrett, L. F. (2007). The role of the amygdala in visual awareness. *Trends in*
19 *Cognitive Sciences* 11, 190-192.
20
21 Ekman, P. (1965). Differential communication of affect by head and body cues. *Journal of*
22 *personality and social psychology* 2 (5), 926-735.
23
24 Ekman, P., & Friesen, W.V. (1976) *Pictures of Facial Affect*. Palo Alto, CA: Consulting
25 *Psychologists Press*.
26
27 Engell, A.D., Haxby, J.V. and Todorov, A. (2007). Implicit trustworthiness decisions:
28 automatic coding of face properties in the human amygdala. *Journal of Cognitive*
29 *Neuroscience* 19 (9), 1508-5019.
30
31 Esteves, F., Dimberg, U., and Öhman, A. (1994a). Automatically elicited fear: conditioned
32 skin conductance responses to masked facial expressions. *Cognition and Emotion* 8, 393 -
33 413.
34
35 Esteves, F., Parra, C., Dimberg, U., and Öhman, A. (1994b). Nonconscious associative
36 learning: Pavlovian conditioning of skin conductance responses to masked fear-relevant facial
37 stimuli. *Psychophysiology* 31, 375-385.
38
39 Frijda, N. (1988). The laws of emotion. *American Psychologist* 43 (5), 349-358.
40
41 Gaigg, S. B., and Bowler, D. M. (2007). Differential fear conditioning in Asperger's
42 syndrome: implications for an amygdala theory of autism. *Neuropsychologia* 45, 2125-2134.
43
44 Gepner, B., Deruelle, C., and Grynfeldt, S. (2001). Motion and emotion: a novel approach to
45 the study of face processing by young autistic children. *J Autism Dev Disord* 31, 37-45.
46
47 Gliga, T., and Dehaene-Lambertz, G. (2005). Structural encoding of body and face in human
48 infants and adults. *Journal of Cognitive Neuroscience* 17, 1328-1340.
49
50 Grèzes, J., Frith, C. and Passingham, R.E. (2004). Brain mechanisms for inferring deceit in
51 the actions of others. *Journal of Neuroscience* 24, 5500-5505.
52
53 Grèzes, J., Pichon, S., and de Gelder, B. (2007). Perceiving fear in dynamic body expressions.
54 *Neuroimage* 35, 959-967.
55
56 Grèzes, J., Wicker, B., Berthoz, S., and de Gelder, B. (2009). A failure to grasp the affective
57 meaning of actions in autism spectrum disorder subjects. *Neuropsychologia* 47, 1816-1825.
58
59 Grossman, E., Donnelly, M., Price, R., Pickens, D., Morgan, V., Neighbor, G., and Blake, R.
60 (2000). Brain areas involved in perception of biological motion. *J Cogn Neurosci* 12, 711-
720.
Hadjikhani, N., and de Gelder, B. (2003). Seeing fearful body expressions activates the
fusiform cortex and amygdala. *Curr Biol* 13, 2201-2205.
Hadjikhani, N., Joseph, R.M., Manoach, D.S., Naik, P., Snyder, J., Dominick, K., Hoge, R.,
Van den Stock, J., Tager Flusberg, H., de Gelder, B. (2009). Body expressions of emotion do
not trigger fear contagion in autism spectrum disorder. *Soc Cogn Affect Neurosci* 4, 70-78.
Hatfield, E., Cacioppo, J. T., and Rapson, R. L. (1994). *Emotional Contagion* (Cambridge,
Cambridge University Press).
Hobson, R. P., Ouston, J., and Lee, A. (1988). What's in a face? The case of autism. *Br J*
Psychol 79, 441-453.

- 1
2
3 Hubert, B., Wicker, B., Moore, D.G., et al. (2007). Brief report: recognition of emotional and
4 non-emotional biological motion in individuals with autistic spectrum disorders. *Journal of*
5 *autism and developmental disorders*, 37, 1386–92.
- 6 Humphreys, K., Minshew, N., Leonard, G. L., and Behrmann, M. (2007). A fine-grained
7 analysis of facial expression processing in high-functioning adults with autism.
8 *Neuropsychologia* 45, 685-695.
- 9 Jolij, J., and Lamme, V. A. (2005). Repression of unconscious information by conscious
10 processing: evidence from affective blindsight induced by transcranial magnetic stimulation.
11 *Proceedings of the National Academy of Sciences of the United States of America* 102,
12 10747-10751.
- 13 Kanwisher, N., McDermott, J., and Chun, M. M. (1997). The fusiform face area: a module in
14 human extrastriate cortex specialized for face perception. *J Neurosci* 17, 4302-4311.
- 15 Kret, M.E., Grèzes, J., Pichon, S. and de Gelder, B. Gender specific brain activations for
16 perceiving threat from dynamic faces and bodies. (submitted)
- 17 Kunst-Wilson, W. R., and Zajonc, R. B. (1980). Affective discrimination of stimuli that
18 cannot be recognized. *Science* 207, 557-558.
- 19 LeDoux, J. E. (1996). *The emotional brain: The mysterious underpinnings of emotional life*
20 (New York, Simon and Schuster).
- 21 Macdonald, H., Rutter, M., Howlin, P., Rios, P., Le Conteur, A., Evered, C., and Folstein, S.
22 (1989). Recognition and expression of emotional cues by autistic and normal adults. *J Child*
23 *Psychol Psychiatry* 30, 865-877.
- 24 Meeren, H. K., Hadjikhani, N., Ahlfors, S. P., Hamalainen, M. S., and de Gelder, B. (2008).
25 Early category-specific cortical activation revealed by visual stimulus inversion. *PLoS ONE*
26 3, e3503.
- 27 Meeren, H. K., van Heijnsbergen, C. C., and de Gelder, B. (2005). Rapid perceptual
28 integration of facial expression and emotional body language. *Proc Natl Acad Sci U S A* 102,
29 16518-16523.
- 30 Mobbs et al. The Kuleshov Effect: the influence of contextual framing on emotional
31 attributions. *Soc Cogn Affect Neurosci*. 2006 1(2):95-106.
- 32 Morris, J. S., DeGelder, B., Weiskrantz, L., and Dolan, R. J. (2001). Differential
33 extrageniculostriate and amygdala responses to presentation of emotional faces in a cortically
34 blind field. *Brain* 124, 1241-1252.
- 35 Morris, J. S., Ohman, A., and Dolan, R. J. (1998). Conscious and unconscious emotional
36 learning in the human amygdala. *Nature* 393, 467-470.
- 37 Müller, J. P. (1840). *Handbuch der Physiologie des Menschen* (Coblenz, H. Ischer).
- 38 Ozonoff, S., Pennington, B. F., and Rogers, S. J. (1990). Are there emotion perception deficits
39 in young autistic children? *J Child Psychol Psychiatry* 31, 343-361.
- 40 Peelen, M. V., and Downing, P. E. (2005). Selectivity for the human body in the fusiform
41 gyrus. *J Neurophysiol* 93, 603-608.
- 42 Peelen, M. V., and Downing, P. E. (2007). The neural basis of visual body perception. *Nat*
43 *Rev Neurosci* 8, 636-648.
- 44 Pegna, A. J., Khateb, A., Lazeyras, F., and Seghier, M. L. (2005). Discriminating emotional
45 faces without primary visual cortices involves the right amygdala. *Nat Neurosci* 8, 24-25.
- 46 Pessoa, L. (2005). To what extent are emotional visual stimuli processed without attention
47 and awareness? *Current Opinion in Neurobiology* 15, 188-196.
- 48 Pickard, N., and Strick, P. L. (1996). Motor Areas of the Medial Wall: A Review of Their
49 Location and Functional Activation. *Cerebral Cortex* 6, 342-353.
- 50 Reed, C. L., Stone, V. E., Bozova, S., and Tanaka, J. (2003). The body-inversion effect.
51 *Psychol Sci* 14, 302-308.
- 52
53
54
55
56
57
58
59
60

- 1
2
3 Sinke, C.B., Sorger, B., Goebel, R. and de Gelder, B. Is it just a tease or a real threat? Judging
4 social interactions from bodily expressions. (submitted)
5
6 Stekelenburg, J. J., and de Gelder, B. (2004). The neural correlates of perceiving human
7 bodies: an ERP study on the body-inversion effect. *Neuroreport* 15, 777-780.
8
9 Tamietto, M., Castelli, L., Vighetti, S., Perozzo, P., Geminiani, G., Weiskrantz, L. and de
10 Gelder, B. Blindly led by emotions. Nonconscious emotional contagion for facial and bodily
11 expressions. (submitted). *Proceedings of the National Academy*.
12
13 Tamietto, M., and de Gelder, B. (2008). Affective blindsight in the intact brain: neural
14 interhemispheric summation for unseen fearful expressions. *Neuropsychologia* 46, 820-828.
15
16 Tantam, D., Monaghan, L., Nicholson, H., and Stirling, J. (1989). Autistic children's ability to
17 interpret faces: a research note. *J Child Psychol Psychiatry* 30, 623-630.
18
19 Tomkins, S.S. (1995). *Exploring affect. The Selected writings of S.S. Tomkins.* Edited by E.
20 Virginia Demos. Cambridge: Cambridge University Press.
21
22 van de Riet, W. A., Grezes, J., and de Gelder, B. (2009). Specific and common brain regions
23 involved in the perception of faces and bodies and the representation of their emotional
24 expressions. *Soc Neurosci* 4, 101-120.
25
26 Van den Stock, J., Grèzes, J., & de Gelder, B. (2008). Human and animal sounds influence
27 recognition of body language. *Brain Res* doi:10.1016/j.brainres.2008.05.040.
28
29 Van den Stock, J., Righart, R., and de Gelder, B. (2007). Body expressions influence
30 recognition of emotions in the face and voice. *Emotion*. 7, 487-494.
31
32 Wallbott (1998).
33
34 Welchew, D. E., Ashwin, C., Berkouk, K., Salvador, R., Suckling, J., Baron-Cohen, S., and
35 Bullmore, E. (2005). Functional disconnectivity of the medial temporal lobe in Asperger's
36 syndrome. *Biol Psychiatry* 57, 991-998.
37
38 Wicker, B., Keysers, C., Plailly, J., Royet, J. P., Gallese, V., and Rizzolatti, G. (2003). Both
39 of us disgusted in my insula: the common neural basis of seeing and feeling disgust. *Neuron*
40 40, 655-664.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60