

# Central Submonads and Notions of Computation

---

Titouan Carette, Louis Lemonnier, Vladimir Zamdzhiev



Laboratoire  
Méthodes  
Formelles

*Inria*

December, 20th 2022 @ SYCO 10

# Overview

- For any monoid  $M$ , its centre  $Z(M)$  is a commutative submonoid;
- For any semiring  $R$ , its centre  $Z(R)$  is a commutative subsemiring.
- For any group  $G$ , its centre  $Z(G)$  is a commutative subgroup (aka abelian subgroup);

# Overview

- For any monoid  $M$ , its centre  $Z(M)$  is a commutative submonoid;
- For any semiring  $R$ , its centre  $Z(R)$  is a commutative subsemiring.
- For any group  $G$ , its centre  $Z(G)$  is a commutative subgroup (aka abelian subgroup);
- What about monads?

# Overview

- For any monoid  $M$ , its centre  $Z(M)$  is a commutative submonoid;
- For any semiring  $R$ , its centre  $Z(R)$  is a commutative subsemiring.
- For any group  $G$ , its centre  $Z(G)$  is a commutative subgroup (aka abelian subgroup);
- What about monads?

Context:

- a symmetric monoidal category  $(\mathbf{C}, I, \otimes)$ ,
- a strong monad  $(\mathcal{T}, \eta, \mu, \tau)$ .

# Overview

- For any monoid  $M$ , its centre  $Z(M)$  is a commutative submonoid;
- For any semiring  $R$ , its centre  $Z(R)$  is a commutative subsemiring.
- For any group  $G$ , its centre  $Z(G)$  is a commutative subgroup (aka abelian subgroup);
- What about monads?

Context:

- a symmetric monoidal category  $(\mathbf{C}, I, \otimes)$ ,
- a strong monad  $(\mathcal{T}, \eta, \mu, \tau)$ .

We wonder:

- Is there a commutative submonad of  $\mathcal{T}$  which is its centre? When does it exist?
- Is there an appropriate computational interpretation?

# Background

---

# The Strength of a Monad

- Given a monoid  $M$ , its centre is defined as

$$Z(M) \stackrel{\text{def}}{=} \{x \in M \mid \forall y \in M. x \cdot y = y \cdot x\}.$$

- Notice there is an implicit *swap* in the arguments.
- *But*, the definition of a monad is independent of any monoidal structure on the base category.
- Unclear how to define a suitable notion of centre for such monads.

# The Strength of a Monad

- Given a monoid  $M$ , its centre is defined as

$$Z(M) \stackrel{\text{def}}{=} \{x \in M \mid \forall y \in M. x \cdot y = y \cdot x\}.$$

- Notice there is an implicit *swap* in the arguments.
- But*, the definition of a monad is independent of any monoidal structure on the base category.
- Unclear how to define a suitable notion of centre for such monads.
- Instead, we introduce the centre for *strong* monads acting on symmetric monoidal categories.
- The monadic strength is a natural transformation  $\tau_{X,Y}: X \otimes TY \rightarrow T(X \otimes Y)$  that satisfies some coherence conditions w.r.t. monoidal structure.
- The monadic left strength is a natural transformation  $\tau'_{X,Y}: TX \otimes Y \rightarrow T(X \otimes Y)$  that may be defined via  $\tau$  and the monoidal symmetry.

# Commutative Monads

## Definition (Commutative Monad)

A strong monad  $\mathcal{T}$  is said to be *commutative* if the following diagram:

$$\begin{array}{ccccc} \mathcal{T}X \otimes \mathcal{T}Y & \xrightarrow{\tau_{\mathcal{T}X, Y}} & \mathcal{T}(\mathcal{T}X \otimes Y) & \xrightarrow{\mathcal{T}\tau'_{X, Y}} & \mathcal{T}^2(X \otimes Y) \\ \tau'_{X, \mathcal{T}Y} \downarrow & & & & \downarrow \mu_{X \otimes Y} \\ \mathcal{T}(X \otimes \mathcal{T}Y) & \xrightarrow{\tau_{\mathcal{T}X, Y}} & \mathcal{T}^2(X \otimes Y) & \xrightarrow{\mu_{X \otimes Y}} & \mathcal{T}(X \otimes Y) \end{array}$$

commutes for every choice of objects  $X$  and  $Y$ .

# The Centre of a Monad on Set

---

## The first example

Given a monoid  $(M, e, m)$ , the writer monad:  $(M \times -) : \mathbf{Set} \rightarrow \mathbf{Set}$  has the following monad structure:

- $\eta_X : X \rightarrow M \times X :: x \mapsto (e, x)$ ;
- $\mu_X : M \times (M \times X) \rightarrow M \times X :: (z, (z', x)) \mapsto (m(z, z'), x)$ ,
- $\tau_{X,Y} : X \times (M \times Y) \rightarrow M \times (X \times Y) :: (x, (z, y)) \mapsto (z, (x, y))$ .

What should be the centre? What about  $Z(M) \times -$ ?

Indeed, it is a commutative submonad of  $(M \times -)$ .

# Commutative Monads in Set

$\mathcal{T} : \mathbf{Set} \rightarrow \mathbf{Set}$  is said to be *commutative* if the following diagram:

$$\begin{array}{ccccc} \mathcal{T}X \times \mathcal{T}Y & \xrightarrow{\tau_{\mathcal{T}X, Y}} & \mathcal{T}(\mathcal{T}X \times Y) & \xrightarrow{\mathcal{T}\tau'_{X, Y}} & \mathcal{T}^2(X \times Y) \\ \tau'_{X, \mathcal{T}Y} \downarrow & & & & \downarrow \mu_{X \times Y} \\ \mathcal{T}(X \times \mathcal{T}Y) & \xrightarrow{\tau_{\mathcal{T}X, Y}} & \mathcal{T}^2(X \times Y) & \xrightarrow{\mu_{X \times Y}} & \mathcal{T}(X \times Y) \end{array}$$

commutes for every choice of sets  $X$  and  $Y$ .

# Commutative Monads in Set

$\mathcal{T} : \mathbf{Set} \rightarrow \mathbf{Set}$  is said to be *commutative* if the following diagram:

$$\begin{array}{ccccc} \mathcal{T}X \times \mathcal{T}Y & \xrightarrow{\tau_{\mathcal{T}X, Y}} & \mathcal{T}(\mathcal{T}X \times Y) & \xrightarrow{\mathcal{T}\tau'_{X, Y}} & \mathcal{T}^2(X \times Y) \\ \tau'_{X, \mathcal{T}Y} \downarrow & & & & \downarrow \mu_{X \times Y} \\ \mathcal{T}(X \times \mathcal{T}Y) & \xrightarrow{\tau_{\mathcal{T}X, Y}} & \mathcal{T}^2(X \times Y) & \xrightarrow{\mu_{X \times Y}} & \mathcal{T}(X \times Y) \end{array}$$

commutes for every choice of sets  $X$  and  $Y$ .

How would you define a central submonad  $\mathcal{Z}$  of  $\mathcal{T}$ ?

The trick is to consider all the monadic elements of  $\mathcal{T}X$  that make the previous diagram commute.

## Definition (Centre)

Given a set  $X$ , the *centre* of  $\mathcal{T}$  at  $X$ , written  $\mathcal{Z}X$ , is defined to be the set

$$\mathcal{Z}X \stackrel{\text{def}}{=} \{t \in \mathcal{T}X \mid \forall Y \in \text{Ob}(\mathbf{Set}). \forall s \in \mathcal{T}Y. \\ \mu(\mathcal{T}\tau'(\tau(t, s))) = \mu(\mathcal{T}\tau(\tau'(t, s)))\}.$$

We write  $\iota_X : \mathcal{Z}X \subseteq \mathcal{T}X$  for the indicated subset inclusion.

- **Lemma:** The assignment  $\mathcal{Z}(-)$  extends to a functor  $\mathcal{Z} : \mathbf{Set} \rightarrow \mathbf{Set}$  when we define

$$\mathcal{Z}f \stackrel{\text{def}}{=} \mathcal{T}f|_{\mathcal{Z}X} : \mathcal{Z}X \rightarrow \mathcal{Z}Y,$$

for any function  $f: X \rightarrow Y$ .

- **Lemma:** For any two sets  $X$  and  $Y$ , the monadic unit  $\eta_X : X \rightarrow \mathcal{T}X$ , the monadic multiplication  $\mu_X : \mathcal{T}^2X \rightarrow \mathcal{T}X$ , and the monadic strength  $\tau_{X,Y} : X \times \mathcal{T}Y \rightarrow \mathcal{T}(X \times Y)$  (co)restrict respectively to functions  $\eta_X^{\mathcal{Z}} : X \rightarrow \mathcal{Z}X$ ,  $\mu_X^{\mathcal{Z}} : \mathcal{Z}^2X \rightarrow \mathcal{Z}X$  and  $\tau_{X,Y}^{\mathcal{Z}} : X \times \mathcal{Z}Y \rightarrow \mathcal{Z}(X \times Y)$ .
- **Theorem:** The assignment  $\mathcal{Z}(-)$  extends to a *commutative submonad*  $(\mathcal{Z}, \eta^{\mathcal{Z}}, \mu^{\mathcal{Z}}, \tau^{\mathcal{Z}})$  of  $\mathcal{T}$  with  $\iota_X : \mathcal{Z}X \subseteq \mathcal{T}X$  the submonad morphism. Furthermore, there exists a canonical<sup>1</sup> isomorphism  $\mathbf{Set}_{\mathcal{Z}} \cong \mathcal{Z}(\mathbf{Set}_{\mathcal{T}})$ .

---

<sup>1</sup>Details later.

- Continuation monad:  $\mathcal{T} = [[-, S], S] : \mathbf{Set} \rightarrow \mathbf{Set}$ .

- Continuation monad:  $\mathcal{T} = [[-, S], S] : \mathbf{Set} \rightarrow \mathbf{Set}$ .
  - $\mathcal{Z}X = \eta_X(X) \cong X$ ,

- Continuation monad:  $\mathcal{T} = [[-, S], S] : \mathbf{Set} \rightarrow \mathbf{Set}$ .
  - $\mathcal{Z}X = \eta_X(X) \cong X$ ,
  - The image of the monadic unit is always in the centre.

- Continuation monad:  $\mathcal{T} = [[-, S], S] : \mathbf{Set} \rightarrow \mathbf{Set}$ .
  - $\mathcal{Z}X = \eta_X(X) \cong X$ ,
  - The image of the monadic unit is always in the centre.
  - The centre is naturally isomorphic to the *identity monad*; therefore the centre is trivial.

# Examples

- Continuation monad:  $\mathcal{T} = [[-, S], S] : \mathbf{Set} \rightarrow \mathbf{Set}$ .
  - $\mathcal{Z}X = \eta_X(X) \cong X$ ,
  - The image of the monadic unit is always in the centre.
  - The centre is naturally isomorphic to the *identity monad*; therefore the centre is trivial.
- If  $\mathcal{T}$  is commutative, its centre is itself.

- Continuation monad:  $\mathcal{T} = [[-, S], S] : \mathbf{Set} \rightarrow \mathbf{Set}$ .
  - $ZX = \eta_X(X) \cong X$ ,
  - The image of the monadic unit is always in the centre.
  - The centre is naturally isomorphic to the *identity monad*; therefore the centre is trivial.
- If  $\mathcal{T}$  is commutative, its centre is itself.
- The centre of  $(M \times -)$  is indeed  $(Z(M) \times -)$ .

## Link with Lawvere theories

- In a Lawvere theory  $\mathbf{T}$ , we say that  $f: A^n \rightarrow A^{n'}$  and  $g: A^m \rightarrow A^{m'}$  commute if and only if  $f^{m'} \circ g^n$  (also written  $f \star g$ ) and  $g^{n'} \circ f^m$  (also written  $g \star f$ ) are equal, up to isomorphism.
- If  $\mathbf{S}$  is a subcategory of  $\mathbf{T}$ , the commutant of  $\mathbf{S}$  in  $\mathbf{T}$  is a subcategory of  $\mathbf{T}$  whose morphisms commute with the morphisms of  $\mathbf{S}$ . This commutant is written  $\mathbf{S}^\perp$ , and is also a Lawvere subtheory of  $\mathbf{T}$ .
- Considering this,  $\mathbf{T}^\perp$  is seen as the *centre* of the Lawvere theory  $\mathbf{T}$ .
- From  $\mathbf{T}$  arises a finitary strong monad  $\mathcal{T}$  on  $\mathbf{Set}$ , and its centre  $\mathcal{Z}$  is the monad of  $\mathbf{T}^\perp$ .

# Central Submonads in Symmetric Monoidal Categories

---

## Definition (Central Cone)

A *central cone* of  $\mathcal{T}$  at  $X$  is given by a pair  $(Z, \iota)$ , an object  $Z$  and a morphism  $\iota : Z \rightarrow \mathcal{T}X$ , such that the diagram:

$$\begin{array}{ccccc}
 Z \otimes TY & \xrightarrow{\iota \otimes TY} & \mathcal{T}X \otimes TY & \xrightarrow{\tau'_{X, TY}} & \mathcal{T}(X \otimes TY) \\
 \downarrow \iota \otimes TY & & & & \downarrow \mathcal{T}\tau_{X, Y} \\
 \mathcal{T}X \otimes TY & & & & \mathcal{T}^2(X \otimes Y) \\
 \downarrow \tau_{\mathcal{T}X, Y} & & & & \downarrow \mu_{X \otimes Y} \\
 \mathcal{T}(\mathcal{T}X \otimes Y) & \xrightarrow{\mathcal{T}\tau'_{X, Y}} & \mathcal{T}^2(X \otimes Y) & \xrightarrow{\mu_{X \otimes Y}} & \mathcal{T}(X \otimes Y)
 \end{array}$$

commutes.

## Definition (Central Submonad)

Given a strong monad  $(\mathcal{S}, \eta^{\mathcal{S}}, \mu^{\mathcal{S}}, \tau^{\mathcal{S}})$  which is a submonad of  $\mathcal{T}$  with monad monomorphism  $\iota$ , we say that  $\mathcal{S}$  is a central submonad of  $\mathcal{T}$  if for any object  $X$ ,  $(\mathcal{S}X, \iota_X)$  is a central cone for  $\mathcal{T}$  at  $X$ . Besides, this last condition implies that  $\mathcal{S}$  is commutative.

## Definition (Central Submonad)

Given a strong monad  $(\mathcal{S}, \eta^{\mathcal{S}}, \mu^{\mathcal{S}}, \tau^{\mathcal{S}})$  which is a submonad of  $\mathcal{T}$  with monad monomorphism  $\iota$ , we say that  $\mathcal{S}$  is a central submonad of  $\mathcal{T}$  if for any object  $X$ ,  $(\mathcal{S}X, \iota_X)$  is a central cone for  $\mathcal{T}$  at  $X$ . Besides, this last condition implies that  $\mathcal{S}$  is commutative.

- There always is at least one central submonad for  $\mathcal{T}$ : the identity functor is one;

## Definition (Central Submonad)

Given a strong monad  $(\mathcal{S}, \eta^{\mathcal{S}}, \mu^{\mathcal{S}}, \tau^{\mathcal{S}})$  which is a submonad of  $\mathcal{T}$  with monad monomorphism  $\iota$ , we say that  $\mathcal{S}$  is a central submonad of  $\mathcal{T}$  if for any object  $X$ ,  $(\mathcal{S}X, \iota_X)$  is a central cone for  $\mathcal{T}$  at  $X$ . Besides, this last condition implies that  $\mathcal{S}$  is commutative.

- There always is at least one central submonad for  $\mathcal{T}$ : the identity functor is one;
- They form a category with strong monad morphisms. If the category has a terminal object, the latter is the centre of  $\mathcal{T}$ .

# Centralisable Monads in Symmetric Monoidal Categories

---

## Centralisable Monad

If  $(Z, \iota)$  and  $(Z', \iota')$  are two central cones of  $\mathcal{T}$  at  $X$ , then a *morphism of central cones*  $\varphi : (Z', \iota') \rightarrow (Z, \iota)$  is a morphism  $\varphi : Z' \rightarrow Z$ , such that  $\iota \circ \varphi = \iota'$ .

## Centralisable Monad

If  $(Z, \iota)$  and  $(Z', \iota')$  are two central cones of  $\mathcal{T}$  at  $X$ , then a *morphism of central cones*  $\varphi : (Z', \iota') \rightarrow (Z, \iota)$  is a morphism  $\varphi : Z' \rightarrow Z$ , such that  $\iota \circ \varphi = \iota'$ .

A *terminal* central cone is a terminal object in the category of central cones. Its morphism component always is a monomorphism.

# Centralisable Monad

If  $(Z, \iota)$  and  $(Z', \iota')$  are two central cones of  $\mathcal{T}$  at  $X$ , then a *morphism of central cones*  $\varphi : (Z', \iota') \rightarrow (Z, \iota)$  is a morphism  $\varphi : Z' \rightarrow Z$ , such that  $\iota \circ \varphi = \iota'$ .

A *terminal* central cone is a terminal object in the category of central cones. Its morphism component always is a monomorphism.

## Definition

We say that the monad  $\mathcal{T}$  is *centralisable* if for any object  $X$ , a terminal central cone of  $\mathcal{T}$  at  $X$  exists. We write  $(\mathcal{Z}X, \iota_X)$  for the terminal central cone of  $\mathcal{T}$  at  $X$ .

# Centralisable Monad

If  $(Z, \iota)$  and  $(Z', \iota')$  are two central cones of  $\mathcal{T}$  at  $X$ , then a *morphism of central cones*  $\varphi : (Z', \iota') \rightarrow (Z, \iota)$  is a morphism  $\varphi : Z' \rightarrow Z$ , such that  $\iota \circ \varphi = \iota'$ .

A *terminal* central cone is a terminal object in the category of central cones. Its morphism component always is a monomorphism.

## Definition

We say that the monad  $\mathcal{T}$  is *centralisable* if for any object  $X$ , a terminal central cone of  $\mathcal{T}$  at  $X$  exists. We write  $(\mathcal{Z}X, \iota_X)$  for the terminal central cone of  $\mathcal{T}$  at  $X$ .

## Theorem

The assignment  $\mathcal{Z}(-)$  extends to a commutative submonad  $(\mathcal{Z}, \eta^{\mathcal{Z}}, \mu^{\mathcal{Z}}, \tau^{\mathcal{Z}})$  of  $\mathcal{T}$  with  $\iota : \mathcal{Z} \Rightarrow \mathcal{T}$  the submonad monomorphism.

Note that a submonad morphism induces a canonical embedding  $\mathcal{I} : \mathbf{C}_{\mathcal{Z}} \rightarrow \mathbf{C}_{\mathcal{T}}$ .

# **Kleisli Categories and Premonoidal Categories**

---

## Premonoidal category

- If  $\mathbf{C}$  is symmetric monoidal and  $\mathcal{T} : \mathbf{C} \rightarrow \mathbf{C}$  a strong monad;
- then  $\mathbf{C}_{\mathcal{T}}$  does not necessarily have a monoidal structure,

## Premonoidal category

- If  $\mathbf{C}$  is symmetric monoidal and  $\mathcal{T} : \mathbf{C} \rightarrow \mathbf{C}$  a strong monad;
- then  $\mathbf{C}_{\mathcal{T}}$  does not necessarily have a monoidal structure,
- $\mathbf{C}_{\mathcal{T}}$  has a *premonoidal structure* [Power and Robinson, 1997].

## Premonoidal category

- If  $\mathbf{C}$  is symmetric monoidal and  $\mathcal{T} : \mathbf{C} \rightarrow \mathbf{C}$  a strong monad;
- then  $\mathbf{C}_{\mathcal{T}}$  does not necessarily have a monoidal structure,
- $\mathbf{C}_{\mathcal{T}}$  has a *premonoidal structure* [Power and Robinson, 1997].
- there are two families of functors  $(- \otimes_l X')$  and  $(X \otimes_r -)$  on  $\mathbf{C}_{\mathcal{T}}$ .

## Premonoidal category

- If  $\mathbf{C}$  is symmetric monoidal and  $\mathcal{T} : \mathbf{C} \rightarrow \mathbf{C}$  a strong monad;
- then  $\mathbf{C}_{\mathcal{T}}$  does not necessarily have a monoidal structure,
- $\mathbf{C}_{\mathcal{T}}$  has a *premonoidal structure* [Power and Robinson, 1997].
- there are two families of functors  $(- \otimes_l X')$  and  $(X \otimes_r -)$  on  $\mathbf{C}_{\mathcal{T}}$ .

### Definition (Central morphism [Power and Robinson, 1997])

A morphism  $f: X \rightarrow Y$  in  $\mathbf{C}_{\mathcal{T}}$  is *central* if for any morphism  $f': X' \rightarrow Y'$

in  $\mathbf{C}_{\mathcal{T}}$ , the following diagram:

$$\begin{array}{ccc} X \otimes X' & \xrightarrow{f \otimes_l X'} & Y \otimes X' \\ \downarrow X \otimes_r f' & & \downarrow Y \otimes_r f \\ X \otimes Y' & \xrightarrow{f \otimes_l Y'} & Y \otimes Y' \end{array}$$

commutes in  $\mathbf{C}_{\mathcal{T}}$ .

## Premonoidal category

- If  $\mathbf{C}$  is symmetric monoidal and  $\mathcal{T} : \mathbf{C} \rightarrow \mathbf{C}$  a strong monad;
- then  $\mathbf{C}_{\mathcal{T}}$  does not necessarily have a monoidal structure,
- $\mathbf{C}_{\mathcal{T}}$  has a *premonoidal structure* [Power and Robinson, 1997].
- there are two families of functors  $(- \otimes_l X')$  and  $(X \otimes_r -)$  on  $\mathbf{C}_{\mathcal{T}}$ .

### Definition (Central morphism [Power and Robinson, 1997])

A morphism  $f : X \rightarrow Y$  in  $\mathbf{C}_{\mathcal{T}}$  is *central* if for any morphism  $f' : X' \rightarrow Y'$

in  $\mathbf{C}_{\mathcal{T}}$ , the following diagram:

$$\begin{array}{ccc} X \otimes X' & \xrightarrow{f \otimes_l X'} & Y \otimes X' \\ \downarrow X \otimes_r f' & & \downarrow Y \otimes_r f \\ X \otimes Y' & \xrightarrow{f \otimes_l Y'} & Y \otimes Y' \end{array}$$

commutes in  $\mathbf{C}_{\mathcal{T}}$ .

Central cones and central morphisms are actually equivalent notions!

- $Z(\mathbf{C}_{\mathcal{T}})$ : the wide subcategory of  $\mathbf{C}_{\mathcal{T}}$  with central morphisms.
- It is symmetric monoidal [Power and Robinson, 1997].

- $Z(\mathbf{C}_{\mathcal{T}})$ : the wide subcategory of  $\mathbf{C}_{\mathcal{T}}$  with central morphisms.
- It is symmetric monoidal [Power and Robinson, 1997].

## Proposition

*If the strong monad  $\mathcal{T}$  is centralisable, then the canonical embedding  $\mathcal{I} : \mathbf{C}_{\mathcal{Z}} \rightarrow \mathbf{C}_{\mathcal{T}}$  corestricts to an isomorphism of categories  $\hat{\mathcal{I}} : \mathbf{C}_{\mathcal{Z}} \rightarrow Z(\mathbf{C}_{\mathcal{T}})$ .*

- $Z(\mathbf{C}_{\mathcal{T}})$ : the wide subcategory of  $\mathbf{C}_{\mathcal{T}}$  with central morphisms.
- It is symmetric monoidal [Power and Robinson, 1997].

## Proposition

*If the strong monad  $\mathcal{T}$  is centralisable, then the canonical embedding  $\mathcal{I} : \mathbf{C}_{\mathcal{Z}} \rightarrow \mathbf{C}_{\mathcal{T}}$  corestricts to an isomorphism of categories  $\hat{\mathcal{I}} : \mathbf{C}_{\mathcal{Z}} \rightarrow Z(\mathbf{C}_{\mathcal{T}})$ .*

This is why we call  $\mathcal{Z}$  the central submonad of  $\mathcal{T}$ .

# Premonoidal adjunction

---

# Kleisli adjunction

- In the Kleisli adjunction between  $\mathbf{C}$  and  $\mathbf{C}_{\mathcal{T}}$ , the left adjoint,  $\mathcal{J} : \mathbf{C} \rightarrow \mathbf{C}_{\mathcal{T}}$  always corestricts to  $\hat{\mathcal{J}} : \mathbf{C} \rightarrow Z(\mathbf{C}_{\mathcal{T}})$ .

# Kleisli adjunction

- In the Kleisli adjunction between  $\mathbf{C}$  and  $\mathbf{C}_{\mathcal{T}}$ , the left adjoint,  $\mathcal{J} : \mathbf{C} \rightarrow \mathbf{C}_{\mathcal{T}}$  always corestricts to  $\hat{\mathcal{J}} : \mathbf{C} \rightarrow \mathcal{Z}(\mathbf{C}_{\mathcal{T}})$ .

## Proposition

*If the strong monad  $\mathcal{T}$  is centralisable, then  $\hat{\mathcal{J}}$  is also a left adjoint and the adjunction induces the central submonad  $\mathcal{Z}$ .*

# Characterisation

---

# The Main Theorem

## Theorem (Centralisability)

Let  $\mathbf{C}$  be a symmetric monoidal category and  $\mathcal{T}$  a strong monad on it. The following are equivalent:

1. For any object  $X$  of  $\mathbf{C}$ ,  $\mathcal{T}$  admits a terminal central cone at  $X$ ;
2. There exists a commutative submonad  $\mathcal{Z}$  of  $\mathcal{T}$  such that the canonical embedding functor  $\mathcal{I} : \mathbf{C}_{\mathcal{Z}} \rightarrow \mathbf{C}_{\mathcal{T}}$  corestricts to an isomorphism of categories  $\mathbf{C}_{\mathcal{Z}} \cong Z(\mathbf{C}_{\mathcal{T}})$ ;
3. The corestriction of the Kleisli left adjoint  $\mathcal{J} : \mathbf{C} \rightarrow \mathbf{C}_{\mathcal{T}}$  to the premonoidal centre  $\hat{\mathcal{J}} : \mathbf{C} \rightarrow Z(\mathbf{C}_{\mathcal{T}})$  also is a left adjoint.

## Some Centralisable Monads and a non Centralisable one

- Using the main theorem, it follows every strong monad on many categories of interest (e.g., **Set**, **DCPO**, **Meas**, **Top**, **Hilb**, **Vect**) is centralisable.
- If  $\mathbf{C}$  is a symmetric monoidal closed category that is total, then every strong monad on it is centralisable.
- If  $\mathcal{T}$  is a commutative monad, then  $\mathcal{T}$  is centralisable and its centre coincides with itself.

Is every strong monad centralisable?

## Some Centralisable Monads and a non Centralisable one

- Using the main theorem, it follows every strong monad on many categories of interest (e.g., **Set**, **DCPO**, **Meas**, **Top**, **Hilb**, **Vect**) is centralisable.
- If  $\mathbf{C}$  is a symmetric monoidal closed category that is total, then every strong monad on it is centralisable.
- If  $\mathcal{T}$  is a commutative monad, then  $\mathcal{T}$  is centralisable and its centre coincides with itself.

Is every strong monad centralisable? No!

Example built with a full subcategory  $\mathbf{C}$  of **Set** where not all subsets of  $\mathcal{T}X$  are objects of  $\mathbf{C}$ .

# More monads with non-trivial centres

## Example

The valuation monad  $\mathcal{V}: \mathbf{DCPO} \rightarrow \mathbf{DCPO}$  is strong, but its commutativity is an open problem [Jones, 1990]. The central submonad of  $\mathcal{V}$  is precisely the "central valuations monad" described in [Jia et al., 2021].

# Computational interpretation

---

# A meta language

Refinement of Moggi's metalanguage;

# A meta language

Refinement of Moggi's metalanguage;

$A, B ::= 1 \mid A \times B \mid A \rightarrow B \mid \mathcal{Z}A \mid \mathcal{T}A$

# A meta language

Refinement of Moggi's metalanguage;

$A, B ::= 1 \mid A \times B \mid A \rightarrow B \mid \mathcal{Z}A \mid \mathcal{T}A$

$$\frac{\Gamma, x : A \vdash M : B}{\Gamma \vdash \lambda x^A. M : A \rightarrow B} \qquad \frac{\Gamma \vdash M : A \rightarrow B \quad \Gamma \vdash N : A}{\Gamma \vdash MN : B}$$

$$\frac{\Gamma \vdash M : A}{\Gamma \vdash \text{ret}_{\mathcal{Z}} M : \mathcal{Z}A} \qquad \frac{\Gamma \vdash M : \mathcal{Z}A \quad \Gamma, x : A \vdash N : \mathcal{Z}B}{\Gamma \vdash \text{do } x \leftarrow_{\mathcal{Z}} M ; N : \mathcal{Z}B}$$

$$\frac{\Gamma \vdash M : \mathcal{Z}A}{\Gamma \vdash \iota M : \mathcal{T}A} \qquad \frac{\Gamma \vdash M : \mathcal{T}A \quad \Gamma, x : A \vdash N : \mathcal{T}B}{\Gamma \vdash \text{do } x \leftarrow_{\mathcal{T}} M ; N : \mathcal{T}B}$$

## Computational use case for the centre of a monad

**do**

x ← op1

y ← op2

f x y

**do**

y ← op2

x ← op1

f x y

If *at least one* of op1 or op2 is central, then the two programs are contextually equivalent!

## Ongoing and Future Work

---

- Completeness and internal language result for the computational interpretation;
- Notion of Commutant for monads in general;
- Link with Garner's results on commutativity.

**Thank you!**

---



Jia, X., Mislove, M. W., and Zamdzhiev, V. (2021).

**The central valuations monad (early ideas).**

In Gadducci, F. and Silva, A., editors, *9th Conference on Algebra and Coalgebra in Computer Science, CALCO 2021, August 31 to September 3, 2021, Salzburg, Austria*, volume 211 of *LIPICs*, pages 18:1–18:5. Schloss Dagstuhl - Leibniz-Zentrum für Informatik.



Jones, C. (1990).

***Probabilistic Non-determinism.***

PhD thesis, University of Edinburgh, UK.



Power, J. and Robinson, E. P. (1997).

**Premonoidal categories and notions of computation.**

*Math. Struct. Comput. Sci.*, 7:453–468.