

Newman's Lemma

Andrei Paskevich, Steffen Frerix, Peter Koepke, Adrian De Lon, Marcel Schütz

2025

Signature 1 (Term). An term is an object. Let $a, b, c, d, u, v, w, x, y, z$ denote terms.

Signature 2 (Rewriting System). A rewriting system is a notion. Let R denote a rewriting system.

Signature 3 (Reduct). A reduct of x in R is a term. Let $x \rightarrow_R y$ stand for y is a reduct of x in R .

Signature 4. $x \rightarrow_R^+ y$ is a relation.

Axiom 5. $x \rightarrow_R^+ y$ iff $x \rightarrow_R y$ or there exists a term z such that $x \rightarrow_R z \rightarrow_R^+ y$.

Axiom 6. If $x \rightarrow_R^+ y \rightarrow_R^+ z$ then $x \rightarrow_R^+ z$.

Definition 7. $x \rightarrow_R^* y$ iff $x = y$ or $x \rightarrow_R^+ y$.

Lemma 8. If $x \rightarrow_R^* y \rightarrow_R^* z$ then $x \rightarrow_R^* z$.

Definition 9 (Confluency). R is confluent iff for all a, b, c such that $a \rightarrow_R^* b, c$ there exists d such that $b, c \rightarrow_R^* d$.

Definition 10 (Local Confluency). R is locally confluent iff for all a, b, c such that $a \rightarrow_R b, c$ there exists d such that $b, c \rightarrow_R^* d$.

Definition 11 (Termination). R is terminating iff for all a, b such that $a \rightarrow_R^+ b$ we have $b \prec a$.

Definition 12 (Normal Form). A normal form of x in R is a term y such that $x \rightarrow_R^* y$ and y has no reducts in R .

Lemma 13. Let R be a terminating rewriting system. Every term x has a normal form in R .

Proof by induction. ■

Theorem 14 (Newman's Lemma). Every locally confluent terminating rewriting system is confluent.

Proof. Let R be a rewriting system. Assume R is locally confluent and terminating.

Let us demonstrate by induction that for all a, b, c such that $a \rightarrow_R^* b, c$ there exists d such that $b, c \rightarrow_R^* d$. Let a, b, c be terms.

Let us show that if $a \rightarrow_R^+ b, c$ then we have the thesis. Assume $a \rightarrow_R^+ b, c$.

Take u such that $a \rightarrow_R u \rightarrow_R^* b$. Take v such that $a \rightarrow_R v \rightarrow_R^* c$. Take w such that $u, v \rightarrow_R^* w$. Take a normal form d of w in R .

$b \rightarrow_R^* d$. Indeed take x such that $b, d \rightarrow_R^* x$. $c \rightarrow_R^* d$. Indeed take y such that $c, d \rightarrow_R^* y$. End. End. ■

Copyright

© Andrei Paskevich (2007), Steffen Frerix (2018), Peter Koepke (2018), Adrian De Lon (2021), Marcel Schütz (2025).