

Pairs & Products

$\{a, b\}$ unordered pair of a, b .


(a, b) ordered pair of a, b .

$$(a, b) = (a', b') \Leftrightarrow a = a' \& b = b'.$$

We could define $(a, b) =_{\text{def}} \{ \{a\}, \{a, b\} \}$.

The product of sets X and Y

$$X \times Y =_{\text{def}} \{ (a, b) \mid a \in X \& b \in Y \}$$



Some laws:


$$A \times (B \cup C) = (A \times B) \cup (A \times C)$$

$$A \times (B \cap C) = (A \times B) \cap (A \times C)$$

$$(A \times B) \cap (C \times D) = (A \cap C) \times (B \cap D)$$

$$(A \times B) \cup (C \times D) \subseteq (A \cup C) \times (B \cup D)$$


A binary relation between sets X, Y
is a subset $R \subseteq X \times Y$



$(x, y) \in R$
often written
 $x R y$.


E.g.

- $\{(x, y) \in \mathbb{R} \times \mathbb{R} \mid x < y\}$



- $\{(x, y) \mid x \text{ is parent of } y\} \subseteq P \times P$

P is set of people



$$f: X \rightarrow Y$$

A function from a set X to set Y
is a relation $f \subseteq X \times Y$
such that:

$$(1) (x, y) \in f \text{ & } (x, y') \in f \Rightarrow y = y'$$

for all $x \in X$, $y, y' \in Y$;

$$(2) \forall x \in X \exists y \in Y (x, y) \in f$$

Write $f(x)$ for the unique y s.t. $(x, y) \in f$.

$$f: X \rightarrow Y$$

A partial function from X to Y is
a relation $f \subseteq X \times Y$ s.t. (1).

Composing relations and functions.

$$R \subseteq X \times Y \quad S \subseteq Y \times Z$$

Their composition:

$$S \circ R =_{\text{def}} \left\{ (x, z) \in X \times Z \mid \exists y \in Y. (x, y) \in R \wedge (y, z) \in S \right\}$$

Identity:

$$\text{id}_X \subseteq X \times X$$

$$\text{id}_X =_{\text{def}} \left\{ (x, x) \mid x \in X \right\}$$

Associativity:

$$R \subseteq X \times Y, \quad S \subseteq Y \times Z, \quad T \subseteq Z \times W$$

$$T \circ (S \circ R) = (T \circ S) \circ R$$

Composition of functions / partial fns
is a function / partial function.

Special functions

Let $f : X \rightarrow Y$.

f is injective (1-1) iff injective function
= injection

$$\forall x, x' \in X. f(x) = f(x') \Rightarrow x = x'$$

f is surjective (onto) iff surjective function
= surjection

$$\forall y \in Y \exists x \in X. y = f(x).$$

f is bijective (1-1 correspondence) iff bijective fn. = bijection

f is injective and surjective.

Proposition 2.9 [P. 30]

$f : X \rightarrow Y$ is bijective iff it has an inverse

i.e. $g : Y \rightarrow X$ s.t. $g(f(x)) = x$ for all $x \in X$

and $f(g(y)) = y$ for all $y \in Y$.

Direct and inverse image

$$R \subseteq X \times Y$$

let $A \subseteq X$. Its direct image under R

$$RA = \{y \in Y \mid \exists x \in A. (x, y) \in R\}$$

let $B \subseteq Y$. Its inverse image under R

$$R^{-1}B = \{x \in X \mid \exists y \in B. (x, y) \in R\}$$

Equivalence relations.

An equivalence relation on a set X is a relation

$$R \subseteq X \times X$$

which is

reflexive: $\forall x \in X. x R x$

symmetric: $\forall x, y \in X. x R y \Rightarrow y R x$


transitive: $\forall x, y, z \in X. x R y \& y R z \Rightarrow x R z$

Let $x \in X$. Its equivalence class

$$\{x\}_R =_{\text{def}} \{y \in X \mid y R x\}$$

Theorem 2.12 [P.32]

$\{\{x\}_R \mid x \in X\}$ is a partition of the set X .




Partition:


- $x \in \{x\}^2_R$
 - $\{x\}^2_R \cap \{y\}^2_R \neq \emptyset \Rightarrow \{x\}^2_R = \{y\}^2_R$
- (1) $\{x\}^2_R \cap \{y\}^2_R \neq \emptyset \Rightarrow x R y$
- (2) $x R y \Rightarrow \{x\}^2_R = \{y\}^2_R$

Size of sets - countability.


Two sets have the same size (or cardinality) iff there is a bijection between them :



A set A is finite iff there is a bijection $f: \{m \in \mathbb{N} \mid m \leq n\} \rightarrow A$ for some $n \in \mathbb{N}$.



A set A is countable iff A is finite or there is a bijection $f: \mathbb{N} \rightarrow A$.



Lemma 2.23 Any subset A of \mathbb{N} is countable.

Proof idea :

Define $f: \mathbb{N} \rightarrow A$ by mathl. ind.

$f(1)$ is least element of A if $A \neq \emptyset$;
undefined otherwise.

$f(n+1)$ is least element of A above $f(n)$
if $f(n)$ is defined & there is
a member of A above $f(n)$;
undefined otherwise.

Corollary 2.24

A set B is countable iff there is
a bijection $g: A \rightarrow B$ where $A \subseteq \mathbb{N}$.

Lemma 2.25

A set B is countable iff there is
an injection $f: B \rightarrow A$ where
 A is countable.

In particular, a subset of a
countable set is countable.

Lemma 2.26 The set $\mathbb{N} \times \mathbb{N}$ is countable.

$$f: \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N} \quad f(m, n) = 2^m \times 3^n$$

Corollary 2.27 The set \mathbb{Q}^+ is countable

$$f: \mathbb{Q}^+ \rightarrow \mathbb{N} \times \mathbb{N} \quad f\left(\frac{m}{n}\right) = (m, n)$$

Lemma 2.28 Suppose $A_1, A_2, \dots, A_n, \dots$
are countable sets. Their union

$$A = \{x \mid \exists n \in \mathbb{N}. x \in A_n\} \text{ is countable.}$$

Theorem 2.33 \mathbb{R} is uncountable.

Proof. By contradiction.

Assume \mathbb{R} is countable.

Then $(0, 1]$ is countable.

$$f(1) = 0. \boxed{d_1^1} d_2^1 d_3^1 \dots d_i^1 \dots$$

$$f(2) = 0. d_1^2 \boxed{d_2^2} d_3^2 \dots d_i^2 \dots$$

$$f(3) = 0. d_1^3 d_2^3 \boxed{d_3^3} \dots d_i^3 \dots$$

$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$

$$f(n) = 0. d_1^n d_2^n d_3^n \dots d_i^n \dots$$

$\vdots \quad \vdots \quad \vdots \quad \vdots \quad \vdots$

$$r = 0. r_1 r_2 r_3 \dots r_i \dots$$

$$r_i = \begin{cases} 1 & \text{if } d_i^i \neq 1 \\ 2 & \text{if } d_i^i = 1. \end{cases}$$