

# ***Lecture 3***

## Constructions on Domains

## Discrete cpo's and flat domains

---

For any set  $X$ , the relation of equality

$$x \sqsubseteq x' \stackrel{\text{def}}{\iff} x = x' \quad (x, x' \in X)$$

makes  $(X, \sqsubseteq)$  into a cpo, called the **discrete** cpo with underlying set  $X$ .

Let  $X_{\perp} \stackrel{\text{def}}{=} X \cup \{\perp\}$ , where  $\perp$  is some element not in  $X$ . Then

$$d \sqsubseteq d' \stackrel{\text{def}}{\iff} (d = d') \vee (d = \perp) \quad (d, d' \in X_{\perp})$$

makes  $(X_{\perp}, \sqsubseteq)$  into a domain (with least element  $\perp$ ), called the **flat** domain determined by  $X$ .

## Binary product of cpo's and domains

---

The **product** of two cpo's  $(D_1, \sqsubseteq_1)$  and  $(D_2, \sqsubseteq_2)$  has underlying set

$$D_1 \times D_2 = \{(d_1, d_2) \mid d_1 \in D_1 \ \& \ d_2 \in D_2\}$$

and partial order  $\sqsubseteq$  defined by

$$(d_1, d_2) \sqsubseteq (d'_1, d'_2) \stackrel{\text{def}}{\iff} d_1 \sqsubseteq_1 d'_1 \ \& \ d_2 \sqsubseteq_2 d'_2 .$$

$$\frac{(x_1, x_2) \sqsubseteq (y_1, y_2)}{x_1 \sqsubseteq_1 y_1 \quad x_2 \sqsubseteq_2 y_2}$$

Lubs of chains are calculated componentwise:

$$\bigsqcup_{n \geq 0} (d_{1,n}, d_{2,n}) = \left( \bigsqcup_{i \geq 0} d_{1,i}, \bigsqcup_{j \geq 0} d_{2,j} \right) .$$

If  $(D_1, \sqsubseteq_1)$  and  $(D_2, \sqsubseteq_2)$  are domains so is  $(D_1 \times D_2, \sqsubseteq)$   
and  $\perp_{D_1 \times D_2} = (\perp_{D_1}, \perp_{D_2})$ .

## Continuous functions of two arguments

---

**Proposition.** Let  $D, E, F$  be cpo's. A function  $f : (D \times E) \rightarrow F$  is monotone if and only if it is monotone in each argument separately:

$$\forall d, d' \in D, e \in E. d \sqsubseteq d' \Rightarrow f(d, e) \sqsubseteq f(d', e)$$

$$\forall d \in D, e, e' \in E. e \sqsubseteq e' \Rightarrow f(d, e) \sqsubseteq f(d, e').$$

Moreover, it is continuous if and only if it preserves lubs of chains in each argument separately:

$$f\left(\bigsqcup_{m \geq 0} d_m, e\right) = \bigsqcup_{m \geq 0} f(d_m, e)$$

$$f\left(d, \bigsqcup_{n \geq 0} e_n\right) = \bigsqcup_{n \geq 0} f(d, e_n).$$

- A couple of derived rules:

$$\frac{x \sqsubseteq x' \quad y \sqsubseteq y'}{f(x, y) \sqsubseteq f(x', y')} \quad (f \text{ monotone})$$

---

$$f(\bigsqcup_m x_m, \bigsqcup_n y_n) = \bigsqcup_k f(x_k, y_k)$$

## Function cpo's and domains

---

Given cpo's  $(D, \sqsubseteq_D)$  and  $(E, \sqsubseteq_E)$ , the **function cpo**  $(D \rightarrow E, \sqsubseteq)$  has underlying set

$$(D \rightarrow E) \stackrel{\text{def}}{=} \{f \mid f : D \rightarrow E \text{ is a } \textit{continuous} \text{ function}\}$$

and partial order:  $f \sqsubseteq f' \stackrel{\text{def}}{\iff} \forall d \in D. f(d) \sqsubseteq_E f'(d)$ .

- A derived rule:

$$\frac{f \sqsubseteq_{(D \rightarrow E)} g \quad x \sqsubseteq_D y}{f(x) \sqsubseteq g(y)}$$

Lubs of chains are calculated 'argumentwise' (using lubs in  $E$ ):

$$\bigsqcup_{n \geq 0} f_n = \lambda d \in D. \bigsqcup_{n \geq 0} f_n(d) .$$

- A derived rule:

---

$$\left( \bigsqcup_n f_n \right) \left( \bigsqcup_m x_m \right) = \left( \bigsqcup_k f_k(x_k) \right)$$

If  $E$  is a domain, then so is  $D \rightarrow E$  and  $\perp_{D \rightarrow E}(d) = \perp_E$ , all  $d \in D$ .



## Continuity of composition

---

For cpo's  $D, E, F$ , the composition function

$$\circ : ((E \rightarrow F) \times (D \rightarrow E)) \longrightarrow (D \rightarrow F)$$

defined by setting, for all  $f \in (D \rightarrow E)$  and  $g \in (E \rightarrow F)$ ,

$$g \circ f = \lambda d \in D. g(f(d))$$

is continuous.

## Continuity of the fixpoint operator

---

Let  $D$  be a domain.

By Tarski's Fixed Point Theorem we know that each continuous function  $f \in (D \rightarrow D)$  possesses a least fixed point,  $fix(f) \in D$ .

**Proposition.** *The function*

$$fix : (D \rightarrow D) \rightarrow D$$

*is continuous.*