

# Partial Recursive Functions

# Aim

A more abstract, machine-independent description of the collection of computable partial functions than provided by register/Turing machines:

they form the smallest collection of partial functions containing some basic functions and closed under some fundamental operations for forming new functions from old—composition, **primitive recursion** and **minimization**.

The characterization is due to Kleene (1936), building on work of Gödel and Herbrand.

# Examples of recursive definitions

$$\begin{cases} f_1(0) & \equiv 0 \\ f_1(x+1) & \equiv f_1(x) + (x+1) \end{cases} \quad f_1(x) = \text{sum of } 0, 1, 2, \dots, x$$

# Examples of recursive definitions

$$\begin{cases} f_1(0) & \equiv 0 \\ f_1(x+1) & \equiv f_1(x) + (x+1) \end{cases} \quad f_1(x) = \text{sum of } 0, 1, 2, \dots, x$$

---

$$\begin{cases} f_2(0) & \equiv 0 \\ f_2(1) & \equiv 1 \\ f_2(x+2) & \equiv f_2(x) + f_2(x+1) \end{cases} \quad f_2(x) = \text{xth Fibonacci number}$$

# Examples of recursive definitions

$$\begin{cases} f_1(0) & \equiv 0 \\ f_1(x+1) & \equiv f_1(x) + (x+1) \end{cases} \quad f_1(x) = \text{sum of } 0, 1, 2, \dots, x$$

---

$$\begin{cases} f_2(0) & \equiv 0 \\ f_2(1) & \equiv 1 \\ f_2(x+2) & \equiv f_2(x) + f_2(x+1) \end{cases} \quad f_2(x) = \text{xth Fibonacci number}$$

---

$$\begin{cases} f_3(0) & \equiv 0 \\ f_3(x+1) & \equiv f_3(x+2) + 1 \end{cases} \quad f_3(x) \text{ undefined except when } x = 0$$

# Examples of recursive definitions

$$\begin{cases} f_1(0) & \equiv 0 \\ f_1(x+1) & \equiv f_1(x) + (x+1) \end{cases} \quad f_1(x) = \text{sum of } 0, 1, 2, \dots, x$$

---

$$\begin{cases} f_2(0) & \equiv 0 \\ f_2(1) & \equiv 1 \\ f_2(x+2) & \equiv f_2(x) + f_2(x+1) \end{cases} \quad f_2(x) = \text{xth Fibonacci number}$$

---

$$\begin{cases} f_3(0) & \equiv 0 \\ f_3(x+1) & \equiv f_3(x+2) + 1 \end{cases} \quad f_3(x) \text{ undefined except when } x = 0$$

---

$$f_4(x) \equiv \textit{if } x > 100 \textit{ then } x - 10 \textit{ else } f_4(f_4(x + 11)) \quad f_4 \text{ is McCarthy's "91 function", which maps } x \text{ to } 91 \text{ if } x \leq 100 \text{ and to } x - 10 \text{ otherwise}$$

# Primitive recursion

**Theorem.** Given  $f \in \mathbb{N}^n \rightarrow \mathbb{N}$  and  $g \in \mathbb{N}^{n+2} \rightarrow \mathbb{N}$ , there is a unique  $h \in \mathbb{N}^{n+1} \rightarrow \mathbb{N}$  satisfying

$$\begin{cases} h(\vec{x}, 0) & \equiv f(\vec{x}) \\ h(\vec{x}, x + 1) & \equiv g(\vec{x}, x, h(\vec{x}, x)) \end{cases}$$

for all  $\vec{x} \in \mathbb{N}^n$  and  $x \in \mathbb{N}$ .

We write  $\rho^n(f, g)$  for  $h$  and call it the partial function defined by primitive recursion from  $f$  and  $g$ .

# Primitive recursion

**Theorem.** Given  $f \in \mathbb{N}^n \rightarrow \mathbb{N}$  and  $g \in \mathbb{N}^{n+2} \rightarrow \mathbb{N}$ , there is a unique  $h \in \mathbb{N}^{n+1} \rightarrow \mathbb{N}$  satisfying

$$(*) \begin{cases} h(\vec{x}, 0) & \equiv f(\vec{x}) \\ h(\vec{x}, x+1) & \equiv g(\vec{x}, x, h(\vec{x}, x)) \end{cases}$$

for all  $\vec{x} \in \mathbb{N}^n$  and  $x \in \mathbb{N}$ .

**Proof (sketch).** *Existence:* the set

$$h \triangleq \{ (\vec{x}, x, y) \in \mathbb{N}^{n+2} \mid \exists y_0, y_1, \dots, y_x \\ f(\vec{x}) = y_0 \wedge (\bigwedge_{i=0}^{x-1} g(\vec{x}, i, y_i) = y_{i+1}) \wedge y_x = y \}$$

defines a partial function satisfying  $(*)$ .

*Uniqueness:* if  $h$  and  $h'$  both satisfy  $(*)$ , then one can prove by induction on  $x$  that  $\forall \vec{x} (h(\vec{x}, x) = h'(\vec{x}, x))$ .



# Example: addition

Addition  $add \in \mathbb{N}^2 \rightarrow \mathbb{N}$  satisfies:

$$\begin{cases} add(x_1, 0) & \equiv x_1 \\ add(x_1, x + 1) & \equiv add(x_1, x) + 1 \end{cases}$$

So  $add = \rho^1(f, g)$  where  $\begin{cases} f(x_1) & \triangleq x_1 \\ g(x_1, x_2, x_3) & \triangleq x_3 + 1 \end{cases}$

Note that  $f = \text{proj}_1^1$  and  $g = \text{succ} \circ \text{proj}_3^3$ ; so  $add$  can be built up from basic functions using composition and primitive recursion:  $add = \rho^1(\text{proj}_1^1, \text{succ} \circ \text{proj}_3^3)$ .

# Example: predecessor

Predecessor  $pred \in \mathbb{N} \rightarrow \mathbb{N}$  satisfies:

$$\begin{cases} pred(0) & \equiv 0 \\ pred(x + 1) & \equiv x \end{cases}$$

So  $pred = \rho^0(f, g)$  where  $\begin{cases} f() & \triangleq 0 \\ g(x_1, x_2) & \triangleq x_1 \end{cases}$

Thus  $pred$  can be built up from basic functions using primitive recursion:  $pred = \rho^0(\text{zero}^0, \text{proj}_1^2)$ .

# Example: multiplication

Multiplication  $\mathit{mult} \in \mathbb{N}^2 \rightarrow \mathbb{N}$  satisfies:

$$\begin{cases} \mathit{mult}(x_1, 0) & \equiv 0 \\ \mathit{mult}(x_1, x + 1) & \equiv \mathit{mult}(x_1, x) + x_1 \end{cases}$$

and thus  $\mathit{mult} = \rho^1(\mathit{zero}^1, \mathit{add} \circ (\mathit{proj}_3^3, \mathit{proj}_1^3))$ .

So  $\mathit{mult}$  can be built up from basic functions using composition and primitive recursion (since  $\mathit{add}$  can be).

**Definition.** A [partial] function  $f$  is **primitive recursive** ( $f \in \mathbf{PRIM}$ ) if it can be built up in finitely many steps from the basic functions by use of the operations of composition and primitive recursion.

In other words, the set **PRIM** of primitive recursive functions is the smallest set (with respect to subset inclusion) of partial functions containing the basic functions and closed under the operations of composition and primitive recursion.

**Definition.** A [partial] function  $f$  is primitive recursive ( $f \in \text{PRIM}$ ) if it can be built up in finitely many steps from the basic functions by use of the operations of composition and primitive recursion.

Every  $f \in \text{PRIM}$  is a total function, because:

- ▶ all the basic functions are total
- ▶ if  $f, g_1, \dots, g_n$  are total, then so is  $f \circ (g_1, \dots, g_n)$  [why?]
- ▶ if  $f$  and  $g$  are total, then so is  $\rho^n(f, g)$  [why?]


**Definition.** A [partial] function  $f$  is primitive recursive ( $f \in \text{PRIM}$ ) if it can be built up in finitely many steps from the basic functions by use of the operations of composition and primitive recursion.


**Theorem.** Every  $f \in \text{PRIM}$  is computable.

**Proof.** Already proved: basic functions are computable; composition preserves computability. So just have to show:

$\rho^n(f, g) \in \mathbb{N}^{n+1} \rightarrow \mathbb{N}$  computable if  $f \in \mathbb{N}^n \rightarrow \mathbb{N}$  and  $g \in \mathbb{N}^{n+1} \rightarrow \mathbb{N}$  are.

Suppose  $f$  and  $g$  are computed by RM programs  $F$  and  $G$  (with our usual I/O conventions). Then the RM specified on the next slide computes  $\rho^n(f, g)$ . (We assume  $X_1, \dots, X_{n+1}, C$  are some registers not mentioned in  $F$  and  $G$ ; and that the latter only use registers  $R_0, \dots, R_N$ , where  $N \geq n + 2$ .)





while  $C < X_{n+1}$  do  
 $(R_0, C) := (g(X_1, \dots, X_n, C, R_0), C+1)$



# Aim

A more abstract, machine-independent description of the collection of computable partial functions than provided by register/Turing machines:

they form the smallest collection of partial functions containing some basic functions and closed under some fundamental operations for forming new functions from old—composition, primitive recursion and **minimization**.

The characterization is due to Kleene (1936), building on work of Gödel and Herbrand.

# Minimization

Given a partial function  $f \in \mathbb{N}^{n+1} \rightarrow \mathbb{N}$ , define  $\mu^n f \in \mathbb{N}^n \rightarrow \mathbb{N}$  by

$\mu^n f(\vec{x}) \triangleq$  least  $x$  such that  $f(\vec{x}, x) = 0$   
and for each  $i = 0, \dots, x - 1$ ,  
 $f(\vec{x}, i)$  is defined and  $> 0$   
(undefined if there is no such  $x$ )

In other words

$$\mu^n f = \{(\vec{x}, x) \in \mathbb{N}^{n+1} \mid \exists y_0, \dots, y_x \\ \left( \bigwedge_{i=0}^x f(\vec{x}, i) = y_i \right) \wedge \left( \bigwedge_{i=0}^{x-1} y_i > 0 \right) \wedge y_x = 0\}$$

# Example of minimization

integer part of  $x_1/x_2$   $\equiv$  least  $x_3$  such that  
(undefined if  $x_2=0$ )  $x_1 < x_2(x_3 + 1)$

# Example of minimization

integer part of  $x_1/x_2$   $\equiv$  least  $x_3$  such that  
(undefined if  $x_2=0$ )  $x_1 < x_2(x_3 + 1)$

$$\equiv \mu^2 f(x_1, x_2)$$

where  $f \in \mathbb{N}^3 \rightarrow \mathbb{N}$  is

$$f(x_1, x_2, x_3) \triangleq \begin{cases} 1 & \text{if } x_1 \geq x_2(x_3 + 1) \\ 0 & \text{if } x_1 < x_2(x_3 + 1) \end{cases}$$

**Definition.** A partial function  $f$  is **partial recursive** ( $f \in \mathbf{PR}$ ) if it can be built up in finitely many steps from the basic functions by use of the operations of composition, primitive recursion and minimization.

In other words, the set **PR** of partial recursive functions is the smallest set (with respect to subset inclusion) of partial functions containing the basic functions and closed under the operations of composition, primitive recursion and minimization.


**Definition.** A partial function  $f$  is **partial recursive** ( $f \in \mathbf{PR}$ ) if it can be built up in finitely many steps from the basic functions by use of the operations of composition, primitive recursion and minimization.

**Theorem.** Every  $f \in \mathbf{PR}$  is computable.

**Proof.** Just have to show:

$\mu^n f \in \mathbb{N}^n \rightarrow \mathbb{N}$  is computable if  $f \in \mathbb{N}^{n+1} \rightarrow \mathbb{N}$  is.

Suppose  $f$  is computed by RM program  $F$  (with our usual I/O conventions). Then the RM specified on the next slide computes  $\mu^n f$ . (We assume  $X_1, \dots, X_n, C$  are some registers not mentioned in  $F$ ; and that the latter only uses registers  $R_0, \dots, R_N$ , where  $N \geq n + 1$ .)



START

$(X_1, \dots, X_n) ::= (R_1, \dots, R_n)$

$(R_1, \dots, R_n, R_{n+1}) ::= (X_1, \dots, X_n, C)$

$(R_0, R_{n+2}, \dots, R_N) ::= (0, 0, \dots, 0)$

$F$

$R_0^-$

$R_0 := f(X_1, \dots, X_n, C);$   
while  $R_0 > 0$  do  
   $C := C + 1;$   
   $R_0 := f(X_1, \dots, X_n, C)$

$R_0 ::= C$

HALT