

Quantum Computing: Exercise Sheet 1

Anuj Dawar

September 26, 2018

Lecture 1: Bits and Qubits

1. Which of the following are possible states of a qubit?

- (a) $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$
- (b) $\frac{\sqrt{3}}{2}|1\rangle - \frac{1}{2}|0\rangle$
- (c) $0.7|0\rangle + 0.3|1\rangle$
- (d) $0.8|0\rangle + 0.6|1\rangle$
- (e) $\cos \theta|0\rangle + i \sin \theta|1\rangle$
- (f) $\cos^2 \theta|0\rangle - \sin^2 \theta|1\rangle$
- (g) $\frac{1}{2}|0\rangle + \frac{i}{2}|0\rangle - \frac{i}{2}|1\rangle + \frac{1}{2}|1\rangle$

For each valid state among the above, give the probabilities of observing $|0\rangle$ and $|1\rangle$ when the system is measured in the standard computational basis.

What are the probabilities of the two outcomes when the state is measured in the basis $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$, $\frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$?

2. A two-qubit system is in the following state

$$\frac{1}{\sqrt{30}}(|00\rangle + 2i|01\rangle - 3|10\rangle - 4i|11\rangle)$$

The first qubit is measured and observed to be 1. What is the state of the system after the measurement? What is the probability that a subsequent measurement of the second qubit will observe a 1?

Lecture 2: Linear Algebra

- 3. Show that the operations of addition and scalar multiplication defined on slide 20 satisfy the axioms of a vector space given on slide 19.
- 4. **Uniqueness of dimension.** This exercise asks you to prove the assertion on slide 21, namely that the size of the basis is uniquely determined by the vector space V . Suppose that $|v_1\rangle, \dots, |v_n\rangle$ is a basis for V . Let $|u_1\rangle, \dots, |u_{n+1}\rangle$ be any collection of $n + 1$ vectors. Show that they cannot all be linearly independent, i.e. one of them must be expressible as a linear combination of the others.

5. For each of the two example linear operators on slide 25, express them in the basis $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle), \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$.
6. Verify that the inner product on \mathbb{C}^n defined on slide 27 satisfies the axioms on slide 26.
7. Express the two linear operators on slide 25 as a linear combination of outer products.
8. Find the eigenvalues and associated eigenvectors of the two linear operators on slide 25.
9. Show that the three Pauli matrices (slides 46 and 47) are unitary. Do the same for the Hadamard matrix (slide 63).
10. If I is the 2-dimensional identity matrix and H is the Hadamard operator, give matrix representations of the operators $I \otimes H$ and $H \otimes I$. Work out, by multiplying these matrices with the corresponding vectors the eight vectors

$$(I \otimes H)|00\rangle, (I \otimes H)|01\rangle, (I \otimes H)|10\rangle, (I \otimes H)|11\rangle, \text{ and}$$

$$(H \otimes I)|00\rangle, (H \otimes I)|01\rangle, (H \otimes I)|10\rangle, (H \otimes I)|11\rangle$$

Lecture 3: Quantum Mechanics

11. (The exercise on slide 44) Prove that unitary operations are norm preserving.
12. For the Pauli matrices X, Y and Z , prove that $XY = iZ$. Show that $X^2 = Y^2 = Z^2 = I$.
13. (The exercise on slide 50). Show that the measurement operators P_0 and P_1 give probabilities $|\alpha|^2$ and $|\beta|^2$ according to the third postulate. Also verify that P_0 and P_1 satisfy the completeness equation on slide 49.
14. Suppose a two-qubit system is in the state $0.8|00\rangle + 0.6|11\rangle$. A Pauli X gate (i.e. a NOT gate) is applied to the second qubit and a measurement performed in the computational basis. What are the probabilities of the possible measurement outcomes?
15. For the two entangled states given on slide 55, prove that they cannot be expressed as the tensor product of single qubit states.
16. The measurement postulate states that the probability of measuring outcome m is $\langle \psi | P_m^\dagger P_m | \psi \rangle$. Prove that when P_m is $|i\rangle\langle i|$, i.e. a measurement in the computational basis, this gives $p(i) = |\langle i | \psi \rangle|^2$.

Lecture 4: Models of Quantum Computation

17. The exercise on slide 62 asks you to describe a unitary operation that implements the Boolean *And* operation. In fact, the unitary operation is not uniquely determined by the description given on that slide. Describe *two* different unitary operations that satisfy the definition of the Boolean *And* in the sense of slide 62.
18. Describe, in matrix form, the action of the controlled- H gate, where H is the Hadamard gate.
19. Show how the controlled-Not gate can be constructed from Hadamard gates and the controlled- Z . Demonstrate that the construction is correct by multiplying the corresponding matrices.
20. Write the Toffoli gate in matrix form.
21. The circuit on slide 69 is formed as the composition of 13 unitary operations, each of them obtained as the tensor product of one and two-qubit operations. Write each of these operators in matrix form and verify that their product is the same as your answer to Question 20.
22. **No-cloning theorem.** The no-cloning theorem states that a quantum state $|\psi\rangle$ cannot be duplicated. That is, there is no quantum operation that will map an arbitrary state $|\psi0\rangle$ to $|\psi\psi\rangle$. Show this by considering any unitary map U that maps $|00\rangle$ to $|00\rangle$ and $|10\rangle$ to $|11\rangle$ and exhibiting a state $|\psi\rangle$ for which $U|\psi0\rangle \neq |\psi\psi\rangle$.
23. **Deutsch-Josza Algorithm.** This exercise is aimed at proving the correctness of the circuit described on slide 72. Consider the operator U_f on slide 71 and show that:
 - (a) if $|y\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$, then $U_f|xy\rangle = (-1)^{f(x)}|xy\rangle$;
 - (b) if, in addition, $|x\rangle = H|0\rangle$, then $U_f|xy\rangle = ((-1)^{f(0)}|0\rangle + (-1)^{f(1)}|1\rangle)|y\rangle$;
 - (c) the final output of the circuit is $(H \otimes I)U_f|xy\rangle$ where $|x\rangle$ and $|y\rangle$ are as above. Show that this is (up to global phase and normalisation) $|0y\rangle$ if f is constant and $|1y\rangle$ if f is balanced.