

Lecture 5

L108 assessment—heads up

Graded exercise sheet
(25% credit)

Available (via Moodle)

Friday 30 October 2020

Solutions due (via Moodle)

Friday 6 Nov 2020 4pm

Exponentials

Given $X, Y \in \mathbf{Set}$, let $Y^X \in \mathbf{Set}$ denote the set of all functions from X to Y .

$$Y^X = \mathbf{Set}(X, Y) = \{f \subseteq X \times Y \mid f \text{ is single-valued and total}\}$$

Aim to characterise Y^X category theoretically.

Exponentials

Given $X, Y \in \mathbf{Set}$, let $Y^X \in \mathbf{Set}$ denote the set of all functions from X to Y .

Aim to characterise Y^X category theoretically.

Function application gives a morphism
 $\mathbf{app} : Y^X \times X \rightarrow Y$ in \mathbf{Set} .

$$\mathbf{app}(f, x) = f x \quad (f \in Y^X, x \in X)$$

so as a set of ordered pairs, \mathbf{app} is
 $\{((f, x), y) \in (Y^X \times X) \times Y \mid (x, y) \in f\}$

Exponentials

Given $X, Y \in \mathbf{Set}$, let $Y^X \in \mathbf{Set}$ denote the set of all functions from X to Y .

Aim to characterise Y^X category theoretically.

Function application gives a morphism
 $\text{app} : Y^X \times X \rightarrow Y$ in \mathbf{Set} .

Currying operation transforms morphisms
 $f : Z \times X \rightarrow Y$ in \mathbf{Set} to morphisms $\text{cur } f : Z \rightarrow Y^X$

$$\text{cur } f \ z \ x = f(z, x) \quad (f \in Y^X, z \in Z, x \in X)$$

$$\begin{aligned} \text{cur } f \ z &= \{(x, y) \mid ((z, x), y) \in f\} \\ \text{cur } f &= \{(z, g) \mid g = \{(x, y) \mid ((z, x), y) \in f\}\} \end{aligned}$$

Haskell Curry

Haskell Brooks Curry (/ˈhæskəl/; September 12, 1900 – September 1, 1982) was an [American mathematician](#) and [logician](#). Curry is best known for his work in [combinatory logic](#); while the initial concept of combinatory logic was based on a single paper by [Moses Schönfinkel](#),^[1] much of the development was done by Curry. Curry is also known for [Curry's paradox](#) and the [Curry–Howard correspondence](#). There are three programming languages named after him, [Haskell](#), [Brook](#) and [Curry](#), as well as the concept of *currying*, a

Haskell Brooks Curry



Born

September 12, 1900
[Millis, Massachusetts](#)

Died

September 1, 1982
(aged 81)
[State College, Pennsylvania](#)

Nationality

American

Alma mater

[Harvard University](#)

Known for

[Combinatory logic](#)
[Curry–Howard correspondence](#)

For each function $f : Z \times X \rightarrow Y$ we get a commutative diagram in **Set**:

$$\begin{array}{ccc}
 Y^X \times X & \xrightarrow{\text{app}} & Y \\
 \text{cur } f \times \text{id}_X \uparrow & & \nearrow f \\
 Z \times X & & \\
 \\
 (\text{cur } f \ z, x) & \longmapsto & \text{cur } f \ z \ x = f(z, x) \\
 \uparrow & & \nearrow \\
 (z, x) & &
 \end{array}$$

For each function $f : Z \times X \rightarrow Y$ we get a commutative diagram in **Set**:

$$\begin{array}{ccc}
 Y^X \times X & \xrightarrow{\text{app}} & Y \\
 \text{cur } f \times \text{id}_X \uparrow & & \nearrow f \\
 Z \times X & &
 \end{array}$$

Furthermore, if any function $g : Z \rightarrow Y^X$ also satisfies

$$\begin{array}{ccc}
 Y^X \times X & \xrightarrow{\text{app}} & Y \\
 g \times \text{id}_X \uparrow & & \nearrow f \\
 Z \times X & &
 \end{array}$$

then $g = \text{cur } f$, because of **function extensionality**...

Function Extensionality

Two functions $f, g \in Y^X$ are equal if (and only if)
 $\forall x \in X, f x = g x$.

This is true of the set-theoretic notion of function, because then

$$\begin{aligned} & \{(x, f x) \mid x \in X\} = \{(x, g x) \mid x \in X\} \\ \text{i.e.} & \{(x, y) \mid (x, y) \in f\} = \{(x, y) \mid (x, y) \in g\} \\ \text{i.e.} & f = g \end{aligned}$$

(in other words it reduces to the extensionality property of sets: two sets are equal iff they have the same elements).

Exponential objects

Suppose a category \mathbf{C} has binary products, that is, for every pair of \mathbf{C} -objects X and Y there is a product diagram $X \xleftarrow{\pi_1} X \times Y \xrightarrow{\pi_2} Y$.

Notation: given $f \in \mathbf{C}(X, X')$ and $f' \in \mathbf{C}(Y, Y')$, then $f \times f' : X \times Y \rightarrow X' \times Y'$ stands for $\langle f \circ \pi_1, f' \circ \pi_2 \rangle$, that is, the unique morphism $g \in \mathbf{C}(X \times Y, X' \times Y')$ satisfying $\pi_1 \circ g = f \circ \pi_1$ and $\pi_2 \circ g = f' \circ \pi_2$.

Exponential objects

Suppose a category \mathbf{C} has binary products.

An **exponential** for \mathbf{C} -objects X and Y is specified by

object Y^X + morphism $\text{app} : Y^X \times X \rightarrow Y$

satisfying the universal property

for all $Z \in \mathbf{C}$ and $f \in \mathbf{C}(Z \times X, Y)$, there is a unique

$g \in \mathbf{C}(Z, Y^X)$ such that

$$\begin{array}{ccc} Y^X \times X & \xrightarrow{\text{app}} & Y \\ g \times \text{id}_X \uparrow & & \nearrow f \\ Z \times X & & \end{array}$$

commutes in \mathbf{C} .

Notation: we write $\boxed{\text{cur } f}$ for the unique g such that $\text{app} \circ (g \times \text{id}_X) = f$.

Exponential objects

The universal property of $\text{app} : Y^X \times X \rightarrow Y$ says that there is a bijection

$$\mathbf{C}(Z, Y^X) \cong \mathbf{C}(Z \times X, Y)$$

$$g \mapsto \text{app} \circ (g \times \text{id}_X)$$

$$\text{cur } f \leftarrow f$$

$$\text{app} \circ (\text{cur } f \times \text{id}_X) = f$$

$$g = \text{cur}(\text{app} \circ (g \times \text{id}_X))$$

Exponential objects

The universal property of $\text{app} : Y^X \times X \rightarrow Y$ says that there is a bijection...

It also says that (Y^X, app) is a terminal object in the following category:

- ▶ objects: (Z, f) where $f \in \mathbf{C}(Z \times X, Y)$
- ▶ morphisms $g : (Z, f) \rightarrow (Z', f')$ are $g \in \mathbf{C}(Z, Z')$ such that $f' \circ (g \times \text{id}_X) = f$
- ▶ composition and identities as in \mathbf{C} .

So when they exist, exponential objects are unique up to (unique) isomorphism.

Cartesian closed category

Definition. \mathcal{C} is a **cartesian closed category (ccc)** if it is a category with a terminal object, binary products and exponentials of any pair of objects.

This is a key concept for the semantics of lambda calculus and for the foundations of functional programming languages.

Notation: an exponential object Y^X is often written as $X \rightarrow Y$

Cartesian closed category

Definition. \mathbf{C} is a **cartesian closed category (ccc)** if it is a category with a terminal object, binary products and exponentials of any pair of objects.

Examples:

- ▶ **Set** is a ccc — as we have seen.
- ▶ **Preord** is a ccc: we already saw that it has a terminal object and binary products; the exponential of (P_1, \sqsubseteq_1) and (P_2, \sqsubseteq_2) is $(P_1 \rightarrow P_2, \sqsubseteq)$ where

$$P_1 \rightarrow P_2 = \mathbf{Preord}((P_1, \sqsubseteq_1), (P_2, \sqsubseteq_2))$$

$$f \sqsubseteq g \Leftrightarrow \forall x \in P_1, f x \sqsubseteq_2 g x$$

(check that this is a pre-order and does give an exponential in **Preord**)