

Topic 3

Constructions on Domains

Discrete cpo's and flat domains

For any set X , the relation of equality

$$x \sqsubseteq x' \stackrel{\text{def}}{\iff} x = x' \quad (x, x' \in X)$$

makes (X, \sqsubseteq) into a cpo, called the **discrete** cpo with underlying set X .

Discrete cpo's and flat domains

For any set X , the relation of equality

$$x \sqsubseteq x' \stackrel{\text{def}}{\iff} x = x' \quad (x, x' \in X)$$

makes (X, \sqsubseteq) into a cpo, called the **discrete** cpo with underlying set X .

Let $X_{\perp} \stackrel{\text{def}}{=} X \cup \{\perp\}$, where \perp is some element not in X . Then

$$d \sqsubseteq d' \stackrel{\text{def}}{\iff} (d = d') \vee (d = \perp) \quad (d, d' \in X_{\perp})$$

makes (X_{\perp}, \sqsubseteq) into a domain (with least element \perp), called the **flat** domain determined by X .

Binary product of cpo's and domains

The **product** of two cpo's (D_1, \sqsubseteq_1) and (D_2, \sqsubseteq_2) has underlying set

$$D_1 \times D_2 = \{(d_1, d_2) \mid d_1 \in D_1 \ \& \ d_2 \in D_2\}$$

and partial order \sqsubseteq defined by

$$(d_1, d_2) \sqsubseteq (d'_1, d'_2) \stackrel{\text{def}}{\iff} d_1 \sqsubseteq_1 d'_1 \ \& \ d_2 \sqsubseteq_2 d'_2 .$$

$$\frac{(x_1, x_2) \sqsubseteq (y_1, y_2)}{x_1 \sqsubseteq_1 y_1 \quad x_2 \sqsubseteq_2 y_2}$$

Lubs of chains are calculated componentwise:

$$\bigsqcup_{n \geq 0} (d_{1,n}, d_{2,n}) = \left(\bigsqcup_{i \geq 0} d_{1,i}, \bigsqcup_{j \geq 0} d_{2,j} \right) .$$

If (D_1, \sqsubseteq_1) and (D_2, \sqsubseteq_2) are domains so is $(D_1 \times D_2, \sqsubseteq)$
and $\perp_{D_1 \times D_2} = (\perp_{D_1}, \perp_{D_2})$.

Continuous functions of two arguments

Proposition. Let D, E, F be cpo's. A function $f : (D \times E) \rightarrow F$ is monotone if and only if it is monotone in each argument separately:

$$\forall d, d' \in D, e \in E. d \sqsubseteq d' \Rightarrow f(d, e) \sqsubseteq f(d', e)$$

$$\forall d \in D, e, e' \in E. e \sqsubseteq e' \Rightarrow f(d, e) \sqsubseteq f(d, e').$$

Moreover, it is continuous if and only if it preserves lubs of chains in each argument separately:

$$f\left(\bigsqcup_{m \geq 0} d_m, e\right) = \bigsqcup_{m \geq 0} f(d_m, e)$$

$$f\left(d, \bigsqcup_{n \geq 0} e_n\right) = \bigsqcup_{n \geq 0} f(d, e_n).$$

- A couple of derived rules:

$$\frac{x \sqsubseteq x' \quad y \sqsubseteq y'}{f(x, y) \sqsubseteq f(x', y')} \quad (f \text{ monotone})$$

$$f(\bigsqcup_m x_m, \bigsqcup_n y_n) = \bigsqcup_k f(x_k, y_k)$$

Function cpo's and domains

Given cpo's (D, \sqsubseteq_D) and (E, \sqsubseteq_E) , the **function cpo** $(D \rightarrow E, \sqsubseteq)$ has underlying set

$$(D \rightarrow E) \stackrel{\text{def}}{=} \{f \mid f : D \rightarrow E \text{ is a } \textit{continuous} \text{ function}\}$$

and partial order: $f \sqsubseteq f' \stackrel{\text{def}}{\iff} \forall d \in D . f(d) \sqsubseteq_E f'(d)$.

Function cpo's and domains

Given cpo's (D, \sqsubseteq_D) and (E, \sqsubseteq_E) , the **function cpo** $(D \rightarrow E, \sqsubseteq)$ has underlying set

$$(D \rightarrow E) \stackrel{\text{def}}{=} \{f \mid f : D \rightarrow E \text{ is a } \textit{continuous} \text{ function}\}$$

and partial order: $f \sqsubseteq f' \stackrel{\text{def}}{\iff} \forall d \in D . f(d) \sqsubseteq_E f'(d)$.

- A derived rule:

$$\frac{f \sqsubseteq_{(D \rightarrow E)} g \quad x \sqsubseteq_D y}{f(x) \sqsubseteq g(y)}$$

Lubs of chains are calculated 'argumentwise' (using lubs in E):

$$\bigsqcup_{n \geq 0} f_n = \lambda d \in D. \bigsqcup_{n \geq 0} f_n(d) .$$

If E is a domain, then so is $D \rightarrow E$ and $\perp_{D \rightarrow E}(d) = \perp_E$, all $d \in D$.

Lubs of chains are calculated 'argumentwise' (using lubs in E):

$$\bigsqcup_{n \geq 0} f_n = \lambda d \in D. \bigsqcup_{n \geq 0} f_n(d) .$$

- A derived rule:

$$\left(\bigsqcup_n f_n \right) \left(\bigsqcup_m x_m \right) = \bigsqcup_k f_k(x_k)$$

If E is a domain, then so is $D \rightarrow E$ and $\perp_{D \rightarrow E}(d) = \perp_E$, all $d \in D$.

Continuity of composition

For cpo's D, E, F , the composition function

$$\circ : ((E \rightarrow F) \times (D \rightarrow E)) \longrightarrow (D \rightarrow F)$$

defined by setting, for all $f \in (D \rightarrow E)$ and $g \in (E \rightarrow F)$,

$$g \circ f = \lambda d \in D. g(f(d))$$

is continuous.

Continuity of the fixpoint operator

Let D be a domain.

By Tarski's Fixed Point Theorem we know that each continuous function $f \in (D \rightarrow D)$ possesses a least fixed point, $fix(f) \in D$.

Proposition. *The function*

$$fix : (D \rightarrow D) \rightarrow D$$

is continuous.