

# Introduction to Probability

Lecture 3: Expectation properties, variance, discrete distributions

Mateja Jamnik, Thomas Sauerwald

University of Cambridge, Department of Computer Science and Technology  
email: {mateja.jamnik,thomas.sauerwald}@cl.cam.ac.uk



UNIVERSITY OF  
CAMBRIDGE

Properties of expectation

Variance

Bernoulli discrete random variable

Binomial discrete random variable



## Properties of expectation: linearity

### Linearity of expectation

Expectations preserve linearity: if  $a$  and  $b$  are constants, then

$$\mathbf{E}[aX + b] = a\mathbf{E}[X] + b$$

**Proof:**

### Example

Let the event be a roll of a 6-sided die,  $X$  its random variable, and  $Y$  another random variable where  $Y = 3X + 1$ . What are the expected values  $\mathbf{E}[X]$  and  $\mathbf{E}[Y]$ ?

Answer



## Properties of expectation: additivity

Additivity of expectation

Expectation of a sum is equal to the sum of expectations: if  $X$  and  $Y$  are any random variables on the same sample space then

$$\mathbf{E}[X + Y] = \mathbf{E}[X] + \mathbf{E}[Y]$$

Example

Let the events be rolls of 2 dice, and  $X$  the random variable for the roll of die 1, and  $Y$  for the roll of die 2. What is the expected value of the sum of the rolls of the two dice?

Answer



## Properties of expectation: LOTUS

Law of the unconscious statistician (LOTUS)

Let  $X$  be a random variable, and  $Y$  another random variable that is a function of  $X$ , so  $Y = g(X)$ . Let  $p(x)$  be a PMF of  $X$ . Then

$$\mathbf{E}[Y] = \mathbf{E}[g(X)] = \sum_{x:p(x)>0} g(x)p(x)$$

Note how now we no longer need to know PMF of  $Y$ .

- LOTUS is also known as **expected value of a function of a random variable**.
- Note that the properties of expectation let you avoid defining difficult PMFs.
- Let  $X$  be a discrete RV, then:
  - $\mathbf{E}[X^2]$  is know as the **second moment of  $X$** .
  - $\mathbf{E}[X^n]$  is know as the  **$n^{\text{th}}$  moment of  $X$** .



## Second moment example

### Example

Let  $X$  be a discrete random variable that ranges over the values  $\{-1, 0, 1\}$ , and respective probabilities  $\mathbf{P}[X = -1] = 0.2$ ,  $\mathbf{P}[X = 0] = 0.5$  and  $\mathbf{P}[X = 1] = 0.3$ . Let another random variable  $Y = X^2$  (second moment). What is  $\mathbf{E}[Y]$ ?

Answer

Note that  $Y = g(X) = X^2$  and  $\mathbf{E}[Y] = \mathbf{E}[g(X)] = \sum_{x:p(x)>0} g(x)p(x)$ , thus



# Outline

---

Properties of expectation

Variance




Bernoulli discrete random variable

Binomial discrete random variable



## Spread in the distribution

Expectation is a useful statistic, but it does not give a detailed view of the PMF. Consider these 3 distributions (PMFs).



- Expectation is the same for all distributions:  $\mathbf{E}[X] = 3$ .
- First has the greatest spread, the third has the least spread.
- But the "spread" or "dispersion" of  $X$  in the distribution is very different!
- **Variance**,  $\mathbf{V}[X]$  defines a formal quantification of "spread".
- Several ways to quantify: it uses average square distance from the mean.





## Definition of variance

### Variance

The variance of a discrete random variable  $X$  with expected value (mean)  $\mu$  is:

$$\mathbf{V}[X] = \mathbf{E}[(X - \mu)^2]$$

When computing the variance, we often use a different form of the same equation:

$$\mathbf{V}[X] = \mathbf{E}[X^2] - (\mathbf{E}[X])^2$$

Proof:

Note:

- $\mathbf{V}[X] \geq 0$
- AKA: Second **central** moment, or square of the standard deviation



## Example with a die roll

### Example

Let  $X$  be the value on one roll of a 6-sided fair die. Recall that  $\mathbf{E}[X] = \frac{7}{2} = 3.5$ . What is  $\mathbf{V}[X]$ ?

Answer

Using  $\mathbf{V}[X] = \mathbf{E}[X^2] - (\mathbf{E}[X])^2$ :

Using  $\mathbf{V}[X] = \mathbf{E}[(X - \mu)^2] = \mathbf{E}[(X - \mathbf{E}[X])^2]$ :



## Example of spread

### Example

Let  $X$ ,  $Y$  and  $Z$  be discrete random variables with the range  $X : \{10\}$  and probability 1, and  $Y : \{11, 9\}$  and  $Z : \{110, -90\}$  with equal probabilities  $\frac{1}{2}$ . Compute expectation and variance for  $X$ ,  $Y$  and  $Z$ .

a)  $\mathbf{E}[X] = \sum_x xp(x) = 10 \cdot 1 = \mathbf{10}$


$$\begin{aligned}\mathbf{V}[X] &= \mathbf{E}[(X - \mathbf{E}[X])^2] = \mathbf{E}[(X - 10)^2] \\ &= (X - 10)^2 p(x) = 0^2 \cdot 1 = \mathbf{0}\end{aligned}$$

b)  $\mathbf{E}[Y] = (11)(0.5) + (9)(0.5) = \mathbf{10}$

$$\begin{aligned}\mathbf{V}[Y] &= \mathbf{E}[(Y - \mathbf{E}[Y])^2] = \mathbf{E}[(Y - 10)^2] \\ &= (11 - 10)^2(0.5) + (9 - 10)^2(0.5) = \mathbf{1}\end{aligned}$$

c)  $\mathbf{E}[Z] = (110)(0.5) + (-90)(0.5) = \mathbf{10}$

$$\begin{aligned}\mathbf{V}[Z] &= \mathbf{E}[(Z - \mathbf{E}[Z])^2] = \mathbf{E}[(Z - 10)^2] = \\ &= (110 - 10)^2(0.5) + (-90 - 10)^2(0.5) \\ &= \mathbf{100^2} = \mathbf{10000}\end{aligned}$$



## Standard deviation

---

- Standard deviation is a kind of average distance of a sample of the mean, i.e., a root mean square (RMS) average.
- Variance is the square of this average distance.

Standard deviation

Standard deviation is defined as a square root of variance:

$$\mathbf{SD} [ X ] = \sqrt{\mathbf{V} [ X ]}$$

Note:

- $\mathbf{E} [ X ]$  and  $\mathbf{V} [ X ]$  are real numbers, not RVs.
- $\mathbf{V} [ X ]$  is expressed in units of the values in the range of  $X^2$ .
- $\mathbf{SD} [ X ]$  is expressed in units of the values in the range of  $X$ .
- For the spread example above:  $\mathbf{SD} [ X ] = 0$ ,  $\mathbf{SD} [ Y ] = 1$ ,  $\mathbf{SD} [ Z ] = 100$ .



- **Property 1:**  $\mathbf{V}[X] = \mathbf{E}[X^2] - (\mathbf{E}[X])^2$
- **Property 2:** variance is **not linear**:  $\mathbf{V}[aX + b] = a^2\mathbf{V}[X]$

Proof:

$$\begin{aligned}\mathbf{V}[aX + b] &= \mathbf{E}[(aX + b)^2] - (\mathbf{E}[aX + b])^2 \\ &= \mathbf{E}[a^2X^2 + 2abX + b^2] - (a\mathbf{E}[X] + b)^2 \\ &= a^2\mathbf{E}[X^2] + 2ab\mathbf{E}[X] + b^2 - (a^2(\mathbf{E}[X])^2 + 2ab\mathbf{E}[X] + b^2) \\ &= a^2\mathbf{E}[X^2] - (a^2(\mathbf{E}[X])^2) = a^2(\mathbf{E}[X^2] - (\mathbf{E}[X])^2) \\ &= a^2\mathbf{V}[X]\end{aligned}$$

## Summary of expectation and variance for discrete RV

---

$$\mathbf{E}[X] = \sum_{x:\mathbf{P}[x]>0} x\mathbf{P}[x] = \sum_x xp(x)$$

### Properties of Expectation

$$\mathbf{E}[X + Y] = \mathbf{E}[X] + \mathbf{E}[Y]$$

$$\mathbf{E}[aX + b] = a\mathbf{E}[X] + b$$

$$\mathbf{E}[g(X)] = \sum_x g(x)p_X(x)$$

### Properties of Variance

$$\mathbf{V}[X] = \mathbf{E}[(X - \mu)^2]$$

$$\mathbf{V}[X] = \mathbf{E}[X^2] - (\mathbf{E}[X])^2$$

$$\mathbf{V}[aX + b] = a^2\mathbf{V}[X]$$



- There is deluge of classic RV abstractions that show up in problems.
- They give rise to significant discrete distributions.
- If problem fits, use precalculated (parametric) PMF, expectation, variance and other properties by providing parameters of the problem.
- We will cover the following RVs:
  1. Bernoulli
  2. Binomial
  3. Poisson
  4. Geometric
  5. Negative Binomial
  6. Hypergeometric



# Outline

---

Properties of expectation

Variance

**Bernoulli discrete random variable**

Binomial discrete random variable





### Bernoulli discrete random variable

A Bernoulli RV  $X$  maps "success" of an experiment to 1 and "failure" to 0. It is AKA indicator RV, boolean RV.  $X$  is "Bernoulli RV with parameter  $p$ ", where  $\mathbf{P}[\text{"success"}] = p$  and so PMF  $p(1) = p$ .

$$\mathbf{X} \sim \mathbf{Ber}(p)$$

$$\text{Range: } \{0, 1\}$$

$$\text{PMF: } \mathbf{P}[X = 1] = p(1) = p$$

$$\mathbf{P}[X = 0] = p(0) = 1 - p$$

$$\text{Expectation: } \mathbf{E}[X] = p$$

$$\text{Variance: } \mathbf{V}[X] = p(1 - p)$$

Examples: coin toss, random binary digit, if someone likes a film, the gender of a newborn baby, pas/fail of you taking an exam.



## Bernoulli examples

### Example

You watch a film on Netflix. At the end you click "like" with probability  $p$ . Define a RV representing this event.

\_\_\_\_\_ Answer \_\_\_\_\_

### Example

Two fair 6-sided dice are rolled. Define a random variable  $X$  for a successful roll of two 6's, and failure for anything else.

\_\_\_\_\_ Answer \_\_\_\_\_



# Outline

---

Properties of expectation

Variance

Bernoulli discrete random variable

**Binomial discrete random variable**



# Binomial

## Binomial discrete random variable

A Binomial RV  $X$  represents the number of successes in  $n$  successive independent trials of a Bernoulli experiment.  $X \sim \text{Bin}(n, p)$  is a Binomial RV, where  $p$  is the probability of success in a given trial:

$$X \sim \text{Bin}(n, p)$$

is distributed as a

Range:  $\{0, 1, \dots, n\}$

with parameters

Probability that  $X$  takes on the value  $k$

$$\mathbf{P}[X = k] = p(k) = \binom{n}{k} p^k (1-p)^{n-k}$$

Expectation:  $\mathbf{E}[X] = np$

Variance:  $\mathbf{V}[X] = np(1-p)$

Probability Mass Function for a Binomial

Examples: # heads in  $n$  coin tosses, # of 1's in randomly generated length  $n$  bit string

Note: by Binomial theorem (revision), we can prove  $\sum_{k=0}^n \mathbf{P}[X = k] = 1$ .



## Binomial example

### Example

Let  $X$  be the number of heads after a coin is tossed three times:  
 $X \sim \text{Bin}(3, 0.5)$ . What is the probability of each of the different values of  $X$ ?

Answer



## Binomial RV is sum of Bernoulli RVs

---

Let  $X$  be a Bernoulli RV:  $X \sim \text{Ber}(p)$ . Let  $Y$  be a Binomial RV:  $Y \sim \text{Bin}(n, p)$ .  
Binomial RV = sum of  $n$  independent Bernoulli RVs:

$$Y = \sum_{i=1}^n X_i, \quad X_i \sim \text{Ber}(p)$$

$$\mathbf{E}[Y] = \mathbf{E}\left[\sum_{i=1}^n X_i\right] = \sum_{i=1}^n \mathbf{E}[X_i] = np$$

Note:  $\text{Ber}(p) = \text{Bin}(1, p)$



## Another example

### Example

An off-licence sells cases of wine, each containing 20 bottles. The probability that a bottle is bad is 0.05. The off-licence gives a money-back guarantee that the case will contain no more than one bad bottle. What is the probability that the off-licence will have to give money back?

Answer

- $X$ : # of bad bottles in a case (20 bottles)
- $\mathbf{P}[\text{have to give money back}] = \mathbf{P}[X \geq 2] = 1 - \mathbf{P}[X = 0] - \mathbf{P}[X = 1]$
- $X$  is a binomial RV with parameters  $X \sim \text{Bin}(n = 20, p = 0.05)$ .
- Bernoulli trial: check if a bottle is bad
- $\mathbf{P}[\text{success}] = \mathbf{P}[\text{bottle is bad}] = 0.05$   
 $\mathbf{P}[\text{failure}] = \mathbf{P}[\text{bottle is good}] = 0.95$
- Recall, when  $X \sim \text{Bin}(n, p)$  then  $\mathbf{P}[X = k] = \binom{n}{k} p^k (1 - p)^{n-k}$  thus

$$\mathbf{P}[X \geq 2] = 1 - \mathbf{P}[X = 0] - \mathbf{P}[X = 1]$$



## Visualising Binomial PMFs

$X \sim \text{Bin}(40, 0.3)$ ;  $X \sim \text{Bin}(40, 0.5)$ ;  $X \sim \text{Bin}(40, 0.7)$

