

# Introduction to Probability

Lecture 1: Conditional probabilities and Bayes' theorem

Mateja Jamnik, Thomas Sauerwald

University of Cambridge, Department of Computer Science and Technology

email: {mateja.jamnik,thomas.sauerwald}@cl.cam.ac.uk



UNIVERSITY OF  
CAMBRIDGE

# Outline

---

Logistics, motivation, background

Conditional probability

Bayes' Theorem

Independence





Mateja Jamnik



Thomas Sauerwald

### Rough syllabus:

- Introduction to probability: 1 lecture
- Discrete and continuous random variables: 6 lectures
- Moments and limit theorems: 3 lectures
- Applications/statistics: 2 lectures

### Recommended reading:

- **Ross, S.M. (2014). A First course in probability. Pearson (9th ed.).**
- **Dekking, F.M., et. al. (2005) A modern introduction to probability and statistics. Springer.**
- Bertsekas, D.P. & Tsitsiklis, J.N. (2008). Introduction to probability. Athena Scientific.
- Grimmett, G. & Welsh, D. (2014). Probability: an Introduction. Oxford University Press (2nd ed.).



## Why probability?


---

- Gives us mathematical tools to deal with uncertain events.
- It is used everywhere, especially in applications of machine learning.
- Machine learning: use **probability** to compute predictions about and from data.
- Probability is not statistics:
  - Both about random processes.
  - Probability: logically self-contained, few rules for computing, one correct answer.
  - Statistics: messier, more art, get experimental data and try to draw probabilistic conclusions, no single correct answer.




# Applications of probability

---





# Applications of probability

---




# Applications of probability

## Ranking Websites




# Applications of probability

## Ranking Websites



## Data Mining





# Applications of probability


## Ranking Websites



## Matching



## Data Mining




# Applications of probability


## Ranking Websites



## Matching




$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$




...

## Data Mining




# Applications of probability


## Ranking Websites



## Matching




$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$



...

## Data Mining




# Applications of probability


## Ranking Websites




## Matching




$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$



## Data Mining




## Particle Processes




# Applications of probability


## Ranking Websites




## Matching




$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$




## Data Mining



## Deep Learning




## Particle Processes




# Applications of probability

## Ranking Websites




## Matching




$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$


## Finance




## Data Mining



## Deep Learning



## Particle Processes




# Applications of probability

## Ranking Websites




## Matching




$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$


## Finance




## Medicine




## Data Mining



## Deep Learning



## Particle Processes



- Set theory
  - Counting: product rule, sum rule, inclusion-exclusion
  - Combinatorics: permutations
  - Probability space: sample space, event space
  - Axioms
  - Union bound
- 
- Look for revision material of above on the course website:  
<https://www.cl.cam.ac.uk/teaching/2526/IntroProb/>



# Outline

---

Logistics, motivation, background

Conditional probability

Bayes' Theorem

Independence



## Definition

---

### Conditional probability

Consider an experiment with sample space  $S$ , and two events  $E$  and  $F$ . Then, the (conditional) probability of event  $E$  given  $F$  has occurred (denoted  $\mathbf{P}[E|F]$ ) with  $\mathbf{P}[F] > 0$  is defined by

$$\mathbf{P}[E|F] = \frac{\mathbf{P}[E \cap F]}{\mathbf{P}[F]} = \frac{\mathbf{P}[EF]}{\mathbf{P}[F]}$$



## Definition

---

### Conditional probability

Consider an experiment with sample space  $S$ , and two events  $E$  and  $F$ . Then, the (conditional) probability of event  $E$  given  $F$  has occurred (denoted  $\mathbf{P}[E|F]$ ) with  $\mathbf{P}[F] > 0$  is defined by

$$\mathbf{P}[E|F] = \frac{\mathbf{P}[E \cap F]}{\mathbf{P}[F]} = \frac{\mathbf{P}[EF]}{\mathbf{P}[F]}$$

Sample space: all possible outcomes consistent with  $F$  (i.e.,  $S \cap F = F$ )



## Definition

---

### Conditional probability

Consider an experiment with sample space  $S$ , and two events  $E$  and  $F$ . Then, the (conditional) probability of event  $E$  given  $F$  has occurred (denoted  $\mathbf{P}[E|F]$ ) with  $\mathbf{P}[F] > 0$  is defined by

$$\mathbf{P}[E|F] = \frac{\mathbf{P}[E \cap F]}{\mathbf{P}[F]} = \frac{\mathbf{P}[EF]}{\mathbf{P}[F]}$$

Sample space: all possible outcomes consistent with  $F$  (i.e.,  $S \cap F = F$ )

Event space: all outcomes in  $E$  consistent with  $F$  (i.e.,  $E \cap F$ )



## Definition

---

### Conditional probability

Consider an experiment with sample space  $S$ , and two events  $E$  and  $F$ . Then, the (conditional) probability of event  $E$  given  $F$  has occurred (denoted  $\mathbf{P}[E|F]$ ) with  $\mathbf{P}[F] > 0$  is defined by

$$\mathbf{P}[E|F] = \frac{\mathbf{P}[E \cap F]}{\mathbf{P}[F]} = \frac{\mathbf{P}[EF]}{\mathbf{P}[F]}$$

Sample space: all possible outcomes consistent with  $F$  (i.e.,  $S \cap F = F$ )

Event space: all outcomes in  $E$  consistent with  $F$  (i.e.,  $E \cap F$ )

Note: we assume that all outcomes are equally likely



## Definition

### Conditional probability

Consider an experiment with sample space  $S$ , and two events  $E$  and  $F$ . Then, the (conditional) probability of event  $E$  given  $F$  has occurred (denoted  $\mathbf{P}[E|F]$ ) with  $\mathbf{P}[F] > 0$  is defined by

$$\mathbf{P}[E|F] = \frac{\mathbf{P}[E \cap F]}{\mathbf{P}[F]} = \frac{\mathbf{P}[EF]}{\mathbf{P}[F]}$$

Sample space: all possible outcomes consistent with  $F$  (i.e.,  $S \cap F = F$ )

Event space: all outcomes in  $E$  consistent with  $F$  (i.e.,  $E \cap F$ )

Note: we assume that all outcomes are equally likely

$$\mathbf{P}[E|F] = \frac{\# \text{ outcomes in } E \cap F}{\# \text{ outcomes in } F} = \frac{\frac{\# \text{ outcomes in } E \cap F}{\# \text{ outcomes in } S}}{\frac{\# \text{ outcomes in } F}{\# \text{ outcomes in } S}} = \frac{\mathbf{P}[E \cap F]}{\mathbf{P}[F]}$$



## Example

### Example

Two dice are rolled yielding value  $D_1$  and  $D_2$ . Let  $E$  be event that  $D_1 + D_2 = 4$ .

1. What is  $\mathbf{P}[E]$ ?
2. Let event  $F$  be  $D_1 = 2$ . What is  $\mathbf{P}[E|F]$ ?

Answer



Chain rule

Rearranging the definition of conditional probability gives us:

$$\mathbf{P} [ EF ] = \mathbf{P} [ E|F ] \mathbf{P} [ F ]$$



Chain rule

Rearranging the definition of conditional probability gives us:

$$\mathbf{P} [ EF ] = \mathbf{P} [ E|F ] \mathbf{P} [ F ]$$

Generalisation of the Chain rule:

Multiplication rule

$$\mathbf{P} [ E_1 E_2 \cdots E_n ] = \mathbf{P} [ E_1 ] \mathbf{P} [ E_2 | E_1 ] \mathbf{P} [ E_3 | E_2 E_1 ] \cdots \mathbf{P} [ E_n | E_1 \cdots E_{n-1} ]$$

## Example

### Example

An ordinary deck of 52 playing cards is randomly divided into 4 piles of 13 cards each. What is the probability that each pile has exactly 1 ace?

Answer



## Example

### Example

An ordinary deck of 52 playing cards is randomly divided into 4 piles of 13 cards each. What is the probability that each pile has exactly 1 ace?

Answer

Define:

$E_1 = ace♥$  is in any one pile

$E_2 = ace♥$  and  $ace♠$  are in different piles

$E_3 = ace♥, ace♠$  and  $ace♣$  are in different piles

$E_4 =$  all aces are in different piles



## Example

### Example

An ordinary deck of 52 playing cards is randomly divided into 4 piles of 13 cards each. What is the probability that each pile has exactly 1 ace?

Answer

Define:

$E_1 = \text{ace}\heartsuit$  is in any one pile

$E_2 = \text{ace}\heartsuit$  and  $\text{ace}\spadesuit$  are in different piles

$E_3 = \text{ace}\heartsuit, \text{ace}\spadesuit$  and  $\text{ace}\clubsuit$  are in different piles

$E_4 =$  all aces are in different piles

$$\mathbf{P} [ E_1 E_2 E_3 E_4 ] = \mathbf{P} [ E_1 ] \mathbf{P} [ E_2 | E_1 ] \mathbf{P} [ E_3 | E_1 E_2 ] \mathbf{P} [ E_4 | E_1 E_2 E_3 ]$$



## Example

### Example

An ordinary deck of 52 playing cards is randomly divided into 4 piles of 13 cards each. What is the probability that each pile has exactly 1 ace?

Answer

Define:

$E_1 = \text{ace}\heartsuit$  is in any one pile

$E_2 = \text{ace}\heartsuit$  and  $\text{ace}\spadesuit$  are in different piles

$E_3 = \text{ace}\heartsuit, \text{ace}\spadesuit$  and  $\text{ace}\clubsuit$  are in different piles

$E_4 =$  all aces are in different piles

$$\mathbf{P} [ E_1 E_2 E_3 E_4 ] = \mathbf{P} [ E_1 ] \mathbf{P} [ E_2 | E_1 ] \mathbf{P} [ E_3 | E_1 E_2 ] \mathbf{P} [ E_4 | E_1 E_2 E_3 ]$$

We have  $\mathbf{P} [ E_1 ] = 1$ . For rest we consider complement of next ace being in the same pile and thus have:



# Outline

---

Logistics, motivation, background

Conditional probability

**Bayes' Theorem**

Independence



## Law of total probability

The law of total probability (a.k.a. Partition theorem)

For events  $E$  and  $F$  where  $\mathbf{P}[F] > 0$ , then for any event  $E$

$$\mathbf{P}[E] = \mathbf{P}[EF] + \mathbf{P}[EF^c] = \mathbf{P}[E|F]\mathbf{P}[F] + \mathbf{P}[E|F^c]\mathbf{P}[F^c]$$

In general, for disjoint events  $F_1, F_2, \dots, F_n$  s.t.  $F_1 \cup F_2 \cup \dots \cup F_n = S$ ,

$$\mathbf{P}[E] = \sum_{i=1}^n \mathbf{P}[E|F_i]\mathbf{P}[F_i]$$

Intuition:

Want to know probability of  $E$ . There are two scenarios,  $F$  and  $F^c$ . If we know these and the probability of  $E$  conditioned on each scenario, we can compute the probability of  $E$ .



## Lightbulb example

### Example

There are 3 boxes each containing a different number of light bulbs. The first box has 10 bulbs of which 4 are dead, the second has 6 bulbs of which 1 is dead, and the third box has 8 bulbs of which 3 are dead. What is the probability of a dead bulb being selected when a bulb is chosen at random from one of the 3 boxes (each box has equal chance of being picked)?

Answer



## Lightbulb example

### Example

There are 3 boxes each containing a different number of light bulbs. The first box has 10 bulbs of which 4 are dead, the second has 6 bulbs of which 1 is dead, and the third box has 8 bulbs of which 3 are dead. What is the probability of a dead bulb being selected when a bulb is chosen at random from one of the 3 boxes (each box has equal chance of being picked)?

Answer

Let event  $E$  = "dead bulb is picked", and  $F_1$  = "bulb is picked from first box",  $F_2$  = "bulb is picked from second box" and  $F_3$  = "bulb is picked from third box". We know:



## Lightbulb example

### Example

There are 3 boxes each containing a different number of light bulbs. The first box has 10 bulbs of which 4 are dead, the second has 6 bulbs of which 1 is dead, and the third box has 8 bulbs of which 3 are dead. What is the probability of a dead bulb being selected when a bulb is chosen at random from one of the 3 boxes (each box has equal chance of being picked)?

Answer

Let event  $E$  = "dead bulb is picked", and  $F_1$  = "bulb is picked from first box",  $F_2$  = "bulb is picked from second box" and  $F_3$  = "bulb is picked from third box". We know:

$$\mathbf{P}[E|F_1] = \frac{4}{10}, \mathbf{P}[E|F_2] = \frac{1}{6}, \mathbf{P}[E|F_3] = \frac{3}{8}$$



## Lightbulb example

### Example

There are 3 boxes each containing a different number of light bulbs. The first box has 10 bulbs of which 4 are dead, the second has 6 bulbs of which 1 is dead, and the third box has 8 bulbs of which 3 are dead. What is the probability of a dead bulb being selected when a bulb is chosen at random from one of the 3 boxes (each box has equal chance of being picked)?

Answer

Let event  $E$  = "dead bulb is picked", and  $F_1$  = "bulb is picked from first box",  $F_2$  = "bulb is picked from second box" and  $F_3$  = "bulb is picked from third box". We know:

$$\mathbf{P}[E|F_1] = \frac{4}{10}, \mathbf{P}[E|F_2] = \frac{1}{6}, \mathbf{P}[E|F_3] = \frac{3}{8}$$

We need to compute  $\mathbf{P}[E]$ , and we know that  $\mathbf{P}[F_i] = \frac{1}{3}$ :



## Lightbulb example

### Example

There are 3 boxes each containing a different number of light bulbs. The first box has 10 bulbs of which 4 are dead, the second has 6 bulbs of which 1 is dead, and the third box has 8 bulbs of which 3 are dead. What is the probability of a dead bulb being selected when a bulb is chosen at random from one of the 3 boxes (each box has equal chance of being picked)?

Answer

Let event  $E$  = "dead bulb is picked", and  $F_1$  = "bulb is picked from first box",  $F_2$  = "bulb is picked from second box" and  $F_3$  = "bulb is picked from third box". We know:

$$\mathbf{P}[E|F_1] = \frac{4}{10}, \mathbf{P}[E|F_2] = \frac{1}{6}, \mathbf{P}[E|F_3] = \frac{3}{8}$$

We need to compute  $\mathbf{P}[E]$ , and we know that  $\mathbf{P}[F_i] = \frac{1}{3}$ :

$$\mathbf{P}[E] = \sum_{i=1}^n \mathbf{P}[E|F_i] \mathbf{P}[F_i] = \frac{4}{10} \frac{1}{3} + \frac{1}{6} \frac{1}{3} + \frac{3}{8} \frac{1}{3} = \frac{113}{360} \approx 0.31$$



## Bayes' theorem

---

How many spam emails contain the word "Dear"?

$$\mathbf{P} [ E|F ] = \mathbf{P} [ \text{"Dear"}|\text{spam} ]$$

But how about what is the probability that an email containing "Dear" is spam?

$$\mathbf{P} [ F|E ] = \mathbf{P} [ \text{spam}|\text{"Dear"} ]$$



## Bayes' theorem

How many spam emails contain the word "Dear"?

$$\mathbf{P}[E|F] = \mathbf{P}[\text{"Dear"}|\text{spam}]$$

But how about what is the probability that an email containing "Dear" is spam?

$$\mathbf{P}[F|E] = \mathbf{P}[\text{spam}|\text{"Dear"}]$$

### Bayes' theorem

For any events  $E$  and  $F$  where  $\mathbf{P}[E] > 0$  and  $\mathbf{P}[F] > 0$ ,

$$\mathbf{P}[F|E] = \frac{\mathbf{P}[E|F]\mathbf{P}[F]}{\mathbf{P}[E]}$$

and in expanded form,

$$\mathbf{P}[F|E] = \frac{\mathbf{P}[E|F]\mathbf{P}[F]}{\mathbf{P}[E|F]\mathbf{P}[F] + \mathbf{P}[E|F^c]\mathbf{P}[F^c]} = \frac{\mathbf{P}[E|F]\mathbf{P}[F]}{\sum_{i=1}^n \mathbf{P}[E|F_i]\mathbf{P}[F_i]}$$

using the Law of Total Probability. Note that all events  $F_i$  must be mutually exclusive (non-overlapping) and exhaustive (their union is the complete sample space).



## Example – Do it at home

### Example

60% of all email in 2022 is spam. 20% of spam contains the word "Dear". 1% of non-spam contains the word "Dear". What is the probability that an email is spam given it contains the word "Dear"?

Answer



## Example – Do it at home

### Example

60% of all email in 2022 is spam. 20% of spam contains the word "Dear". 1% of non-spam contains the word "Dear". What is the probability that an email is spam given it contains the word "Dear"?

Answer

- Let event  $E$  = "Dear", event  $F$  = spam.



$$\mathbf{P}[F|E] = \frac{\mathbf{P}[E|F] \cdot \mathbf{P}[F]}{\mathbf{P}[E]}$$

$F$ : hypothesis,  $E$ : evidence



posterior

$$\mathbf{P}[F|E] = \frac{\mathbf{P}[E|F] \cdot \mathbf{P}[F]}{\mathbf{P}[E]}$$

$F$ : hypothesis,  $E$ : evidence

$$\begin{array}{c} \text{posterior} \\ \mathbf{P}[F|E] \end{array} = \frac{\mathbf{P}[E|F] \cdot \mathbf{P}[F]}{\mathbf{P}[E]} \begin{array}{c} \text{prior} \end{array}$$

$F$ : hypothesis,  $E$ : evidence

$\mathbf{P}[F]$ : "prior probability" of hypothesis

## Bayes' terminology

---

$$\begin{array}{ccc} \text{posterior} & \text{likelihood} & \text{prior} \\ \text{P}[F|E] & = & \frac{\text{P}[E|F] \cdot \text{P}[F]}{\text{P}[E]} \end{array}$$

$F$ : hypothesis,  $E$ : evidence

$\text{P}[F]$ : "prior probability" of hypothesis

$\text{P}[E|F]$ : probability of evidence given hypothesis (likelihood)



## Bayes' terminology

The diagram shows the equation for Bayes' theorem with callouts for each term:

$$\mathbf{P}[F|E] = \frac{\mathbf{P}[E|F] \cdot \mathbf{P}[F]}{\mathbf{P}[E]}$$

Callouts:

- posterior:  $\mathbf{P}[F|E]$
- likelihood:  $\mathbf{P}[E|F]$
- prior:  $\mathbf{P}[F]$
- normalisation constant:  $\mathbf{P}[E]$

$F$ : hypothesis,  $E$ : evidence

$\mathbf{P}[F]$ : "prior probability" of hypothesis

$\mathbf{P}[E|F]$ : probability of evidence given hypothesis (likelihood)

$\mathbf{P}[E]$ : calculated by making sure that probabilities of all outcomes sum to 1 (they are "normalised")

## Confusion matrix (error matrix)

Used in classification tasks for predicting output error.

|                     |                                    | True condition                               |   |
|---------------------|------------------------------------|--|---|
|                     |                                    | Condition positive $F$                       | Condition negative $F^c$                      |
| Predicted condition | Predicted condition positive $E$   | <b>True positive</b><br>$\mathbf{P}[E F]$    | <b>False positive</b><br>$\mathbf{P}[E F^c]$  |
|                     | Predicted condition negative $E^c$ | <b>False negative</b><br>$\mathbf{P}[E^c F]$ | <b>True negative</b><br>$\mathbf{P}[E^c F^c]$ |



## Medical testing example

### Example

- A test is 98% effective at detecting the disease COVID-19 ("true positive").
- The test has a "false positive" rate of 1%.
- 0.5% of the population has COVID-19.
- What is the likelihood you have COVID-19 if you test positive?

---

Answer



## Medical testing example

### Example

- A test is 98% effective at detecting the disease COVID-19 ("true positive").
- The test has a "false positive" rate of 1%.
- 0.5% of the population has COVID-19.
- What is the likelihood you have COVID-19 if you test positive?

Answer

- Let  $E$ : test positive,  $F$ : actually have COVID-19.
- Need to find  $\mathbf{P} [ F|E ]$ .



## Bayesian intuition

---

- 33% chance of having COVID-19 after testing positive may seem surprising.



## Bayesian intuition

---

- 33% chance of having COVID-19 after testing positive may seem surprising.
- But the space of facts is now **conditioned** on a positive test result (people who test positive and have COVID-19 **and** people who test positive and don't have COVID-19).



## Bayesian intuition

- 33% chance of having COVID-19 after testing positive may seem surprising.
- But the space of facts is now **conditioned** on a positive test result (people who test positive and have COVID-19 **and** people who test positive and don't have COVID-19).

|             | $F$ yes disease                              | $F^c$ no disease                              |
|-------------|--|---|
| $E$ test+   | True positive<br>$\mathbf{P}[E F] = 0.98$    | False positive<br>$\mathbf{P}[E F^c] = 0.01$  |
| $E^c$ test- | False negative<br>$\mathbf{P}[E^c F] = 0.02$ | True negative<br>$\mathbf{P}[E^c F^c] = 0.99$ |



## Bayesian intuition

- 33% chance of having COVID-19 after testing positive may seem surprising.
- But the space of facts is now **conditioned** on a positive test result (people who test positive and have COVID-19 **and** people who test positive and don't have COVID-19).

|             | $F$ yes disease                              | $F^c$ no disease                              |
|-------------|--|---|
| $E$ test+   | True positive<br>$\mathbf{P}[E F] = 0.98$    | False positive<br>$\mathbf{P}[E F^c] = 0.01$  |
| $E^c$ test- | False negative<br>$\mathbf{P}[E^c F] = 0.02$ | True negative<br>$\mathbf{P}[E^c F^c] = 0.99$ |

- But what is a chance of having COVID-19 if you test and it comes back negative?



## Bayesian intuition

- 33% chance of having COVID-19 after testing positive may seem surprising.
- But the space of facts is now **conditioned** on a positive test result (people who test positive and have COVID-19 **and** people who test positive and don't have COVID-19).

|             | $F$ yes disease                              | $F^c$ no disease                              |
|-------------|--|---|
| $E$ test+   | True positive<br>$\mathbf{P}[E F] = 0.98$    | False positive<br>$\mathbf{P}[E F^c] = 0.01$  |
| $E^c$ test- | False negative<br>$\mathbf{P}[E^c F] = 0.02$ | True negative<br>$\mathbf{P}[E^c F^c] = 0.99$ |

- But what is a chance of having COVID-19 if you test and it comes back negative?

$$\mathbf{P}[F|E^c] = \frac{\mathbf{P}[E^c|F]\mathbf{P}[F]}{\mathbf{P}[E^c|F]\mathbf{P}[F] + \mathbf{P}[E^c|F^c]\mathbf{P}[F^c]} \approx 0.0001$$



## Bayesian intuition

- 33% chance of having COVID-19 after testing positive may seem surprising.
- But the space of facts is now **conditioned** on a positive test result (people who test positive and have COVID-19 **and** people who test positive and don't have COVID-19).

|             | $F$ yes disease                              | $F^c$ no disease                              |
|-------------|--|---|
| $E$ test+   | True positive<br>$\mathbf{P}[E F] = 0.98$    | False positive<br>$\mathbf{P}[E F^c] = 0.01$  |
| $E^c$ test- | False negative<br>$\mathbf{P}[E^c F] = 0.02$ | True negative<br>$\mathbf{P}[E^c F^c] = 0.99$ |

- But what is a chance of having COVID-19 if you test and it comes back negative?

$$\mathbf{P}[F|E^c] = \frac{\mathbf{P}[E^c|F]\mathbf{P}[F]}{\mathbf{P}[E^c|F]\mathbf{P}[F] + \mathbf{P}[E^c|F^c]\mathbf{P}[F^c]} \approx 0.0001$$

- We update our beliefs with Bayes' theorem:  
I have 0.5% chance of having COVID-19. I take the test:
  - Test is positive: **I now have 33% chance of having COVID-19.**
  - Test is negative: **I now have 0.01% chance of having COVID-19.**
- So it makes sense to take the test.



# Outline

---

Logistics, motivation, background

Conditional probability

Bayes' Theorem

Independence



### Independence

Two events  $E$  and  $F$  are independent if and only if

$$\mathbf{P}[EF] = \mathbf{P}[E]\mathbf{P}[F]$$

Otherwise, they are called dependent events.

In general,  $n$  events  $E_1, E_2, \dots, E_n$  are mutually independent if for every subset of these events with  $r$  elements (where  $r \leq n$ ) it holds that

$$\mathbf{P}[E_a E_b \cdots E_r] = \mathbf{P}[E_a]\mathbf{P}[E_b] \cdots \mathbf{P}[E_r]$$



## Independent events

### Independence

Two events  $E$  and  $F$  are independent if and only if

$$\mathbf{P}[EF] = \mathbf{P}[E]\mathbf{P}[F]$$

Otherwise, they are called dependent events.

In general,  $n$  events  $E_1, E_2, \dots, E_n$  are mutually independent if for every subset of these events with  $r$  elements (where  $r \leq n$ ) it holds that

$$\mathbf{P}[E_a E_b \cdots E_r] = \mathbf{P}[E_a]\mathbf{P}[E_b] \cdots \mathbf{P}[E_r]$$

Therefore for 3 events  $E, F, G$  to be independent, we must have

$$\mathbf{P}[EFG] = \mathbf{P}[E]\mathbf{P}[F]\mathbf{P}[G]$$

$$\mathbf{P}[EF] = \mathbf{P}[E]\mathbf{P}[F]$$

$$\mathbf{P}[EG] = \mathbf{P}[E]\mathbf{P}[G]$$

$$\mathbf{P}[FG] = \mathbf{P}[F]\mathbf{P}[G]$$



## Independence of complement

---

Notice an equivalent definition for independent events  $E$  and  $F$  ( $\mathbf{P}[F] > 0$ )

$$\mathbf{P}[E|F] = \mathbf{P}[E]$$

Proof:



## Independence of complement

---

Notice an equivalent definition for independent events  $E$  and  $F$  ( $\mathbf{P}[F] > 0$ )

$$\mathbf{P}[E|F] = \mathbf{P}[E]$$

Proof:

Independence of complement

If events  $E$  and  $F$  are independent, then  $E$  and  $F^c$  are independent:

$$\mathbf{P}[EF^c] = \mathbf{P}[E]\mathbf{P}[F^c]$$

Proof:



## Example

### Example

Each roll of a die is an independent trial. We have two rolls of  $D_1$  and  $D_2$ . Let event  $E : D_1 = 1$ ,  $F : D_2 = 6$  and event  $G : D_1 + D_2 = 7$  (thus  $G = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$ ).

1. Are  $E$  and  $F$  independent?
2. Are  $E$  and  $G$  independent?
3. Are  $E, F, G$  independent?

Answer



### Conditional independence

Two events  $E$  and  $F$  are called conditionally independent given a third event  $G$  if

$$\mathbf{P}[EF|G] = \mathbf{P}[E|G]\mathbf{P}[F|G]$$

Or equivalently,

$$\mathbf{P}[E|FG] = \mathbf{P}[E|G]$$

Notice that:

- Dependent events can become conditionally independent.
- Independent events can become conditionally dependent.
- Knowing when conditioning breaks or creates independence is a big part of building complex probabilistic models.



## Example revisited

### Example

Each roll of a die is an independent trial. We have two rolls of  $D_1$  and  $D_2$ . Let event  $E : D_1 = 1$ ,  $F : D_2 = 6$  and event  $G : D_1 + D_2 = 7$  (thus  $G = \{(1, 6), (2, 5), (3, 4), (4, 3), (5, 2), (6, 1)\}$ ).

1. Are  $E$  and  $F$  independent?
2. Are  $E$  and  $F$  independent given  $G$ ?

Answer



## Summary of conditional probability

---

Conditioning on event  $G$ :

| Name of rule             | Original rule  | Conditional rule   |
|--------------------------|--|--|
| 1st axiom of probability | $0 \leq \mathbf{P}[E] \leq 1$  | $0 \leq \mathbf{P}[E G] \leq 1$  |
| Complement               | $\mathbf{P}[E] = 1 - \mathbf{P}[E^c]$                                  | $\mathbf{P}[E G] = 1 - \mathbf{P}[E^c G]$                                    |
| Chain rule               | $\mathbf{P}[EF] = \mathbf{P}[E F]\mathbf{P}[F]$                        | $\mathbf{P}[EF G] = \mathbf{P}[E FG]\mathbf{P}[F G]$                         |
| Bayes' theorem           | $\mathbf{P}[F E] = \frac{\mathbf{P}[E F]\mathbf{P}[F]}{\mathbf{P}[E]}$ | $\mathbf{P}[F EG] = \frac{\mathbf{P}[E FG]\mathbf{P}[F G]}{\mathbf{P}[E G]}$ |

