

Introduction to Probability

Lectures 9: Central Limit Theorem

Mateja Jamnik, [Thomas Sauerwald](#)

University of Cambridge, Department of Computer Science and Technology
email: {mateja.jamnik,thomas.sauerwald}@cl.cam.ac.uk

Easter 2026



UNIVERSITY OF
CAMBRIDGE

Outline

Recap: Weak Law of Large Numbers


Central Limit Theorem

Illustrations

Examples

Weak Law of Large Numbers (4/4)


Weak Law of Large Numbers: For any $\epsilon > 0$, $\lim_{n \rightarrow \infty} \mathbf{P} \left[|\bar{X}_n - \mu| > \epsilon \right] = 0$



Weak Law of Large Numbers (4/4)

Weak Law of Large Numbers: For any $\epsilon > 0$, $\lim_{n \rightarrow \infty} \mathbf{P} \left[|\bar{X}_n - \mu| > \epsilon \right] = 0$


$$\Rightarrow \epsilon = 0.2, \delta = 0.25, \exists N. \forall n \geq N. \mathbf{P} \left[|\bar{X}_n - \mu| > 0.2 \right] \leq 0.25$$



Weak Law of Large Numbers (4/4)

Weak Law of Large Numbers: For any $\epsilon > 0$, $\lim_{n \rightarrow \infty} \mathbf{P} \left[|\bar{X}_n - \mu| > \epsilon \right] = 0$

$$\Rightarrow \epsilon = 0.2, \delta = 0.25, \exists N. \forall n \geq N. \mathbf{P} \left[|\bar{X}_n - \mu| > 0.2 \right] \leq 0.25$$




Weak Law of Large Numbers (4/4)

Weak Law of Large Numbers: For any $\epsilon > 0$, $\lim_{n \rightarrow \infty} \mathbf{P} \left[|\bar{X}_n - \mu| > \epsilon \right] = 0$

$$\Rightarrow \epsilon = 0.2, \delta = 0.25, \exists N. \forall n \geq N. \mathbf{P} \left[|\bar{X}_n - \mu| > 0.2 \right] \leq 0.25$$

$$N = \frac{\sigma^2}{\delta \epsilon^2} = \frac{1}{0.25 \cdot 0.2^2} = 100$$




Weak Law of Large Numbers (4/4)

Weak Law of Large Numbers: For any $\epsilon > 0$, $\lim_{n \rightarrow \infty} \mathbf{P} \left[|\bar{X}_n - \mu| > \epsilon \right] = 0$

$$\Rightarrow \epsilon = 0.2, \delta = 0.25, \exists N. \forall n \geq N. \mathbf{P} \left[|\bar{X}_n - \mu| > 0.2 \right] \leq 0.25$$

$$N = \frac{\sigma^2}{\delta \epsilon^2} = \frac{1}{0.25 \cdot 0.2^2} = 100$$




Weak Law of Large Numbers (4/4)

Weak Law of Large Numbers: For any $\epsilon > 0$, $\lim_{n \rightarrow \infty} \mathbf{P} \left[|\bar{X}_n - \mu| > \epsilon \right] = 0$

$$\Rightarrow \epsilon = 0.2, \delta = 0.25, \exists N. \forall n \geq N. \mathbf{P} \left[|\bar{X}_n - \mu| > 0.2 \right] \leq 0.25$$

$$N = \frac{\sigma^2}{\delta \epsilon^2} = \frac{1}{0.25 \cdot 0.2^2} = 100$$




Weak Law of Large Numbers (4/4)

Weak Law of Large Numbers: For any $\epsilon > 0$, $\lim_{n \rightarrow \infty} \mathbf{P} \left[|\bar{X}_n - \mu| > \epsilon \right] = 0$

$$\Rightarrow \epsilon = 0.2, \delta = 0.25, \exists N. \forall n \geq N. \mathbf{P} \left[|\bar{X}_n - \mu| > 0.2 \right] \leq 0.25$$

$$N = \frac{\sigma^2}{\delta \epsilon^2} = \frac{1}{0.25 \cdot 0.2^2} = 100$$



Outline

Recap: Weak Law of Large Numbers

Central Limit Theorem

Illustrations

Examples

Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2

Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2



Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2

The Sum

- Let $\tilde{X}_n := \sum_{i=1}^n X_i$ (often denoted by S_n)
- The variance is $\mathbf{V}[\tilde{X}_n] = n\sigma^2 \rightarrow \infty$




Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2

The Sum

- Let $\tilde{X}_n := \sum_{i=1}^n X_i$ (often denoted by S_n)
- The variance is $\mathbf{V}[\tilde{X}_n] = n\sigma^2 \rightarrow \infty$




Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2

The Sum

- Let $\tilde{X}_n := \sum_{i=1}^n X_i$ (often denoted by S_n)
- The variance is $\mathbf{V}[\tilde{X}_n] = n\sigma^2 \rightarrow \infty$



The Sample Average (Sample Mean)


- Let $\bar{X}_n := \frac{1}{n} \cdot \sum_{i=1}^n X_i$
- The variance is $\mathbf{V}[\bar{X}_n] = \sigma^2/n \rightarrow 0$

Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2


The Sum

- Let $\tilde{X}_n := \sum_{i=1}^n X_i$ (often denoted by S_n)
- The variance is $\mathbf{V}[\tilde{X}_n] = n\sigma^2 \rightarrow \infty$



The Sample Average (Sample Mean)

- Let $\bar{X}_n := \frac{1}{n} \cdot \sum_{i=1}^n X_i$
- The variance is $\mathbf{V}[\bar{X}_n] = \sigma^2/n \rightarrow 0$




Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2


The Sum

- Let $\tilde{X}_n := \sum_{i=1}^n X_i$ (often denoted by S_n)
- The variance is $\mathbf{V}[\tilde{X}_n] = n\sigma^2 \rightarrow \infty$



The Sample Average (Sample Mean)

- Let $\bar{X}_n := \frac{1}{n} \cdot \sum_{i=1}^n X_i$
- The variance is $\mathbf{V}[\bar{X}_n] = \sigma^2/n \rightarrow 0$



The Proper Scaling (Standardising, see Lec. 5)


- Let $Z_n := \frac{1}{\sqrt{n} \cdot \sigma} \cdot \sum_{i=1}^n X_i$
- The variance is $\mathbf{V}[Z_n] = 1$

Towards the CLT: Finding the Right Scaling

- Let X_1, X_2, \dots i.i.d. with $\mu = 0$ and finite σ^2


The Sum

- Let $\tilde{X}_n := \sum_{i=1}^n X_i$ (often denoted by S_n)
- The variance is $\mathbf{V}[\tilde{X}_n] = n\sigma^2 \rightarrow \infty$




The Sample Average (Sample Mean)

- Let $\bar{X}_n := \frac{1}{n} \cdot \sum_{i=1}^n X_i$
- The variance is $\mathbf{V}[\bar{X}_n] = \sigma^2/n \rightarrow 0$



The Proper Scaling (Standardising, see Lec. 5)

- Let $Z_n := \frac{1}{\sqrt{n} \cdot \sigma} \cdot \sum_{i=1}^n X_i$
- The variance is $\mathbf{V}[Z_n] = 1$



Central Limit Theorem



A. de Moivre (1667-1754) P.-S. de Laplace (1749-1827) C. Gauss (1777-1855) A. Lyapunov (1857-1918) C. Lindeberg (1876-1932)

Central Limit Theorem



A. de Moivre (1667-1754) P.-S. de Laplace (1749-1827) C. Gauss (1777-1855) A. Lyapunov (1857-1918) C. Lindeberg (1876-1932)

Central Limit Theorem

Let X_1, X_2, \dots be any sequence of independent identically distributed random variables with finite expectation μ and finite variance σ^2 . Let

$$Z_n := \sqrt{n} \cdot \frac{\bar{X}_n - \mu}{\sigma}$$

Central Limit Theorem



A. de Moivre (1667-1754) P.-S. de Laplace (1749-1827) C. Gauss (1777-1855) A. Lyapunov (1857-1918) C. Lindeberg (1876-1932)

Central Limit Theorem

Let X_1, X_2, \dots be any sequence of independent identically distributed random variables with finite expectation μ and finite variance σ^2 . Let

$$Z_n := \sqrt{n} \cdot \frac{\bar{X}_n - \mu}{\sigma} = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right)$$

Central Limit Theorem



A. de Moivre (1667-1754) P.-S. de Laplace (1749-1827) C. Gauss (1777-1855) A. Lyapunov (1857-1918) C. Lindeberg (1876-1932)

Central Limit Theorem

Let X_1, X_2, \dots be any sequence of independent identically distributed random variables with finite expectation μ and finite variance σ^2 . Let

$$Z_n := \sqrt{n} \cdot \frac{\bar{X}_n - \mu}{\sigma} = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right)$$

Then for any number $a \in \mathbb{R}$, it holds that

$$\lim_{n \rightarrow \infty} F_{Z_n}(a) = \Phi(a)$$

where Φ is the distribution function of the $\mathcal{N}(0, 1)$ distribution.

Central Limit Theorem



A. de Moivre (1667-1754) P.-S. de Laplace (1749-1827) C. Gauss (1777-1855) A. Lyapunov (1857-1918) C. Lindeberg (1876-1932)

Central Limit Theorem

Let X_1, X_2, \dots be any sequence of independent identically distributed random variables with finite expectation μ and finite variance σ^2 . Let

$$Z_n := \sqrt{n} \cdot \frac{\bar{X}_n - \mu}{\sigma} = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right)$$

Then for any number $a \in \mathbb{R}$, it holds that

$$\lim_{n \rightarrow \infty} F_{Z_n}(a) = \Phi(a) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx,$$

where Φ is the distribution function of the $\mathcal{N}(0, 1)$ distribution.

Central Limit Theorem



A. de Moivre (1667-1754) P.-S. de Laplace (1749-1827) C. Gauss (1777-1855) A. Lyapunov (1857-1918) C. Lindeberg (1876-1932)

Central Limit Theorem

Let X_1, X_2, \dots be any sequence of independent identically distributed random variables with finite expectation μ and finite variance σ^2 . Let

$$Z_n := \sqrt{n} \cdot \frac{\bar{X}_n - \mu}{\sigma} = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right)$$

Then for any number $a \in \mathbb{R}$, it holds that

$$\lim_{n \rightarrow \infty} F_{Z_n}(a) = \Phi(a) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx,$$

where Φ is the distribution function of the $\mathcal{N}(0, 1)$ distribution.

In words: the distribution of Z_n **always** converges to the distribution function Φ of the standard normal distribution.

Comments on the CLT

- one of the most remarkable results in probability/statistics
- extremely useful tool in data analysis or physical measurements
 - we may not know the actual distribution in real-world, and CLT says we don't have to(!)
 - adding up independent noises in measurements leads to an error following the Normal distribution
 - applies also to sums of random variables which may be unbounded

Comments on the CLT

- one of the most remarkable results in probability/statistics
- extremely useful tool in data analysis or physical measurements
 - we may not know the actual distribution in real-world, and CLT says we don't have to(!)
 - adding up independent noises in measurements leads to an error following the Normal distribution
 - applies also to sums of random variables which may be unbounded

- catch: the CLT only holds **approximately**, i.e., for large n

When is the approximation good?

Comments on the CLT

- one of the most remarkable results in probability/statistics
- extremely useful tool in data analysis or physical measurements
 - we may not know the actual distribution in real-world, and CLT says we don't have to(!)
 - adding up independent noises in measurements leads to an error following the Normal distribution
 - applies also to sums of random variables which may be unbounded
- catch: the CLT only holds **approximately**, i.e., for large n

When is the approximation good?

- usually $n \geq 10$ or $n \geq 15$ is sufficient in practice
- approximation tends to be worse when threshold a is far from 0, distribution of X_i 's asymmetric, bimodal or discrete
- (for a result quantifying the approximation error: Berry-Esseen-Theorem)

Outline

Recap: Weak Law of Large Numbers

Central Limit Theorem

Illustrations

Examples

Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^1 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^2 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^3 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^4 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^5 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^6 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^7 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^8 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^9 X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{10} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{11} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{12} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{13} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{14} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{15} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{16} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{17} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{18} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{19} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{20} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{21} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{22} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{23} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{24} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{25} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{26} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{27} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{28} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{29} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$\mathbf{P} \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$




Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$

By the CLT:

$$Z_n = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right) \xrightarrow{n \rightarrow \infty} Z \sim \mathcal{N}(0, 1)$$




Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$

By the CLT:

$$Z_n = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right) \xrightarrow{n \rightarrow \infty} Z \sim \mathcal{N}(0, 1)$$
$$\Rightarrow \sum_{i=1}^n X_i$$




Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$

By the CLT:

$$Z_n = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right) \xrightarrow{n \rightarrow \infty} Z \sim \mathcal{N}(0, 1)$$

$$\Rightarrow \sum_{i=1}^n X_i \approx \sqrt{n} \cdot \sigma Z$$




Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$

By the CLT:

$$Z_n = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right) \xrightarrow{n \rightarrow \infty} Z \sim \mathcal{N}(0, 1)$$

$$\Rightarrow \sum_{i=1}^n X_i \approx \sqrt{n} \cdot \sigma Z \sim \mathcal{N}(0, n \cdot \sigma^2)$$




Illustration of CLT (1/4)

$$P \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$

By the CLT:

$$Z_n = \frac{1}{\sqrt{n} \cdot \sigma} \cdot \left(\sum_{i=1}^n X_i - n \cdot \mu \right) \xrightarrow{n \rightarrow \infty} Z \sim \mathcal{N}(0, 1)$$

$$\Rightarrow \sum_{i=1}^n X_i \approx \sqrt{n} \cdot \sigma Z \sim \mathcal{N}(0, n \cdot \sigma^2)$$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^1 X_j = x \right]$$

$$\blacksquare \mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$$

$$\blacksquare \sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^2 X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^3 X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^4 X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$



Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^5 X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^6 X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^7 X_j = x \right]$$

$$\blacksquare \mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$$

$$\blacksquare \sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^8 X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^9 X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{10} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{11} X_j = x \right]$$

$$\blacksquare \mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$$

$$\blacksquare \sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$$




Illustration of CLT (2/4)

$$P \left[\sum_{j=1}^{12} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{13} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{14} X_j = x \right]$$

$$\blacksquare \mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$$

$$\blacksquare \sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{15} X_j = x \right]$$

$$\blacksquare \mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$$

$$\blacksquare \sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{16} X_j = x \right]$$

$$\blacksquare \mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$$

$$\blacksquare \sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{17} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{18} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{19} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{20} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{21} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{22} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{23} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{24} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$P \left[\sum_{j=1}^{25} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{26} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$P \left[\sum_{j=1}^{27} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{28} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{29} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (2/4)

$$\mathbf{P} \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = 0.15 \cdot (-3) + 0.1 \cdot (-2) + 0.05 \cdot (-1) + 0.7 \cdot 1 = 0$
- $\sigma^2 = 0.15 \cdot 9 + 0.1 \cdot 4 + 0.05 \cdot 1 + 0.7 \cdot 1 = 2.5$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^1 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^2 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^3 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^4 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^5 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^6 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^7 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^8 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^9 X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{10} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{11} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{12} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{13} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{14} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{15} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{16} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{17} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{18} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{19} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{20} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{21} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{22} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{23} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{24} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{25} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{26} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{27} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{28} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{29} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$P \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{30} X_j = x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{30} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part I) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{30} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^1 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^2 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^3 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^4 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^5 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^6 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^7 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^8 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^9 X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{10} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{11} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{12} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{13} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{14} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{15} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{16} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{17} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{18} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{19} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{20} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{21} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{22} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{23} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{24} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{25} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{26} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{27} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{28} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{29} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (3, Part II) (Distribution from Lecture 8)

$$\mathbf{P} \left[\sum_{j=1}^{30} X_j \leq x \right]$$

- $\mu = \frac{1}{2} \cdot (-1) + \frac{1}{2} \cdot 1 = 0$
- $\sigma^2 = \frac{1}{2} \cdot (-1)^2 + \frac{1}{2} \cdot 1^2 = 1$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[X_1 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$\mathbf{P}[X_1 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_1 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_1 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_1 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_1 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_2 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_3 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_4 = x]$

- First switch to random variable $Z_n := \frac{\sqrt{n}}{\sigma} \cdot \bar{X}_n = \frac{1}{\sqrt{n}\sigma} \cdot \sum_{i=1}^n X_i$
- This random variable takes values which are $\frac{1}{\sqrt{n}\sigma}$ apart
- In histogram, always adjust probability so that total area equals 1




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_5 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_6 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_7 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_8 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_9 = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{10} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{11} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{12} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{13} = x]$

- First switch to random variable $Z_n := \frac{\sqrt{n}}{\sigma} \cdot \bar{X}_n = \frac{1}{\sqrt{n}\sigma} \cdot \sum_{i=1}^n X_i$
- This random variable takes values which are $\frac{1}{\sqrt{n}\sigma}$ apart
- In histogram, always adjust probability so that total area equals 1




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{14} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{15} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{16} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{17} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{18} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{19} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{20} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{21} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{22} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{23} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{24} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{25} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{26} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{27} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{28} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{29} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{30} = x]$




Illustration of CLT (4, Part I) with Standardising

$$\blacksquare \mu = \frac{1}{3} \cdot (-1) + \frac{1}{3} \cdot 0 + \frac{1}{3} \cdot 1 = 0$$

$$\blacksquare \sigma^2 = \frac{1}{3} \cdot (-1)^2 + \frac{1}{3} \cdot 0^2 + \frac{1}{3} \cdot 1^2 = \frac{2}{3}$$

$P[Z_{30} = x]$




Illustration of CLT (4, Part II) with Standardising




Fig. 14.2. Densities of standardized averages Z_n . Left column: from a gamma density; right column: from a bimodal density. Dotted line: $N(0, 1)$ probability density.

Source: Dekking et al., Modern Introduction to Statistics

Outline

Recap: Weak Law of Large Numbers

Central Limit Theorem

Illustrations

Examples

Recall: Standard Normal Table

Section 5.4 Normal Random Variables 201

TABLE 5.1: AREA $\Phi(x)$ UNDER THE STANDARD NORMAL CURVE TO THE LEFT OF X

X	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

Source: Ross, Probability 8th ed.

$$Z \sim \mathcal{N}(0, 1) \quad \mathbf{P}[Z \leq x] = \Phi(x)$$

Recall: Standard Normal Table

Section 5.4 Normal Random Variables 201

TABLE 5.1: AREA $\Phi(x)$ UNDER THE STANDARD NORMAL CURVE TO THE LEFT OF X

X	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9988	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

Source: Ross, Probability 8th ed.

Question: What if we need $\Phi(x)$ for negative x ?

$$Z \sim \mathcal{N}(0, 1) \quad \mathbf{P}[Z \leq x] = \Phi(x)$$

Recall: Standard Normal Table

Section 5.4 Normal Random Variables 201

TABLE 5.1: AREA $\Phi(x)$ UNDER THE STANDARD NORMAL CURVE TO THE LEFT OF X

X	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998

Source: Ross, Probability 8th ed.

Question: What if we need $\Phi(x)$ for negative x ?

$$Z \sim \mathcal{N}(0, 1) \quad \mathbf{P}[Z \leq x] = \Phi(x)$$

Due to symmetry of density we have $\Phi(x) = 1 - \Phi(-x)$.

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

_____ Answer _____

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1 - p) = 3/16$.

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1 - p) = 3/16$.
- Applying the **CLT** yields:

$$\mathbf{P}[X \geq 6] = \mathbf{P}\left[\sum_{i=1}^n X_i \geq 6\right]$$

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1 - p) = 3/16$.
- Applying the **CLT** yields:

$$\begin{aligned}\mathbf{P}[X \geq 6] &= \mathbf{P}\left[\sum_{i=1}^n X_i \geq 6\right] \\ &= \mathbf{P}\left[\frac{\sum_{i=1}^n X_i - n\mu}{\sqrt{n\sigma}} \geq \frac{6 - n\mu}{\sqrt{n\sigma}}\right]\end{aligned}$$

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1 - p) = 3/16$.
- Applying the **CLT** yields:

$$\begin{aligned}\mathbf{P}[X \geq 6] &= \mathbf{P}\left[\sum_{i=1}^n X_i \geq 6\right] \\ &= \mathbf{P}\left[\frac{\sum_{i=1}^n X_i - n\mu}{\sqrt{n\sigma}} \geq \frac{6 - n\mu}{\sqrt{n\sigma}}\right] \\ &= \mathbf{P}\left[Z_{10} \geq \frac{6 - 2.5}{\sqrt{10} \cdot \sqrt{3/16}}\right]\end{aligned}$$

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1 - p) = 3/16$.
- Applying the **CLT** yields:

$$\begin{aligned}\mathbf{P}[X \geq 6] &= \mathbf{P}\left[\sum_{i=1}^n X_i \geq 6\right] \\ &= \mathbf{P}\left[\frac{\sum_{i=1}^n X_i - n\mu}{\sqrt{n\sigma}} \geq \frac{6 - n\mu}{\sqrt{n\sigma}}\right] \\ &= \mathbf{P}\left[Z_{10} \geq \frac{6 - 2.5}{\sqrt{10} \cdot \sqrt{3/16}}\right] \approx 1 - \Phi(2.56) \approx 0.0052.\end{aligned}$$

Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1 - p) = 3/16$.
- Applying the **CLT** yields:

$$\begin{aligned}\mathbf{P}[X \geq 6] &= \mathbf{P}\left[\sum_{i=1}^n X_i \geq 6\right] \\ &= \mathbf{P}\left[\frac{\sum_{i=1}^n X_i - n\mu}{\sqrt{n\sigma}} \geq \frac{6 - n\mu}{\sqrt{n\sigma}}\right] \\ &= \mathbf{P}\left[Z_{10} \geq \frac{6 - 2.5}{\sqrt{10} \cdot \sqrt{3/16}}\right] \approx 1 - \Phi(2.56) \approx 0.0052.\end{aligned}$$

True value is 0.0197. Error lies in the discretisation!


Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1 - p) = 3/16$.
- Applying the **CLT** yields:


$$\begin{aligned} \mathbf{P}[X \geq 6] &= \mathbf{P}\left[\sum_{i=1}^n X_i \geq 6\right] \\ &= \mathbf{P}\left[\frac{\sum_{i=1}^n X_i - n\mu}{\sqrt{n\sigma}} \geq \frac{6 - n\mu}{\sqrt{n\sigma}}\right] \\ &= \mathbf{P}\left[Z_{10} \geq \frac{6 - 2.5}{\sqrt{10} \cdot \sqrt{3/16}}\right] \approx 1 - \Phi(2.56) \approx 0.0052. \end{aligned}$$

True value is 0.0197. Error lies in the discretisation!


Normal Approximation of the Binomial Distribution

Example 1

Suppose you are attending a multiple-choice exam of 10 questions and you are completely unprepared. Each question has 4 choices, and you are going to pass the exam if you **guess** at least 6 correct answers. Use the normal approximation to estimate the probability of passing.

Answer

- Let $X \sim \text{Bin}(10, 1/4)$. We are interested in $\mathbf{P}[X \geq 6]$.
- Note $X := \sum_{i=1}^n X_i$, where each $X_i \sim \text{Ber}(p)$ and $n = 10, p = 1/4$.
 $\Rightarrow \mu = 1/4$ and $\sigma^2 = p(1-p) = 3/16$.
- Applying the **CLT** yields:


$$\begin{aligned} \mathbf{P}[X \geq 6] &= \mathbf{P}\left[\sum_{i=1}^n X_i \geq 6\right] \\ &= \mathbf{P}\left[\frac{\sum_{i=1}^n X_i - n\mu}{\sqrt{n\sigma}} \geq \frac{6 - n\mu}{\sqrt{n\sigma}}\right] \\ &= \mathbf{P}\left[Z_{10} \geq \frac{6 - 2.5}{\sqrt{10} \cdot \sqrt{3/16}}\right] \approx 1 - \Phi(2.56) \approx 0.0052. \end{aligned}$$

continuity correction: a better approximation is obtained by $\mathbf{P}\left[\sum_{i=1}^n X_i \geq 5.5\right] \rightsquigarrow \approx 0.0143$

True value is 0.0197. Error lies in the discretisation!

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?

Answer

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?

Answer

- We have $X_1, X_2, \dots, X_n \sim \text{Exp}(1/2)$, where n is unknown.

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?

Answer

- We have $X_1, X_2, \dots, X_n \sim \text{Exp}(1/2)$, where n is unknown.
- Recall that $\mu = \sigma = 2$.

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?

Answer

- We have $X_1, X_2, \dots, X_n \sim \text{Exp}(1/2)$, where n is unknown.
- Recall that $\mu = \sigma = 2$.
- By the CLT,

$$\mathbf{P} \left[\sum_{i=1}^n X_i \leq 100 \right]$$

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?

Answer

- We have $X_1, X_2, \dots, X_n \sim \text{Exp}(1/2)$, where n is unknown.
- Recall that $\mu = \sigma = 2$.
- By the CLT,

$$\mathbf{P} \left[\sum_{i=1}^n X_i \leq 100 \right] = \mathbf{P} \left[\frac{\sum_{i=1}^n X_i - 2n}{2\sqrt{n}} \leq \frac{100 - 2n}{2\sqrt{n}} \right]$$

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?

Answer

- We have $X_1, X_2, \dots, X_n \sim \text{Exp}(1/2)$, where n is unknown.
- Recall that $\mu = \sigma = 2$.
- By the CLT,

$$\begin{aligned} \mathbf{P} \left[\sum_{i=1}^n X_i \leq 100 \right] &= \mathbf{P} \left[\frac{\sum_{i=1}^n X_i - 2n}{2\sqrt{n}} \leq \frac{100 - 2n}{2\sqrt{n}} \right] \\ &\approx \Phi \left(\frac{100 - 2n}{2\sqrt{n}} \right) \stackrel{!}{=} 0.95. \end{aligned}$$

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?

Answer

- We have $X_1, X_2, \dots, X_n \sim \text{Exp}(1/2)$, where n is unknown.
- Recall that $\mu = \sigma = 2$.
- By the CLT,

$$\begin{aligned} \mathbf{P} \left[\sum_{i=1}^n X_i \leq 100 \right] &= \mathbf{P} \left[\frac{\sum_{i=1}^n X_i - 2n}{2\sqrt{n}} \leq \frac{100 - 2n}{2\sqrt{n}} \right] \\ &\approx \Phi \left(\frac{100 - 2n}{2\sqrt{n}} \right) \stackrel{!}{=} 0.95. \end{aligned}$$

- Using a normal table (looking for value 0.95) yields: $\frac{100 - 2n}{2\sqrt{n}} = 1.645$.

A “Reverse” Application of the CLT

Example 2

Suppose we are sequentially loading one container with packets, whose weights are i.i.d. exponential variables with parameter $\lambda = 1/2$. The container has a capacity of 100 weight units. How many packets can we load so that we meet the capacity threshold with at least .95 probability?


Answer

- We have $X_1, X_2, \dots, X_n \sim \text{Exp}(1/2)$, where n is unknown.
- Recall that $\mu = \sigma = 2$.
- By the CLT,

$$\begin{aligned} \mathbf{P} \left[\sum_{i=1}^n X_i \leq 100 \right] &= \mathbf{P} \left[\frac{\sum_{i=1}^n X_i - 2n}{2\sqrt{n}} \leq \frac{100 - 2n}{2\sqrt{n}} \right] \\ &\approx \Phi \left(\frac{100 - 2n}{2\sqrt{n}} \right) \stackrel{!}{=} 0.95. \end{aligned}$$

- Using a normal table (looking for value 0.95) yields: $\frac{100-2n}{2\sqrt{n}} = 1.645$.
- ⇒ Solving the quadratic gives $n \leq 39.6$ (so $n \leq 39$)

A Sample of 100 Exponential Random Variables $Exp(1/2)$



Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

-
- Answer _____
- Markov: $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

-
- Answer
- Markov: $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

-
- Answer
- Markov: $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.
 $\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666$.
 - Chebyshev: $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

-
- Answer
- Markov: $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- Chebyshev: $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- Markov: $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- Chebyshev: $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

- _____ Answer _____
- Markov: $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- Chebyshev: $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- Central Limit Theorem: First standardise:

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- Markov: $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- Chebyshev: $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- Central Limit Theorem: First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n} \cdot 1/2}$

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- **Markov:** $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- **Chebyshev:** $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- **Central Limit Theorem:** First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n} \cdot 1/2}$

$$\mathbf{P}[X \geq 74.5] = \mathbf{P}\left[Z_n \geq \frac{74.5 - n \cdot 1/2}{\sqrt{n} \cdot 1/2}\right] \approx 1 - \Phi(4.9) = 4.79 \cdot 10^{-7}$$

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- **Markov:** $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- **Chebyshev:** $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- **Central Limit Theorem:** First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n} \cdot 1/2}$

$$\mathbf{P}[X \geq 74.5] = \mathbf{P}\left[Z_n \geq \frac{74.5 - n \cdot 1/2}{\sqrt{n} \cdot 1/2}\right] \approx 1 - \Phi(4.9) = 4.79 \cdot 10^{-7}$$

- exact probability is $2.82 \cdot 10^{-7}$

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- **Markov:** $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- **Chebyshev:** $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- **Central Limit Theorem:** First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n} \cdot 1/2}$

$$\mathbf{P}[X \geq 74.5] = \mathbf{P}\left[Z_n \geq \frac{74.5 - n \cdot 1/2}{\sqrt{n} \cdot 1/2}\right] \approx 1 - \Phi(4.9) = 4.79 \cdot 10^{-7}$$

- exact probability is $2.82 \cdot 10^{-7}$

CLT gives a much better result (but requires i.i.d.)

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- **Markov:** $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- **Chebyshev:** $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- **Central Limit Theorem:** First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n} \cdot 1/2}$

$$\mathbf{P}[X \geq 74.5] = \mathbf{P}\left[Z_n \geq \frac{74.5 - n \cdot 1/2}{\sqrt{n} \cdot 1/2}\right] \approx 1 - \Phi(4.9) = 4.79 \cdot 10^{-7}$$

- exact probability is $2.82 \cdot 10^{-7}$

CLT gives a much better result (but requires i.i.d.)

- **Side Note:** without continuity correction, we have 75 instead 74.5:

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- **Markov:** $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- **Chebyshev:** $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- **Central Limit Theorem:** First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n} \cdot 1/2}$

$$\mathbf{P}[X \geq 74.5] = \mathbf{P}\left[Z_n \geq \frac{74.5 - n \cdot 1/2}{\sqrt{n} \cdot 1/2}\right] \approx 1 - \Phi(4.9) = 4.79 \cdot 10^{-7}$$

- exact probability is $2.82 \cdot 10^{-7}$

CLT gives a much better result (but requires i.i.d.)

- **Side Note:** without continuity correction, we have 75 instead 74.5:

- This leads to $1 - \Phi(5) = 2.86 \cdot 10^{-7}$

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- **Markov:** $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- **Chebyshev:** $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- **Central Limit Theorem:** First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n} \cdot 1/2}$

$$\mathbf{P}[X \geq 74.5] = \mathbf{P}\left[Z_n \geq \frac{74.5 - n \cdot 1/2}{\sqrt{n} \cdot 1/2}\right] \approx 1 - \Phi(4.9) = 4.79 \cdot 10^{-7}$$

- exact probability is $2.82 \cdot 10^{-7}$

CLT gives a much better result (but requires i.i.d.)

- **Side Note:** without continuity correction, we have 75 instead 74.5:

- This leads to $1 - \Phi(5) = 2.86 \cdot 10^{-7}$
- Issue: threshold too large ($\mathbf{P}[X \geq a] \approx \mathbf{P}[X = a]$) \Rightarrow CLT less precise

Comparison between Markov, Chebyshev and CLT

Example 3

Consider $n = 100$ independent coin flips. Estimate the probability that the number of heads is greater or equal than 75.

Answer

- **Markov:** $X = \sum_{i=1}^{100} X_i$, $X_i \in \{0, 1\}$ and $\mathbf{E}[X] = 100 \cdot \frac{1}{2} = 50$.

$$\mathbf{P}[X \geq 3/2 \cdot \mathbf{E}[X]] \leq 2/3 = 0.666.$$

- **Chebyshev:** $\mathbf{V}[X] = \sum_{i=1}^{100} \mathbf{V}[X_i] = 100 \cdot (1/2)^2 = 25$.

$$\mathbf{P}[|X - \mu| \geq 25] \leq \frac{\mathbf{V}[X]}{25^2} = \frac{1}{25} = 0.04.$$

As X is symmetric, we could deduce probability is at most 0.02.

- **Central Limit Theorem:** First standardise: $Z_n = \frac{X - n \cdot 1/2}{\sqrt{n \cdot 1/2}}$

$$\mathbf{P}[X \geq 74.5] = \mathbf{P}\left[Z_n \geq \frac{74.5 - n \cdot 1/2}{\sqrt{n \cdot 1/2}}\right] \approx 1 - \Phi(4.9) = 4.79 \cdot 10^{-7}$$


- exact probability is $2.82 \cdot 10^{-7}$

CLT gives a much better result (but requires i.i.d.)

- **Side Note:** without continuity correction, we have 75 instead 74.5:

- This leads to $1 - \Phi(5) = 2.86 \cdot 10^{-7}$
- Issue: threshold too large ($\mathbf{P}[X \geq a] \approx \mathbf{P}[X = a]$) \Rightarrow CLT less precise
- In this region, 75 gives a better approximation than 74.5, but for smaller values (e.g., ≤ 63) the continuity corrections gives significantly better results.

A Distribution whose Average does not converge



$\text{Cau}(2, 1)$ distribution, Source: Dekking et al., Modern Introduction to Statistics

The **Cauchy distribution** has “too heavy” tails (no expectation), in particular the average does not converge.