

2005 Paper 11 Question 7

Continuous Mathematics

- (a) Let $f(x)$ be a periodic function of period 2π . Give expressions for the Fourier coefficients a_r ($r = 0, 1, \dots$) and b_r ($r = 1, 2, \dots$) of $f(x)$ where

$$\frac{a_0}{2} + \sum_{r=1}^{\infty} (a_r \cos rx + b_r \sin rx)$$

is the Fourier series representation of $f(x)$. [2 marks]

- (b) Show that the Fourier series in part (a) can also be written as a complex Fourier series

$$\sum_{r=-\infty}^{r=\infty} c_r e^{irx}$$

by deriving expressions for the complex Fourier coefficients c_r ($r = 0, \pm 1, \pm 2, \dots$) in terms of a_r and b_r . [3 marks]

- (c) Use your expressions for a_r and b_r in part (a) and for c_r in part (b) to show that

$$c_r = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-irx} dx \quad (r = 0, \pm 1, \pm 2, \dots).$$

[3 marks]

- (d) Show that the complex Fourier coefficients of $f(x - \alpha)$ (where α is a constant) are given by $e^{-ir\alpha} c_r$ ($r = 0, \pm 1, \pm 2, \dots$). [6 marks]

- (e) Suppose that $g(x)$ is another periodic function of period 2π with complex Fourier coefficients d_r ($r = 0, \pm 1, \pm 2, \dots$) and define $h(x)$ by

$$h(x) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x - y)g(y)dy.$$

Show that $h(x)$ is a periodic function of period 2π and that its complex Fourier coefficients are given by $h_r = c_r d_r$ ($r = 0, \pm 1, \pm 2, \dots$). [6 marks]

[You may assume that the periodic functions in this question satisfy the Dirichlet conditions. Euler's equation may be used without proof but should be stated precisely.]